

Fatou's Lemma and the Lebesgue's Convergence Theorem

Noboru Endou
 Gifu National College of Technology
 Japan

Keiko Narita
 Hirosaki-city
 Aomori, Japan

Yasunari Shidama
 Shinshu University
 Nagano, Japan

Summary. In this article we prove the Fatou's Lemma and Lebesgue's Convergence Theorem [10].

MML identifier: MESFUN10, version: 7.9.01 4.101.1015

The articles [15], [1], [16], [14], [11], [5], [12], [2], [3], [4], [8], [9], [13], [6], [7], and [17] provide the terminology and notation for this paper.

1. FATOU'S LEMMA

For simplicity, we adopt the following rules: X denotes a non empty set, F denotes a sequence of partial functions from X into $\overline{\mathbb{R}}$ with the same dom, s_1, s_2, s_3 denote sequences of extended reals, x denotes an element of X , a, r denote extended real numbers, and n, m, k denote natural numbers.

We now state several propositions:

- (1) If for every natural number n holds $s_2(n) \leq s_3(n)$, then $\inf \text{rng } s_2 \leq \inf \text{rng } s_3$.
- (2) Suppose that for every natural number n holds $s_2(n) \leq s_3(n)$. Then
 - (i) (the inferior real sequence of s_2)(k) $\leq$ (the inferior real sequence of s_3)(k), and
 - (ii) (the superior real sequence of s_2)(k) $\leq$ (the superior real sequence of s_3)(k).

- (3) If for every natural number n holds $s_2(n) \leq s_3(n)$, then $\liminf s_2 \leq \liminf s_3$ and $\limsup s_2 \leq \limsup s_3$.
- (4) If for every natural number n holds $s_1(n) \geq a$, then $\inf s_1 \geq a$.
- (5) If for every natural number n holds $s_1(n) \leq a$, then $\sup s_1 \leq a$.
- (6) For every element n of $\mathbb{N}$ such that $x \in \text{dom } \inf(F \uparrow n)$ holds $(\inf(F \uparrow n))(x) = \inf((F \# x) \uparrow n)$.

In the sequel S is a σ -field of subsets of X , M is a σ -measure on S , and E is an element of S .

We now state the proposition

- (7) Suppose $E = \text{dom } F(0)$ and for every n holds $F(n)$ is non-negative and $F(n)$ is measurable on E . Then there exists a sequence I of extended reals such that for every n holds $I(n) = \int F(n) dM$ and $\int \liminf F dM \leq \liminf I$.

2. LEBESGUE'S CONVERGENCE THEOREM

We now state three propositions:

- (8) For all non empty subsets X, Y of $\overline{\mathbb{R}}$ and for every extended real number r such that $X = \{r\}$ and $r \in \mathbb{R}$ holds $\sup(X + Y) = \sup X + \sup Y$.
- (9) For all non empty subsets X, Y of $\overline{\mathbb{R}}$ and for every extended real number r such that $X = \{r\}$ and $r \in \mathbb{R}$ holds $\inf(X + Y) = \inf X + \inf Y$.
- (10) If $r \in \mathbb{R}$ and for every natural number n holds $s_2(n) = r + s_3(n)$, then $\liminf s_2 = r + \liminf s_3$ and $\limsup s_2 = r + \limsup s_3$.

We follow the rules: F_1, F_2 are sequences of partial functions from X into $\overline{\mathbb{R}}$ and f, g, P are partial functions from X to $\overline{\mathbb{R}}$.

We now state several propositions:

- (11) Suppose that
 - (i) $\text{dom } F_1(0) = \text{dom } F_2(0)$,
 - (ii) F_1 has the same dom,
 - (iii) F_2 has the same dom,
 - (iv) $f^{-1}(\{+\infty\}) = \emptyset$,
 - (v) $f^{-1}(\{-\infty\}) = \emptyset$, and
 - (vi) for every natural number n holds $F_1(n) = f + F_2(n)$.

Then $\liminf F_1 = f + \liminf F_2$ and $\limsup F_1 = f + \limsup F_2$.

- (12) $s_1 \uparrow 0 = s_1$.
- (13) If f is integrable on M and g is integrable on M , then $f - g$ is integrable on M .
- (14) Suppose f is integrable on M and g is integrable on M . Then there exists an element E of S such that $E = \text{dom } f \cap \text{dom } g$ and $\int f - g dM = \int f \upharpoonright E dM + \int (-g) \upharpoonright E dM$.

- (15) If for every natural number n holds $s_2(n) = -s_3(n)$, then $\liminf s_3 = -\limsup s_2$ and $\limsup s_3 = -\liminf s_2$.
- (16) Suppose $\text{dom } F_1(0) = \text{dom } F_2(0)$ and F_1 has the same dom and F_2 has the same dom and for every natural number n holds $F_1(n) = -F_2(n)$. Then $\liminf F_1 = -\limsup F_2$ and $\limsup F_1 = -\liminf F_2$.
- (17) Suppose that
- (i) $E = \text{dom } F(0)$,
 - (ii) $E = \text{dom } P$,
 - (iii) for every natural number n holds $F(n)$ is measurable on E ,
 - (iv) P is integrable on M ,
 - (v) P is non-negative, and
 - (vi) for every element x of X and for every natural number n such that $x \in E$ holds $|F(n)|(x) \leq P(x)$.
- Then
- (vii) for every natural number n holds $|F(n)|$ is integrable on M ,
 - (viii) $|\liminf F|$ is integrable on M , and
 - (ix) $|\limsup F|$ is integrable on M .
- (18) Suppose that
- (i) $E = \text{dom } F(0)$,
 - (ii) $E = \text{dom } P$,
 - (iii) for every natural number n holds $F(n)$ is measurable on E ,
 - (iv) P is integrable on M ,
 - (v) P is non-negative, and
 - (vi) for every element x of X and for every natural number n such that $x \in E$ holds $|F(n)|(x) \leq P(x)$.
- Then there exists a sequence I of extended reals such that
- (vii) for every natural number n holds $I(n) = \int F(n) \, dM$,
 - (viii) $\liminf I \geq \int \liminf F \, dM$,
 - (ix) $\limsup I \leq \int \limsup F \, dM$, and
 - (x) if for every element x of X such that $x \in E$ holds $F \# x$ is convergent, then I is convergent and $\lim I = \int \lim F \, dM$.
- (19) Suppose that
- (i) $E = \text{dom } F(0)$,
 - (ii) for every n holds $F(n)$ is non-negative and $F(n)$ is measurable on E ,
 - (iii) for all x, n, m such that $x \in E$ and $n \leq m$ holds $F(n)(x) \geq F(m)(x)$, and
 - (iv) $\int F(0) \upharpoonright E \, dM < +\infty$.
- Then there exists a sequence I of extended reals such that for every natural number n holds $I(n) = \int F(n) \, dM$ and I is convergent and $\lim I = \int \lim F \, dM$.

Let X be a set and let F be a sequence of partial functions from X into $\overline{\mathbb{R}}$. We say that F is uniformly bounded if and only if:

- (Def. 1) There exists a real number K such that for every natural number n and for every set x such that $x \in \text{dom } F(0)$ holds $|F(n)(x)| \leq K$.

Next we state the proposition

- (20) Suppose that
- (i) $M(E) < +\infty$,
 - (ii) $E = \text{dom } F(0)$,
 - (iii) for every natural number n holds $F(n)$ is measurable on E ,
 - (iv) F is uniformly bounded, and
 - (v) for every element x of X such that $x \in E$ holds $F\#x$ is convergent.

Then

- (vi) for every natural number n holds $F(n)$ is integrable on M ,
- (vii) $\lim F$ is integrable on M , and
- (viii) there exists a sequence I of extended reals such that for every natural number n holds $I(n) = \int F(n) dM$ and I is convergent and $\lim I = \int \lim F dM$.

Let X be a set, let F be a sequence of partial functions from X into $\overline{\mathbb{R}}$, and let f be a partial function from X to $\overline{\mathbb{R}}$. We say that F is uniformly convergent to f if and only if the conditions (Def. 2) are satisfied.

- (Def. 2)(i) F has the same dom,
- (ii) $\text{dom } F(0) = \text{dom } f$, and
 - (iii) for every real number e such that $e > 0$ there exists a natural number N such that for every natural number n and for every set x such that $n \geq N$ and $x \in \text{dom } F(0)$ holds $|F(n)(x) - f(x)| < e$.

One can prove the following two propositions:

- (21) Suppose F_1 is uniformly convergent to f . Let x be an element of X . If $x \in \text{dom } F_1(0)$, then $F_1\#x$ is convergent and $\lim(F_1\#x) = f(x)$.
- (22) Suppose that
- (i) $M(E) < +\infty$,
 - (ii) $E = \text{dom } F(0)$,
 - (iii) for every natural number n holds $F(n)$ is integrable on M , and
 - (iv) F is uniformly convergent to f .

Then

- (v) f is integrable on M , and
- (vi) there exists a sequence I of extended reals such that for every natural number n holds $I(n) = \int F(n) dM$ and I is convergent and $\lim I = \int f dM$.

REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [2] Józef Białas. Infimum and supremum of the set of real numbers. Measure theory. *Formalized Mathematics*, 2(1):163–171, 1991.
- [3] Józef Białas. Series of positive real numbers. Measure theory. *Formalized Mathematics*, 2(1):173–183, 1991.
- [4] Józef Białas. The σ -additive measure theory. *Formalized Mathematics*, 2(2):263–270, 1991.
- [5] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [6] Noboru Endou and Yasunari Shidama. Integral of measurable function. *Formalized Mathematics*, 14(2):53–70, 2006.
- [7] Noboru Endou, Yasunari Shidama, and Keiko Narita. Egoroff's theorem. *Formalized Mathematics*, 16(1):57–63, 2008.
- [8] Noboru Endou, Katsumi Wasaki, and Yasunari Shidama. Basic properties of extended real numbers. *Formalized Mathematics*, 9(3):491–494, 2001.
- [9] Noboru Endou, Katsumi Wasaki, and Yasunari Shidama. Definitions and basic properties of measurable functions. *Formalized Mathematics*, 9(3):495–500, 2001.
- [10] P. R. Halmos. *Measure Theory*. Springer-Verlag, 1987.
- [11] Jarosław Kotowicz. Monotone real sequences. Subsequences. *Formalized Mathematics*, 1(3):471–475, 1990.
- [12] Andrzej Nędzusiak. σ -fields and probability. *Formalized Mathematics*, 1(2):401–407, 1990.
- [13] Beata Perkowska. Functional sequence from a domain to a domain. *Formalized Mathematics*, 3(1):17–21, 1992.
- [14] Andrzej Trybulec. Domains and their Cartesian products. *Formalized Mathematics*, 1(1):115–122, 1990.
- [15] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [16] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.
- [17] Hiroshi Yamazaki, Noboru Endou, Yasunari Shidama, and Hiroyuki Okazaki. Inferior limit, superior limit and convergence of sequences of extended real numbers. *Formalized Mathematics*, 15(4):231–236, 2007.

Received July 22, 2008
