

# Propositional Calculus for Boolean Valued Functions. Part VIII

Shunichi Kobayashi  
Matsumoto University  
Nagano

**Summary.** In this paper, we proved some elementary propositional calculus formulae for Boolean valued functions.

MML Identifier: BVFUNC26.

The articles [5], [6], [8], [7], [9], [1], [4], [3], and [2] provide the notation and terminology for this paper.

In this paper  $Y$  denotes a non empty set and  $a, b, c$  denote elements of  $Boolean^Y$ .

Let  $p, q$  be boolean-valued functions. The functor  $p \text{ 'nand' } q$  yielding a function is defined as follows:

(Def. 1)  $\text{dom}(p \text{ 'nand' } q) = \text{dom } p \cap \text{dom } q$  and for every set  $x$  such that  $x \in \text{dom}(p \text{ 'nand' } q)$  holds  $(p \text{ 'nand' } q)(x) = p(x) \text{ 'nand' } q(x)$ .

Let us observe that the functor  $p \text{ 'nand' } q$  is commutative. The functor  $p \text{ 'nor' } q$  yielding a function is defined as follows:

(Def. 2)  $\text{dom}(p \text{ 'nor' } q) = \text{dom } p \cap \text{dom } q$  and for every set  $x$  such that  $x \in \text{dom}(p \text{ 'nor' } q)$  holds  $(p \text{ 'nor' } q)(x) = p(x) \text{ 'nor' } q(x)$ .

Let us note that the functor  $p \text{ 'nor' } q$  is commutative.

Let  $p, q$  be boolean-valued functions. Note that  $p \text{ 'nand' } q$  is boolean-valued and  $p \text{ 'nor' } q$  is boolean-valued.

Let  $A$  be a non empty set and let  $p, q$  be elements of  $Boolean^A$ . Then  $p \text{ 'nand' } q$  is an element of  $Boolean^A$  and it can be characterized by the condition:

(Def. 3) For every element  $x$  of  $A$  holds  $(p \text{ 'nand' } q)(x) = p(x) \text{ 'nand' } q(x)$ .

Then  $p \text{ 'nor' } q$  is an element of  $\text{Boolean}^A$  and it can be characterized by the condition:

(Def. 4) For every element  $x$  of  $A$  holds  $(p \text{ 'nor' } q)(x) = p(x) \text{ 'nor' } q(x)$ .

Let us consider  $Y$  and let  $a, b$  be elements of  $\text{BVF}(Y)$ . Then  $a \text{ 'nand' } b$  is an element of  $\text{BVF}(Y)$ . Then  $a \text{ 'nor' } b$  is an element of  $\text{BVF}(Y)$ .

We now state a number of propositions:

- (1)  $a \text{ 'nand' } b = \neg(a \wedge b)$ .
- (2)  $a \text{ 'nor' } b = \neg(a \vee b)$ .
- (3)  $\text{true}(Y) \text{ 'nand' } a = \neg a$ .
- (4)  $\text{false}(Y) \text{ 'nand' } a = \text{true}(Y)$ .
- (5)  $\text{false}(Y) \text{ 'nand' } \text{false}(Y) = \text{true}(Y)$  and  $\text{false}(Y) \text{ 'nand' } \text{true}(Y) = \text{true}(Y)$  and  $\text{true}(Y) \text{ 'nand' } \text{true}(Y) = \text{false}(Y)$ .
- (6)  $a \text{ 'nand' } a = \neg a$  and  $\neg(a \text{ 'nand' } a) = a$ .
- (7)  $\neg(a \text{ 'nand' } b) = a \wedge b$ .
- (8)  $a \text{ 'nand' } \neg a = \text{true}(Y)$  and  $\neg(a \text{ 'nand' } \neg a) = \text{false}(Y)$ .
- (9)  $a \text{ 'nand' } b \wedge c = \neg(a \wedge b \wedge c)$ .
- (10)  $a \text{ 'nand' } b \wedge c = a \wedge b \text{ 'nand' } c$ .
- (11)  $a \text{ 'nand' } (b \vee c) = \neg(a \wedge b) \wedge \neg(a \wedge c)$ .
- (12)  $a \text{ 'nand' } (b \oplus c) = a \wedge b \Leftrightarrow a \wedge c$ .
- (13)  $a \text{ 'nand' } (b \text{ 'nand' } c) = \neg a \vee b \wedge c$  and  $a \text{ 'nand' } (b \text{ 'nand' } c) = a \Rightarrow b \wedge c$ .
- (14)  $a \text{ 'nand' } (b \text{ 'nor' } c) = \neg a \vee b \vee c$  and  $a \text{ 'nand' } (b \text{ 'nor' } c) = a \Rightarrow b \vee c$ .
- (15)  $a \text{ 'nand' } (b \Leftrightarrow c) = a \Rightarrow b \oplus c$ .
- (16)  $a \text{ 'nand' } a \wedge b = a \text{ 'nand' } b$ .
- (17)  $a \text{ 'nand' } (a \vee b) = \neg a \wedge \neg(a \wedge b)$ .
- (18)  $a \text{ 'nand' } (a \Leftrightarrow b) = a \Rightarrow a \oplus b$ .
- (19)  $a \text{ 'nand' } (a \text{ 'nand' } b) = \neg a \vee b$  and  $a \text{ 'nand' } (a \text{ 'nand' } b) = a \Rightarrow b$ .
- (20)  $a \text{ 'nand' } (a \text{ 'nor' } b) = \text{true}(Y)$ .
- (21)  $a \text{ 'nand' } (a \Leftrightarrow b) = \neg a \vee \neg b$ .
- (22)  $a \wedge b = a \text{ 'nand' } b \text{ 'nand' } (a \text{ 'nand' } b)$ .
- (23)  $a \text{ 'nand' } b \text{ 'nand' } (a \text{ 'nand' } c) = a \wedge (b \vee c)$ .
- (24)  $a \text{ 'nand' } (b \Rightarrow c) = (\neg a \vee b) \wedge \neg(a \wedge c)$ .
- (25)  $a \text{ 'nand' } (a \Rightarrow b) = \neg(a \wedge b)$ .
- (26)  $\text{true}(Y) \text{ 'nor' } a = \text{false}(Y)$ .
- (27)  $\text{false}(Y) \text{ 'nor' } a = \neg a$ .
- (28)  $\text{false}(Y) \text{ 'nor' } \text{false}(Y) = \text{true}(Y)$  and  $\text{false}(Y) \text{ 'nor' } \text{true}(Y) = \text{false}(Y)$  and  $\text{true}(Y) \text{ 'nor' } \text{true}(Y) = \text{false}(Y)$ .
- (29)  $a \text{ 'nor' } a = \neg a$  and  $\neg(a \text{ 'nor' } a) = a$ .

- (30)  $\neg(a \text{ 'nor' } b) = a \vee b.$
- (31)  $a \text{ 'nor' } \neg a = \text{false}(Y)$  and  $\neg(a \text{ 'nor' } \neg a) = \text{true}(Y).$
- (32)  $\neg a \wedge (a \oplus b) = \neg a \wedge b.$
- (33)  $a \text{ 'nor' } b \wedge c = \neg(a \vee b) \vee \neg(a \vee c).$
- (34)  $a \text{ 'nor' } (b \vee c) = \neg(a \vee b \vee c).$
- (35)  $a \text{ 'nor' } (b \Leftrightarrow c) = \neg a \wedge (b \oplus c).$
- (36)  $a \text{ 'nor' } (b \Rightarrow c) = \neg a \wedge b \wedge \neg c.$
- (37)  $a \text{ 'nor' } (b \text{ 'nand' } c) = \neg a \wedge b \wedge c.$
- (38)  $a \text{ 'nor' } (b \text{ 'nor' } c) = \neg a \wedge (b \vee c).$
- (39)  $a \text{ 'nor' } a \wedge b = \neg(a \wedge (a \vee b)).$
- (40)  $a \text{ 'nor' } (a \vee b) = \neg(a \vee b).$
- (41)  $a \text{ 'nor' } (a \Leftrightarrow b) = \neg a \wedge b.$
- (42)  $a \text{ 'nor' } (a \Rightarrow b) = \text{false}(Y).$
- (43)  $a \text{ 'nor' } (a \text{ 'nand' } b) = \text{false}(Y).$
- (44)  $a \text{ 'nor' } (a \text{ 'nor' } b) = \neg a \wedge b.$
- (45)  $\text{false}(Y) \Leftrightarrow \text{false}(Y) = \text{true}(Y).$
- (46)  $\text{false}(Y) \Leftrightarrow \text{true}(Y) = \text{false}(Y).$
- (47)  $\text{true}(Y) \Leftrightarrow \text{true}(Y) = \text{true}(Y).$
- (48)  $a \Leftrightarrow a = \text{true}(Y)$  and  $\neg(a \Leftrightarrow a) = \text{false}(Y).$
- (49)  $a \Leftrightarrow a \vee b = a \vee \neg b.$
- (50)  $a \wedge (b \text{ 'nand' } c) = a \wedge \neg b \vee a \wedge \neg c.$
- (51)  $a \vee (b \text{ 'nand' } c) = a \vee \neg b \vee \neg c.$
- (52)  $a \oplus (b \text{ 'nand' } c) = \neg a \wedge \neg(b \wedge c) \vee a \wedge b \wedge c.$
- (53)  $a \Leftrightarrow b \text{ 'nand' } c = a \wedge \neg(b \wedge c) \vee \neg a \wedge b \wedge c.$
- (54)  $a \Rightarrow b \text{ 'nand' } c = \neg(a \wedge b \wedge c).$
- (55)  $a \text{ 'nor' } (b \text{ 'nand' } c) = \neg(a \vee \neg b \vee \neg c).$
- (56)  $a \wedge (a \text{ 'nand' } b) = a \wedge \neg b.$
- (57)  $a \vee (a \text{ 'nand' } b) = \text{true}(Y).$
- (58)  $a \oplus (a \text{ 'nand' } b) = \neg a \vee b.$
- (59)  $a \Leftrightarrow a \text{ 'nand' } b = a \wedge \neg b.$
- (60)  $a \Rightarrow a \text{ 'nand' } b = \neg(a \wedge b).$
- (61)  $a \text{ 'nor' } (a \text{ 'nand' } b) = \text{false}(Y).$
- (62)  $a \wedge (b \text{ 'nor' } c) = a \wedge \neg b \wedge \neg c.$
- (63)  $a \vee (b \text{ 'nor' } c) = (a \vee \neg b) \wedge (a \vee \neg c).$
- (64)  $a \oplus (b \text{ 'nor' } c) = (a \vee \neg(b \vee c)) \wedge (\neg a \vee b \vee c).$
- (65)  $a \Leftrightarrow b \text{ 'nor' } c = (a \vee b \vee c) \wedge (\neg a \vee \neg(b \vee c)).$
- (66)  $a \Rightarrow b \text{ 'nor' } c = \neg(a \wedge (b \vee c)).$

$$(67) \quad a \text{ 'nand' } (b \text{ 'nor' } c) = \neg a \vee b \vee c.$$

$$(68) \quad a \wedge (a \text{ 'nor' } b) = \text{false}(Y).$$

$$(69) \quad a \vee (a \text{ 'nor' } b) = a \vee \neg b.$$

$$(70) \quad a \oplus (a \text{ 'nor' } b) = a \vee \neg b.$$

$$(71) \quad a \Leftrightarrow a \text{ 'nor' } b = \neg a \wedge b.$$

$$(72) \quad a \Rightarrow a \text{ 'nor' } b = \neg(a \vee a \wedge b).$$

$$(73) \quad a \text{ 'nand' } (a \text{ 'nor' } b) = \text{true}(Y).$$

#### ACKNOWLEDGMENTS

This research was partially supported by the research funds of the Matsumoto University.

#### REFERENCES

- [1] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [2] Shunichi Kobayashi. On the calculus of binary arithmetics. *Formalized Mathematics*, 11(4):417–419, 2003.
- [3] Shunichi Kobayashi and Kui Jia. A theory of Boolean valued functions and partitions. *Formalized Mathematics*, 7(2):249–254, 1998.
- [4] Andrzej Trybulec. Function domains and Frænkel operator. *Formalized Mathematics*, 1(3):495–500, 1990.
- [5] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [6] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [7] Edmund Woronowicz. Interpretation and satisfiability in the first order logic. *Formalized Mathematics*, 1(4):739–743, 1990.
- [8] Edmund Woronowicz. Many-argument relations. *Formalized Mathematics*, 1(4):733–737, 1990.
- [9] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.

*Received September 5, 2004*

---