

Laplace Expansion

Karol Pąk
 Institute of Computer Science
 University of Białystok
 Poland

Andrzej Trybulec
 Institute of Computer Science
 University of Białystok
 Poland

Summary. In the article the formula for Laplace expansion is proved.

MML identifier: LAPLACE, version: 7.8.05 4.87.985

The notation and terminology used in this paper are introduced in the following articles: [23], [11], [29], [20], [12], [30], [31], [6], [9], [7], [3], [4], [21], [28], [26], [15], [22], [10], [5], [13], [24], [14], [33], [25], [18], [34], [1], [8], [2], [16], [17], [27], [19], and [32].

1. PRELIMINARIES

For simplicity, we follow the rules: x, y are sets, N is an element of $\mathbb{N}$, c , i, j, k, m, n are natural numbers, D is a non empty set, s is an element of $2\text{SetSeg}(n+2)$, p is an element of the permutations of n -element set, p_1, q_1 are elements of the permutations of $(n+1)$ -element set, p_2 is an element of the permutations of $(n+2)$ -element set, K is a field, a, b are elements of K , f is a finite sequence of elements of K , A is a matrix over K , A_1 is a matrix over D of dimension $n \times m$, p_3 is a finite sequence of elements of D , and M is a matrix over K of dimension n .

The following propositions are true:

- (1) For every finite sequence f and for every natural number i such that $i \in \text{dom } f$ holds $\text{len}(f_{\setminus i}) = \text{len } f - 1$.
- (2) Let i, j, n be natural numbers and M be a matrix over K of dimension n . If $i \in \text{dom } M$, then $\text{len}(\text{the deleting of } i\text{-row and } j\text{-column in } M) = n - 1$.
- (3) If $j \in \text{Seg width } A$, then $\text{width}(\text{the deleting of } j\text{-column in } A) = \text{width } A - 1$.

- (4) For every natural number i such that $\text{len } A > 1$ holds $\text{width } A = \text{width (the deleting of } i\text{-row in } A)$.
- (5) For every natural number i such that $j \in \text{Seg width } M$ holds $\text{width (the deleting of } i\text{-row and } j\text{-column in } M) = n -' 1$.

Let G be a non empty groupoid, let B be a function from $\{ \text{the carrier of } G, \mathbb{N} \}$ into the carrier of G , let g be an element of G , and let i be a natural number. Then $B(g, i)$ is an element of G .

One can prove the following propositions:

- (6) $\overline{\text{the permutations of } n\text{-element set}} = n!$.
- (7) $\overline{\text{For all } i, j \text{ such that } i \in \text{Seg}(n+1) \text{ and } j \in \text{Seg}(n+1) \text{ holds } \{p_1 : p_1(i) = j\}} = n!$.
- (8) Let K be a Fanoian field, given p_2 , and X, Y be elements of $\text{Fin } 2\text{Set Seg}(n+2)$. Suppose $Y = \{s : s \in X \wedge (\text{Part-sgn}(p_2, K))(s) = -\mathbf{1}_K\}$. Then $(\text{the multiplication of } K) \cdot \sum_X \text{Part-sgn}(p_2, K) = \text{power}_K(-\mathbf{1}_K, \text{card } Y)$.
- (9) Let K be a Fanoian field and given p_2, i, j . Suppose $i \in \text{Seg}(n+2)$ and $p_2(i) = j$. Then there exists an element X of $\text{Fin } 2\text{Set Seg}(n+2)$ such that $X = \{\{N, i\} : \{N, i\} \in 2\text{Set Seg}(n+2)\}$ and $(\text{the multiplication of } K) \cdot \sum_X \text{Part-sgn}(p_2, K) = \text{power}_K(-\mathbf{1}_K, i+j)$.
- (10) Let given i, j . Suppose $i \in \text{Seg}(n+1)$ and $j \in \text{Seg}(n+1)$ and $n \geq 2$. Then there exists a function P_1 from $2\text{Set Seg } n$ into $2\text{Set Seg}(n+1)$ such that
 - (i) $\text{rng } P_1 = 2\text{Set Seg}(n+1) \setminus \{\{N, i\} : \{N, i\} \in 2\text{Set Seg}(n+1)\}$,
 - (ii) P_1 is one-to-one, and
 - (iii) for all k, m such that $k < m$ and $\{k, m\} \in 2\text{Set Seg } n$ holds if $m < i$ and $k < i$, then $P_1(\{k, m\}) = \{k, m\}$ and if $m \geq i$ and $k < i$, then $P_1(\{k, m\}) = \{k, m+1\}$ and if $m \geq i$ and $k \geq i$, then $P_1(\{k, m\}) = \{k+1, m+1\}$.
- (11) If $n < 2$, then for every element p of the permutations of n -element set holds p is even and $p = \text{idseq}(n)$.
- (12) Let X, Y, D be non empty sets, f be a function from X into $\text{Fin } Y$, g be a function from $\text{Fin } Y$ into D , and F be a binary operation on D . Suppose that
 - (i) for all elements A, B of $\text{Fin } Y$ such that A misses B holds $F(g(A), g(B)) = g(A \cup B)$,
 - (ii) F is commutative and associative and has a unity, and
 - (iii) $g(\emptyset) = \mathbf{1}_F$.

Let I be an element of $\text{Fin } X$. Suppose that for all x, y such that $x \in I$ and $y \in I$ and $f(x)$ meets $f(y)$ holds $x = y$. Then $F \cdot \sum_I g \cdot f = F \cdot \sum_{f \circ I} g$ and $F \cdot \sum_{f \circ I} g = g(\bigcup(f \circ I))$ and $\bigcup(f \circ I)$ is an element of $\text{Fin } Y$.

2. AUXILIARY NOTIONS

Let i, j, n be natural numbers, let us consider K , and let M be a matrix over K of dimension n . Let us assume that $i \in \text{Seg } n$ and $j \in \text{Seg } n$. The functor $\text{Delete}(M, i, j)$ yielding a matrix over K of dimension $n - 1$ is defined as follows:

(Def. 1) $\text{Delete}(M, i, j)$ = the deleting of i -row and j -column in M .

The following propositions are true:

- (13) Let given i, j . Suppose $i \in \text{Seg } n$ and $j \in \text{Seg } n$. Let given k, m such that $k \in \text{Seg}(n - 1)$ and $m \in \text{Seg}(n - 1)$. Then
 - (i) if $k < i$ and $m < j$, then $(\text{Delete}(M, i, j))_{k,m} = M_{k,m}$,
 - (ii) if $k < i$ and $m \geq j$, then $(\text{Delete}(M, i, j))_{k,m} = M_{k,m+1}$,
 - (iii) if $k \geq i$ and $m < j$, then $(\text{Delete}(M, i, j))_{k,m} = M_{k+1,m}$, and
 - (iv) if $k \geq i$ and $m \geq j$, then $(\text{Delete}(M, i, j))_{k,m} = M_{k+1,m+1}$.
- (14) For all i, j such that $i \in \text{Seg } n$ and $j \in \text{Seg } n$ holds $(\text{Delete}(M, i, j))^T = \text{Delete}(M^T, j, i)$.
- (15) For every finite sequence f of elements of K and for all i, j such that $i \in \text{Seg } n$ and $j \in \text{Seg } n$ holds $\text{Delete}(M, i, j) = \text{Delete}(\text{RLine}(M, i, f), i, j)$.

Let us consider c, n, m, D , let M be a matrix over D of dimension $n \times m$, and let p_3 be a finite sequence of elements of D . The functor $\text{ReplaceCol}(M, c, p_3)$ yielding a matrix over D of dimension $n \times m$ is defined by:

- (Def. 2)(i) $\text{len } \text{ReplaceCol}(M, c, p_3) = \text{len } M$ and $\text{width } \text{ReplaceCol}(M, c, p_3) = \text{width } M$ and for all i, j such that $\langle i, j \rangle \in$ the indices of M holds if $j \neq c$, then $(\text{ReplaceCol}(M, c, p_3))_{i,j} = M_{i,j}$ and if $j = c$, then $(\text{ReplaceCol}(M, c, p_3))_{i,c} = p_3(i)$ if $\text{len } p_3 = \text{len } M$,
- (ii) $\text{ReplaceCol}(M, c, p_3) = M$, otherwise.

Let us consider c, n, m, D , let M be a matrix over D of dimension $n \times m$, and let p_3 be a finite sequence of elements of D . We introduce $\text{RCol}(M, c, p_3)$ as a synonym of $\text{ReplaceCol}(M, c, p_3)$.

We now state four propositions:

- (16) For every i such that $i \in \text{Seg width } A_1$ holds if $i = c$ and $\text{len } p_3 = \text{len } A_1$, then $(\text{RCol}(A_1, c, p_3))_{\square, i} = p_3$ and if $i \neq c$, then $(\text{RCol}(A_1, c, p_3))_{\square, i} = (A_1)_{\square, i}$.
- (17) If $c \notin \text{Seg width } A_1$, then $\text{RCol}(A_1, c, p_3) = A_1$.
- (18) $\text{RCol}(A_1, c, (A_1)_{\square, c}) = A_1$.
- (19) Let A be a matrix over D of dimension $n \times m$ and A' be a matrix over D of dimension $m \times n$. If $A' = A^T$ and if $m = 0$, then $n = 0$, then $\text{ReplaceCol}(A, c, p_3) = (\text{ReplaceLine}(A', c, p_3))^T$.

3. PERMUTATIONS

Let us consider i, n and let p_4 be an element of the permutations of $(n+1)$ -element set. Let us assume that $i \in \text{Seg}(n+1)$. The functor $\text{Rem}(p_4, i)$ yielding an element of the permutations of n -element set is defined by the condition (Def. 3).

(Def. 3) Let given k such that $k \in \text{Seg } n$. Then

- (i) if $k < i$, then if $p_4(k) < p_4(i)$, then $(\text{Rem}(p_4, i))(k) = p_4(k)$ and if $p_4(k) \geq p_4(i)$, then $(\text{Rem}(p_4, i))(k) = p_4(k) - 1$, and
- (ii) if $k \geq i$, then if $p_4(k+1) < p_4(i)$, then $(\text{Rem}(p_4, i))(k) = p_4(k+1)$ and if $p_4(k+1) \geq p_4(i)$, then $(\text{Rem}(p_4, i))(k) = p_4(k+1) - 1$.

One can prove the following three propositions:

- (20) Let given i, j . Suppose $i \in \text{Seg}(n+1)$ and $j \in \text{Seg}(n+1)$. Let P be a set. Suppose $P = \{p_1 : p_1(i) = j\}$. Then there exists a function P_1 from P into the permutations of n -element set such that P_1 is bijective and for every q_1 such that $q_1(i) = j$ holds $P_1(q_1) = \text{Rem}(q_1, i)$.
- (21) For all i, j such that $i \in \text{Seg}(n+1)$ and $p_1(i) = j$ holds $(-1)^{\text{sgn}(p_1)} a = \text{power}_K(-\mathbf{1}_K, i+j) \cdot (-1)^{\text{sgn}(\text{Rem}(p_1, i))} a$.
- (22) Let given i, j . Suppose $i \in \text{Seg}(n+1)$ and $p_1(i) = j$. Let M be a matrix over K of dimension $n+1$ and D_1 be a matrix over K of dimension n . Suppose $D_1 = \text{Delete}(M, i, j)$. Then (the product on paths of M)(p_1) = $\text{power}_K(-\mathbf{1}_K, i+j) \cdot M_{i,j} \cdot (\text{the product on paths of } D_1)(\text{Rem}(p_1, i))$.

4. MINORS AND COFACTORS

Let i, j, n be natural numbers, let us consider K , and let M be a matrix over K of dimension n . The functor $\text{Minor}(M, i, j)$ yielding an element of K is defined by:

(Def. 4) $\text{Minor}(M, i, j) = \text{Det Delete}(M, i, j)$.

Let i, j, n be natural numbers, let us consider K , and let M be a matrix over K of dimension n . The functor $\text{Cofactor}(M, i, j)$ yielding an element of K is defined as follows:

(Def. 5) $\text{Cofactor}(M, i, j) = \text{power}_K(-\mathbf{1}_K, i+j) \cdot \text{Minor}(M, i, j)$.

The following propositions are true:

- (23) Let given i, j . Suppose $i \in \text{Seg } n$ and $j \in \text{Seg } n$. Let P be an element of Fin (the permutations of n -element set). Suppose $P = \{p : p(i) = j\}$. Let M be a matrix over K of dimension n . Then (the addition of K)- $\sum_P$ (the product on paths of M) = $M_{i,j} \cdot \text{Cofactor}(M, i, j)$.
- (24) For all i, j such that $i \in \text{Seg } n$ and $j \in \text{Seg } n$ holds $\text{Minor}(M, i, j) = \text{Minor}(M^T, j, i)$.

Let us consider n, K and let M be a matrix over K of dimension n . The matrix of cofactor M yielding a matrix over K of dimension n is defined by the condition (Def. 6).

- (Def. 6) Let i, j be natural numbers. Suppose $\langle i, j \rangle \in$ the indices of the matrix of cofactor M . Then (the matrix of cofactor M) $_{i,j} = \text{Cofactor}(M, i, j)$.

5. LAPLACE EXPANSION

Let us consider n, i, K and let M be a matrix over K of dimension n . The functor $\text{LaplaceExpL}(M, i)$ yields a finite sequence of elements of K and is defined as follows:

- (Def. 7) $\text{len LaplaceExpL}(M, i) = n$ and for every j such that $j \in \text{dom LaplaceExpL}(M, i)$ holds
 $(\text{LaplaceExpL}(M, i))(j) = M_{i,j} \cdot \text{Cofactor}(M, i, j)$.

Let us consider n , let j be a natural number, let us consider K , and let M be a matrix over K of dimension n . The functor $\text{LaplaceExpC}(M, j)$ yields a finite sequence of elements of K and is defined by:

- (Def. 8) $\text{len LaplaceExpC}(M, j) = n$ and for every i such that $i \in \text{dom LaplaceExpC}(M, j)$ holds $(\text{LaplaceExpC}(M, j))(i) = M_{i,j} \cdot \text{Cofactor}(M, i, j)$.

One can prove the following propositions:

- (25) For every natural number i and for every matrix M over K of dimension n such that $i \in \text{Seg } n$ holds $\text{Det } M = \sum \text{LaplaceExpL}(M, i)$.
- (26) For every i such that $i \in \text{Seg } n$ holds $\text{LaplaceExpC}(M, i) = \text{LaplaceExpL}(M^T, i)$.
- (27) For every natural number j and for every matrix M over K of dimension n such that $j \in \text{Seg } n$ holds $\text{Det } M = \sum \text{LaplaceExpC}(M, j)$.
- (28) For all p, i such that $\text{len } f = n$ and $i \in \text{Seg } n$ holds $\text{Line}(\text{the matrix of cofactor } M, i) \bullet f = \text{LaplaceExpL}(\text{RLine}(M, i, f), i)$.
- (29) If $i \in \text{Seg } n$, then $\text{Line}(M, j) \cdot ((\text{the matrix of cofactor } M)^T)_{\square, i} = \text{Det RLine}(M, i, \text{Line}(M, j))$.
- (30) If $\text{Det } M \neq 0_K$, then $M \cdot (\text{Det } M^{-1} \cdot (\text{the matrix of cofactor } M)^T) = \begin{pmatrix} 1 & & 0 \\ & \ddots & \\ 0 & & 1 \end{pmatrix}_{n \times n}^K$.
- (31) For all f, i such that $\text{len } f = n$ and $i \in \text{Seg } n$ holds $(\text{the matrix of cofactor } M)_{\square, i} \bullet f = \text{LaplaceExpL}(\text{RLine}(M^T, i, f), i)$.
- (32) If $i \in \text{Seg } n$ and $j \in \text{Seg } n$, then $\text{Line}((\text{the matrix of cofactor } M)^T, i) \cdot M_{\square, j} = \text{Det RLine}(M^T, i, \text{Line}(M^T, j))$.

- (33) If $\text{Det } M \neq 0_K$, then $\text{Det } M^{-1} \cdot (\text{the matrix of cofactor } M)^T \cdot M =$

$$\begin{pmatrix} 1 & & 0 \\ & \ddots & \\ 0 & & 1 \end{pmatrix}_{n \times n}^K.$$
- (34) M is invertible iff $\text{Det } M \neq 0_K$.
- (35) If $\text{Det } M \neq 0_K$, then $M^\smile = \text{Det } M^{-1} \cdot (\text{the matrix of cofactor } M)^T$.
- (36) Let M be a matrix over K of dimension n . Suppose M is invertible. Let given i, j . If $\langle i, j \rangle \in$ the indices of $M^\smile$, then $M^\smile_{i,j} = \text{Det } M^{-1} \cdot \text{power}_K(-\mathbf{1}_K, i + j) \cdot \text{Minor}(M, j, i)$.
- (37) Let A be a matrix over K of dimension n . Suppose $\text{Det } A \neq 0_K$. Let x, b be matrices over K . Suppose $\text{len } x = n$ and $A \cdot x = b$. Then $x = A^\smile \cdot b$ and for all i, j such that $\langle i, j \rangle \in$ the indices of x holds $x_{i,j} = \text{Det } A^{-1} \cdot \text{Det ReplaceCol}(A, i, b_{\square, j})$.
- (38) Let A be a matrix over K of dimension n . Suppose $\text{Det } A \neq 0_K$. Let x, b be matrices over K . Suppose $\text{width } x = n$ and $x \cdot A = b$. Then $x = b \cdot A^\smile$ and for all i, j such that $\langle i, j \rangle \in$ the indices of x holds $x_{i,j} = \text{Det } A^{-1} \cdot \text{Det ReplaceLine}(A, j, \text{Line}(b, i))$.

6. PRODUCT BY A VECTOR

Let D be a non empty set and let f be a finite sequence of elements of D . Then $\langle f \rangle$ is a matrix over D of dimension $1 \times \text{len } f$.

Let us consider K , let M be a matrix over K , and let f be a finite sequence of elements of K . The functor $M \cdot f$ yielding a matrix over K is defined by:

$$(\text{Def. 9}) \quad M \cdot f = M \cdot \langle f \rangle^T.$$

The functor $f \cdot M$ yields a matrix over K and is defined by:

$$(\text{Def. 10}) \quad f \cdot M = \langle f \rangle \cdot M.$$

Next we state two propositions:

- (39) Let A be a matrix over K of dimension n . Suppose $\text{Det } A \neq 0_K$. Let x, b be finite sequences of elements of K . Suppose $\text{len } x = n$ and $A \cdot x = \langle b \rangle^T$. Then $\langle x \rangle^T = A^\smile \cdot b$ and for every i such that $i \in \text{Seg } n$ holds $x(i) = \text{Det } A^{-1} \cdot \text{Det ReplaceCol}(A, i, b)$.
- (40) Let A be a matrix over K of dimension n . Suppose $\text{Det } A \neq 0_K$. Let x, b be finite sequences of elements of K . Suppose $\text{len } x = n$ and $x \cdot A = \langle b \rangle$. Then $\langle x \rangle = b \cdot A^\smile$ and for every i such that $i \in \text{Seg } n$ holds $x(i) = \text{Det } A^{-1} \cdot \text{Det ReplaceLine}(A, i, b)$.

REFERENCES

- [1] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(2):377–382, 1990.
- [2] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [4] Czesław Byliński. Binary operations. *Formalized Mathematics*, 1(1):175–180, 1990.
- [5] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(3):529–536, 1990.
- [6] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [7] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [8] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Formalized Mathematics*, 1(3):521–527, 1990.
- [9] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [10] Czesław Byliński. Semigroup operations on finite subsets. *Formalized Mathematics*, 1(4):651–656, 1990.
- [11] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [12] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(1):165–167, 1990.
- [13] Katarzyna Jankowska. Matrices. Abelian group of matrices. *Formalized Mathematics*, 2(4):475–480, 1991.
- [14] Katarzyna Jankowska. Transpose matrices and groups of permutations. *Formalized Mathematics*, 2(5):711–717, 1991.
- [15] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Formalized Mathematics*, 1(2):335–342, 1990.
- [16] Rafał Kwiatek. Factorial and Newton coefficients. *Formalized Mathematics*, 1(5):887–890, 1990.
- [17] Yatsuka Nakamura. Determinant of some matrices of field elements. *Formalized Mathematics*, 14(1):1–5, 2006.
- [18] Takaya Nishiyama and Yasuho Mizuhara. Binary arithmetics. *Formalized Mathematics*, 4(1):83–86, 1993.
- [19] Karol Pąk. Basic properties of determinants of square matrices over a field. *Formalized Mathematics*, 15(1):17–25, 2007.
- [20] Andrzej Trybulec. Subsets of complex numbers. *To appear in Formalized Mathematics*.
- [21] Andrzej Trybulec. Binary operations applied to functions. *Formalized Mathematics*, 1(2):329–334, 1990.
- [22] Andrzej Trybulec. Semilattice operations on finite subsets. *Formalized Mathematics*, 1(2):369–376, 1990.
- [23] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [24] Andrzej Trybulec and Agata Darmochwał. Boolean domains. *Formalized Mathematics*, 1(1):187–190, 1990.
- [25] Wojciech A. Trybulec. Binary operations on finite sequences. *Formalized Mathematics*, 1(5):979–981, 1990.
- [26] Wojciech A. Trybulec. Groups. *Formalized Mathematics*, 1(5):821–827, 1990.
- [27] Wojciech A. Trybulec. Non-contiguous substrings and one-to-one finite sequences. *Formalized Mathematics*, 1(3):569–573, 1990.
- [28] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(2):291–296, 1990.
- [29] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [30] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [31] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.
- [32] Xiaopeng Yue, Xiquan Liang, and Zhongpin Sun. Some properties of some special matrices. *Formalized Mathematics*, 13(4):541–547, 2005.
- [33] Katarzyna Zawadzka. The sum and product of finite sequences of elements of a field. *Formalized Mathematics*, 3(2):205–211, 1992.

- [34] Katarzyna Zawadzka. The product and the determinant of matrices with entries in a field. *Formalized Mathematics*, 4(**1**):1–8, 1993.

Received August 13, 2007
