FORMALIZED MATHEMATICS
2007, Vol. **15**, No. **4**, Pages 167–170
DOI: 10.2478/v10037-007-0019-2

Alexandroff One Point Compactification

Czesław Byliński
University of Białystok
Poland

Summary. In the article, I introduce the notions of the compactification of topological spaces and the Alexandroff one point compactification. Some properties of the locally compact spaces and one point compactification are proved.

MML identifier: `COMPACT1`, version: `7.8.05 4.87.985`

The articles [15], [5], [16], [17], [4], [18], [1], [8], [14], [13], [19], [7], [9], [10], [6], [12], [2], [3], and [11] provide the notation and terminology for this paper.

Let X be a topological space and let P be a family of subsets of X. We say that P is compact if and only if:

(Def. 1) For every subset U of X such that $U \in P$ holds U is compact.

Let X be a topological space and let U be a subset of X. We say that U is relatively-compact if and only if:

(Def. 2) $\overline{U}$ is compact.

Let X be a topological space. Note that $\emptyset_X$ is relatively-compact.

Let X be a topological space. Observe that there exists a subset of X which is relatively-compact.

Let X be a topological space and let U be a relatively-compact subset of X. Observe that $\overline{U}$ is compact.

Let X be a topological space and let U be a subset of X. We introduce U is pre-compact as a synonym of U is relatively-compact.

Let X be a non empty topological space. We introduce X is liminally-compact as a synonym of X is locally-compact.

Let X be a non empty topological space. Let us observe that X is liminally-compact if and only if:

(Def. 3) For every point x of X holds there exists a generalized basis of x which is compact.

© 2007 University of Białystok
ISSN 1426–2630

Let X be a non empty topological space. We say that X is locally-relatively-compact if and only if:

(Def. 4) For every point x of X holds there exists a neighbourhood of x which is relatively-compact.

Let X be a non empty topological space. We say that X is locally-closed/compact if and only if:

(Def. 5) For every point x of X holds there exists a neighbourhood of x which is closed and compact.

Let X be a non empty topological space. We say that X is locally-compact if and only if:

(Def. 6) For every point x of X holds there exists a neighbourhood of x which is compact.

Let us observe that every non empty topological space which is liminally-compact is also locally-compact.

Let us note that every non empty T_3 topological space which is locally-compact is also liminally-compact.

One can verify that every non empty topological space which is locally-relatively-compact is also locally-closed/compact.

Let us observe that every non empty topological space which is locally-closed/compact is also locally-relatively-compact.

Let us observe that every non empty topological space which is locally-relatively-compact is also locally-compact.

One can verify that every non empty Hausdorff topological space which is locally-compact is also locally-relatively-compact.

One can check that every non empty topological space which is compact is also locally-compact.

Let us observe that every non empty topological space which is discrete is also locally-compact.

Let us mention that there exists a topological space which is discrete and non empty.

Let X be a locally-compact non empty topological space and let C be a closed non empty subset of X. Note that $X\restriction C$ is locally-compact.

Let X be a locally-compact non empty T_3 topological space and let P be an open non empty subset of X. Note that $X\restriction P$ is locally-compact.

One can prove the following two propositions:

(1) Let X be a Hausdorff non empty topological space and E be a non empty subset of X. If $X\restriction E$ is dense and locally-compact, then $X\restriction E$ is open.

(2) For all topological spaces X, Y and for every subset A of X such that $\Omega_X \subseteq \Omega_Y$ holds $(\mathrm{incl}(X,Y))^\circ A = A$.

Let X, Y be topological spaces and let f be a function from X into Y. We say that f is embedding if and only if:

(Def. 7) There exists a function h from X into $Y \restriction \operatorname{rng} f$ such that $h = f$ and h is a homeomorphism.

The following proposition is true

(3) Let X, Y be topological spaces. Suppose $\Omega_X \subseteq \Omega_Y$ and there exists a subset X_1 of Y such that $X_1 = \Omega_X$ and the topology of $Y \restriction X_1 =$ the topology of X. Then $\operatorname{incl}(X, Y)$ is embedding.

Let X be a topological space, let Y be a topological space, and let h be a function from X into Y. We say that h is compactification if and only if:

(Def. 8) h is embedding and Y is compact and $h^\circ(\Omega_X)$ is dense.

Let X be a topological space and let Y be a topological space. Note that every function from X into Y which is compactification is also embedding.

Let X be a topological structure. The one-point compactification of X yields a strict topological structure and is defined by the conditions (Def. 9).

(Def. 9)(i) The carrier of the one-point compactification of $X = \operatorname{succ}(\Omega_X)$, and

(ii) the topology of the one-point compactification of $X =$ (the topology of $X) \cup \{U \cup \{\Omega_X\}; U$ ranges over subsets of X: U is open $\wedge\, U^{\mathrm{c}}$ is compact$\}$.

Let X be a topological structure. Note that the one-point compactification of X is non empty.

We now state the proposition

(4) For every topological structure X holds $\Omega_X \subseteq \Omega_{\text{the one-point compactification of } X}$.

Let X be a topological space. Note that the one-point compactification of X is topological space-like.

Next we state the proposition

(5) Every topological structure X is a subspace of the one-point compactification of X.

Let X be a topological space. One can verify that the one-point compactification of X is compact.

One can prove the following propositions:

(6) Let X be a non empty topological space. Then X is Hausdorff and locally-compact if and only if the one-point compactification of X is Hausdorff.

(7) Let X be a non empty topological space. Then X is non compact if and only if there exists a subset X' of the one-point compactification of X such that $X' = \Omega_X$ and X' is dense.

(8) Let X be a non empty topological space. Suppose X is non compact. Then $\operatorname{incl}(X,$ the one-point compactification of $X)$ is compactification.

References

[1] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(**1**):91–96, 1990.
[2] Grzegorz Bancerek. The "way-below" relation. *Formalized Mathematics*, 6(**1**):169–176, 1997.
[3] Grzegorz Bancerek. Bases and refinements of topologies. *Formalized Mathematics*, 7(**1**):35–43, 1998.
[4] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(**1**):55–65, 1990.
[5] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(**1**):47–53, 1990.
[6] Agata Darmochwał. Compact spaces. *Formalized Mathematics*, 1(**2**):383–386, 1990.
[7] Agata Darmochwał. Families of subsets, subspaces and mappings in topological spaces. *Formalized Mathematics*, 1(**2**):257–261, 1990.
[8] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(**1**):165–167, 1990.
[9] Zbigniew Karno. Separated and weakly separated subspaces of topological spaces. *Formalized Mathematics*, 2(**5**):665–674, 1991.
[10] Zbigniew Karno. On nowhere and everywhere dense subspaces of topological spaces. *Formalized Mathematics*, 4(**1**):137–146, 1993.
[11] Artur Korniłowicz. Introduction to meet-continuous topological lattices. *Formalized Mathematics*, 7(**2**):279–283, 1998.
[12] Beata Padlewska. Locally connected spaces. *Formalized Mathematics*, 2(**1**):93–96, 1991.
[13] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(**1**):223–230, 1990.
[14] Andrzej Trybulec. Domains and their Cartesian products. *Formalized Mathematics*, 1(**1**):115–122, 1990.
[15] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(**1**):9–11, 1990.
[16] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(**1**):67–71, 1990.
[17] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(**1**):73–83, 1990.
[18] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(**1**):181–186, 1990.
[19] Mirosław Wysocki and Agata Darmochwał. Subsets of topological spaces. *Formalized Mathematics*, 1(**1**):231–237, 1990.

Received August 13, 2007