FORMALIZED MATHEMATICS
2007, Vol. **15**, No. **4**, Pages 171–174
DOI: 10.2478/v10037-007-0020-9

Arrow's Impossibility Theorem

Freek Wiedijk
Institute for Computing and Information Sciences
Radboud University Nijmegen
Toernooiveld 1, 6525 ED Nijmegen, The Netherlands

Summary. A formalization of the first proof from [6].

MML identifier: `ARROW`, version: `7.8.05 4.87.985`

The terminology and notation used here are introduced in the following articles: [11], [13], [12], [10], [9], [5], [2], [3], [1], [8], [4], and [7].

1. Preliminaries

Let A, B' be non empty sets, let B be a non empty subset of B', let f be a function from A into B, and let x be an element of A. Then $f(x)$ is an element of B.

Next we state two propositions:

(1) For every finite set A such that $\operatorname{card} A \geq 2$ and for every element a of A there exists an element b of A such that $b \neq a$.

(2) Let A be a finite set. Suppose $\operatorname{card} A \geq 3$. Let a, b be elements of A. Then there exists an element c of A such that $c \neq a$ and $c \neq b$.

2. Linear Preorders and Linear Orders

In the sequel A denotes a non empty set and a, b, c denote elements of A.

Let us consider A. The functor $\operatorname{LinPreorders} A$ is defined by the condition (Def. 1).

© 2007 University of Białystok
ISSN 1426–2630

(Def. 1) Let R be a set. Then $R \in \mathrm{LinPreorders}\, A$ if and only if the following conditions are satisfied:

(i) R is a binary relation on A,

(ii) for all a, b holds $\langle a, b\rangle \in R$ or $\langle b, a\rangle \in R$, and

(iii) for all a, b, c such that $\langle a, b\rangle \in R$ and $\langle b, c\rangle \in R$ holds $\langle a, c\rangle \in R$.

Let us consider A. Note that $\mathrm{LinPreorders}\, A$ is non empty.

Let us consider A. The functor $\mathrm{LinOrders}\, A$ yielding a subset of $\mathrm{LinPreorders}\, A$ is defined by:

(Def. 2) For every element R of $\mathrm{LinPreorders}\, A$ holds $R \in \mathrm{LinOrders}\, A$ iff for all a, b such that $\langle a, b\rangle \in R$ and $\langle b, a\rangle \in R$ holds $a = b$.

Let A be a set. One can verify that there exists an order in A which is connected.

Let us consider A. Then $\mathrm{LinOrders}\, A$ can be characterized by the condition:

(Def. 3) For every set R holds $R \in \mathrm{LinOrders}\, A$ iff R is a connected order in A.

Let us consider A. One can verify that $\mathrm{LinOrders}\, A$ is non empty.

In the sequel o, o' are elements of $\mathrm{LinPreorders}\, A$ and o'' is an element of $\mathrm{LinOrders}\, A$.

Let us consider A, o, a, b. The predicate $a \leq_o b$ is defined by:

(Def. 4) $\langle a, b\rangle \in o$.

Let us consider A, o, a, b. We introduce $b \geq_o a$ as a synonym of $a \leq_o b$. We introduce $b <_o a$ as an antonym of $a \leq_o b$. We introduce $a >_o b$ as an antonym of $a \leq_o b$.

We now state a number of propositions:

(3) $a \leq_o a$.

(4) $a \leq_o b$ or $b \leq_o a$.

(5) If $a \leq_o b$ or $a <_o b$ and if $b \leq_o c$ or $b <_o c$, then $a \leq_o c$.

(6) If $a \leq_{o''} b$ and $b \leq_{o''} a$, then $a = b$.

(7) If $a \neq b$ and $b \neq c$ and $a \neq c$, then there exists o such that $a <_o b$ and $b <_o c$.

(8) There exists o such that for every a such that $a \neq b$ holds $b <_o a$.

(9) There exists o such that for every a such that $a \neq b$ holds $a <_o b$.

(10) If $a \neq b$ and $a \neq c$, then there exists o such that $a <_o b$ and $a <_o c$ and $b <_o c$ iff $b <_{o'} c$ and $c <_o b$ iff $c <_{o'} b$.

(11) If $a \neq b$ and $a \neq c$, then there exists o such that $b <_o a$ and $c <_o a$ and $b <_o c$ iff $b <_{o'} c$ and $c <_o b$ iff $c <_{o'} b$.

(12) Let o, o' be elements of $\mathrm{LinOrders}\, A$. Then $a <_o b$ iff $a <_{o'} b$ and $b <_o a$ iff $b <_{o'} a$ if and only if $a <_o b$ iff $a <_{o'} b$.

(13) Let o be an element of $\mathrm{LinOrders}\, A$ and o' be an element of $\mathrm{LinPreorders}\, A$. Then for all a, b such that $a <_o b$ holds $a <_{o'} b$ if and only

if for all a, b holds $a <_o b$ iff $a <_{o'} b$.

3. Arrow's Theorem

For simplicity, we follow the rules: A, N are finite non empty sets, a, b are elements of A, i, n are elements of N, p, p' are elements of $(\mathrm{LinPreorders}\, A)^N$, and f is a function from $(\mathrm{LinPreorders}\, A)^N$ into $\mathrm{LinPreorders}\, A$.

We now state the proposition

(14) Suppose that
  (i) for all p, a, b such that for every i holds $a <_{p(i)} b$ holds $a <_{f(p)} b$,
  (ii) for all p, p', a, b such that for every i holds $a <_{p(i)} b$ iff $a <_{p'(i)} b$ and $b <_{p(i)} a$ iff $b <_{p'(i)} a$ holds $a <_{f(p)} b$ iff $a <_{f(p')} b$, and
  (iii) $\mathrm{card}\, A \geq 3$.

  Then there exists n such that for all p, a, b such that $a <_{p(n)} b$ holds $a <_{f(p)} b$.

In the sequel p, p' denote elements of $(\mathrm{LinOrders}\, A)^N$ and f denotes a function from $(\mathrm{LinOrders}\, A)^N$ into $\mathrm{LinPreorders}\, A$.

One can prove the following proposition

(15) Suppose that
  (i) for all p, a, b such that for every i holds $a <_{p(i)} b$ holds $a <_{f(p)} b$,
  (ii) for all p, p', a, b such that for every i holds $a <_{p(i)} b$ iff $a <_{p'(i)} b$ holds $a <_{f(p)} b$ iff $a <_{f(p')} b$, and
  (iii) $\mathrm{card}\, A \geq 3$.

  Then there exists n such that for all p, a, b holds $a <_{p(n)} b$ iff $a <_{f(p)} b$.

References

[1] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(**2**):377–382, 1990.
[2] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(**1**):55–65, 1990.
[3] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(**1**):153–164, 1990.
[4] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(**1**):47–53, 1990.
[5] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(**1**):165–167, 1990.
[6] John Geanakoplos. Three brief proofs of arrow's impossibility theorem. Cowles Foundation Discussion Papers 1123R3, Cowles Foundation, Yale University, April 1996. Available at http://ideas.repec.org/p/cwl/cwldpp/1123r3.html.
[7] Andrzej Trybulec. Subsets of complex numbers. *To appear in Formalized Mathematics*.
[8] Andrzej Trybulec. Function domains and Frænkel operator. *Formalized Mathematics*, 1(**3**):495–500, 1990.
[9] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(**1**):9–11, 1990.
[10] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(**1**):67–71, 1990.
[11] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(**1**):73–83, 1990.
[12] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(**1**):181–186, 1990.

[13] Edmund Woronowicz and Anna Zalewska. Properties of binary relations. *Formalized Mathematics*, 1(**1**):85–89, 1990.

Received August 13, 2007