FORMALIZED MATHEMATICS
2007, Vol. **15**, No. **4**, Pages 189–198
DOI: 10.2478/v10037-007-0023-6

Several Integrability Formulas of Special Functions

Cuiying Peng
Qingdao University of Science and Technology
China

Fuguo Ge
Qingdao University of Science and Technology
China

Xiquan Liang
Qingdao University of Science and Technology
China

Summary. In this article, we give several integrability formulas of special and composite functions including trigonometric function, inverse trigonometric function, hyperbolic function and logarithmic function.

MML identifier: `INTEGRA8`, version: `7.8.05 4.87.985`

The notation and terminology used here are introduced in the following papers: [21], [20], [7], [12], [6], [25], [3], [8], [26], [24], [5], [22], [18], [19], [17], [9], [16], [11], [14], [1], [15], [23], [13], [10], [2], and [4].

1. Preliminaries

For simplicity, we adopt the following convention: f, f_1, f_2, g denote partial functions from $\mathbb{R}$ to $\mathbb{R}$, A denotes a closed-interval subset of $\mathbb{R}$, r, x, x_0 denote real numbers, n denotes an element of $\mathbb{N}$, and Z denotes an open subset of $\mathbb{R}$.

The following propositions are true:

(1) $\sin(x + 2 \cdot n \cdot \pi) = \sin x$.

(2) $\sin(x + (2 \cdot n + 1) \cdot \pi) = -\sin x$.

(3) $\cos(x + 2 \cdot n \cdot \pi) = \cos x$.

© 2007 University of Białystok
ISSN 1426–2630

(4) $\cos(x + (2 \cdot n + 1) \cdot \pi) = -\cos x$.

(5) If $\sin(\frac{x}{2}) \geq 0$, then $\sin(\frac{x}{2}) = \sqrt{\frac{1-\cos x}{2}}$.

(6) If $\sin(\frac{x}{2}) < 0$, then $\sin(\frac{x}{2}) = -\sqrt{\frac{1-\cos x}{2}}$.

(7) $\sin(\frac{\pi}{4}) = \frac{\sqrt{2}}{2}$.

(8) $\sin(-\frac{\pi}{4}) = -\frac{\sqrt{2}}{2}$.

(9) $[-\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}] \subseteq]-1, 1[$.

(10) $\arcsin(\frac{\sqrt{2}}{2}) = \frac{\pi}{4}$.

(11) $\arcsin(-\frac{\sqrt{2}}{2}) = -\frac{\pi}{4}$.

(12) If $\cos(\frac{x}{2}) \geq 0$, then $\cos(\frac{x}{2}) = \sqrt{\frac{1+\cos x}{2}}$.

(13) $\cos(\frac{\pi}{4}) = \frac{\sqrt{2}}{2}$.

(14) $\cos(\frac{3 \cdot \pi}{4}) = -\frac{\sqrt{2}}{2}$.

(15) $\arccos(\frac{\sqrt{2}}{2}) = \frac{\pi}{4}$.

(16) $\arccos(-\frac{\sqrt{2}}{2}) = \frac{3 \cdot \pi}{4}$.

(17) (The function sinh)(1) $= \frac{e^{\mathbf{2}}-1}{2 \cdot e}$.

(18) (The function cosh)(0) $= 1$.

(19) (The function cosh)(1) $= \frac{e^{\mathbf{2}}+1}{2 \cdot e}$.

(20) For every linear function L_1 holds $-L_1$ is a linear function.

(21) For every rest R_1 holds $-R_1$ is a rest.

(22) For all f_1, x_0 such that f_1 is differentiable in x_0 holds $-f_1$ is differentiable in x_0 and $(-f_1)'(x_0) = -{f_1}'(x_0)$.

(23) Let given f_1, Z. Suppose $Z \subseteq \mathrm{dom}(-f_1)$ and f_1 is differentiable on Z. Then $-f_1$ is differentiable on Z and for every x such that $x \in Z$ holds $(-f_1)'_{\restriction Z}(x) = -{f_1}'(x)$.

(24) $-$the function sin is differentiable on $\mathbb{R}$.

(25) $-$the function cos is differentiable in x and $(-\text{the function cos})'(x) =$ (the function sin)(x).

(26)(i) $-$the function cos is differentiable on $\mathbb{R}$, and

(ii) for every x such that $x \in \mathbb{R}$ holds $(-\text{the function cos})'(x) =$ (the function sin)(x).

(27) $(\text{The function sin})'_{\restriction \mathbb{R}} =$ the function cos.

(28) $(\text{The function cos})'_{\restriction \mathbb{R}} = -$the function sin.

(29) $(-\text{the function cos})'_{\restriction \mathbb{R}} =$ the function sin.

(30) $(\text{The function sinh})'_{\restriction \mathbb{R}} =$ the function cosh.

(31) $(\text{The function cosh})'_{\restriction \mathbb{R}} =$ the function sinh.

(32) $(\text{The function exp})'_{\restriction \mathbb{R}} =$ the function exp.

(33) Suppose $Z \subseteq \mathrm{dom}\,(\text{the function tan})$ and for every x such that $x \in Z$ holds $f(x) = \frac{1}{(\text{the function cos})(x)^2}$ and $(\text{the function cos})(x) \neq 0$. Then

(i) the function tan is differentiable on Z, and

(ii) for every x such that $x \in Z$ holds $(\text{the function tan})'_{\restriction Z}(x) = \frac{1}{(\text{the function cos})(x)^2}$.

(34) Suppose that

(i) $Z \subseteq \mathrm{dom}\,(\text{the function cot})$, and

(ii) for every x such that $x \in Z$ holds $f(x) = -\frac{1}{(\text{the function sin})(x)^2}$ and $(\text{the function sin})(x) \neq 0$.

Then

(iii) the function cot is differentiable on Z, and

(iv) for every x such that $x \in Z$ holds $(\text{the function cot})'_{\restriction Z}(x) = -\frac{1}{(\text{the function sin})(x)^2}$.

(35) For every real number r holds $\mathrm{dom}(\mathbb{R} \longmapsto r) = \mathbb{R}$ and $\mathrm{rng}(\mathbb{R} \longmapsto r) \subseteq \mathbb{R}$.

Let r be a real number. The functor $\mathrm{Cst}\, r$ yielding a function from $\mathbb{R}$ into $\mathbb{R}$ is defined as follows:

(Def. 1) $\mathrm{Cst}\, r = \mathbb{R} \longmapsto r$.

We now state two propositions:

(36) For all real numbers a, b and for every closed-interval subset A of $\mathbb{R}$ holds $\chi_{A,A} = \mathrm{Cst}\, 1 \restriction A$.

(37) For all real numbers a, b and for every closed-interval subset A of $\mathbb{R}$ such that $A = [a, b]$ holds $\sup A = b$ and $\inf A = a$.

2. Several Integrability Formulas of Special Functions

The following propositions are true:

(38) For all real numbers a, b such that $a \leq b$ holds $\int_a^b \mathrm{Cst}\, 1(x)dx = b - a$.

(39) $\int_A (\text{the function cos})(x)dx = (\text{the function sin})(\sup A) - (\text{the function sin})(\inf A)$.

(40) If $A = [0, \frac{\pi}{2}]$, then $\int_A (\text{the function cos})(x)dx = 1$.

(41) If $A = [0, \pi]$, then $\int_A (\text{the function cos})(x)dx = 0$.

(42) If $A = [0, \frac{\pi \cdot 3}{2}]$, then $\int_A (\text{the function cos})(x)dx = -1$.

(43) If $A = [0, \pi \cdot 2]$, then $\int\limits_A (\text{the function cos})(x)dx = 0$.

(44) If $A = [2 \cdot n \cdot \pi, (2 \cdot n + 1) \cdot \pi]$, then $\int\limits_A (\text{the function cos})(x)dx = 0$.

(45) If $A = [x + 2 \cdot n \cdot \pi, x + (2 \cdot n + 1) \cdot \pi]$, then $\int\limits_A (\text{the function cos})(x)dx = -2 \cdot \sin x$.

(46) $\int\limits_A (-\text{the function sin})(x)dx = (\text{the function cos})(\sup A) - (\text{the function cos})(\inf A)$.

(47) If $A = [0, \frac{\pi}{2}]$, then $\int\limits_A (-\text{the function sin})(x)dx = -1$.

(48) If $A = [0, \pi]$, then $\int\limits_A (-\text{the function sin})(x)dx = -2$.

(49) If $A = [0, \frac{\pi \cdot 3}{2}]$, then $\int\limits_A (-\text{the function sin})(x)dx = -1$.

(50) If $A = [0, \pi \cdot 2]$, then $\int\limits_A (-\text{the function sin})(x)dx = 0$.

(51) If $A = [2 \cdot n \cdot \pi, (2 \cdot n + 1) \cdot \pi]$, then $\int\limits_A (-\text{the function sin})(x)dx = -2$.

(52) If $A = [x + 2 \cdot n \cdot \pi, x + (2 \cdot n + 1) \cdot \pi]$, then $\int\limits_A (-\text{the function sin})(x)dx = -2 \cdot \cos x$.

(53) $\int\limits_A (\text{the function exp})(x)dx = (\text{the function exp})(\sup A) - (\text{the function exp})(\inf A)$.

(54) If $A = [0, 1]$, then $\int\limits_A (\text{the function exp})(x)dx = e - 1$.

(55) $\int\limits_A (\text{the function sinh})(x)dx = (\text{the function cosh})(\sup A) - (\text{the function cosh})(\inf A)$.

(56) If $A = [0, 1]$, then $\int\limits_A (\text{the function sinh})(x)dx = \dfrac{(e-1)^{\mathbf{2}}}{2 \cdot e}$.

(57) $\int\limits_A (\text{the function cosh})(x)dx = (\text{the function sinh})(\sup A) - (\text{the function}$

sinh)(inf A).

(58) If $A = [0, 1]$, then $\int_A$ (the function cosh)$(x)dx = \frac{e^2 - 1}{2 \cdot e}$.

(59) Suppose that
(i) $A \subseteq Z$,
(ii) dom (the function tan) $= Z$,
(iii) dom (the function tan) $=$ dom f_2,
(iv) for every x such that $x \in Z$ holds $f_2(x) = \frac{1}{(\text{the function cos})(x)^2}$ and (the function cos)$(x) \neq 0$, and
(v) f_2 is continuous on A.
Then $\int_A f_2(x)dx =$ (the function tan)(sup A) $-$ (the function tan)(inf A).

(60) Suppose that
(i) $A \subseteq Z$,
(ii) dom (the function cot) $= Z$,
(iii) dom (the function cot) $=$ dom f_2,
(iv) for every x such that $x \in Z$ holds $f_2(x) = -\frac{1}{(\text{the function sin})(x)^2}$ and (the function sin)$(x) \neq 0$, and
(v) f_2 is continuous on A.
Then $\int_A f_2(x)dx =$ (the function cot)(sup A) $-$ (the function cot)(inf A).

(61) Suppose dom (the function tanh) $=$ dom f_2 and for every x such that $x \in \mathbb{R}$ holds $f_2(x) = \frac{1}{(\text{the function cosh})(x)^2}$ and f_2 is continuous on A. Then $\int_A f_2(x)dx =$ (the function tanh)(sup A) $-$ (the function tanh)(inf A).

(62) Suppose $A \subseteq]-1, 1[$ and dom((the function arcsin)$'_{]-1,1[}$) $=$ dom f_2 and for every x holds $x \in]-1, 1[$ and $f_2(x) = \frac{1}{\sqrt{1-x^2}}$ and f_2 is continuous on A. Then $\int_A f_2(x)dx =$ (the function arcsin)(sup A) $-$ (the function arcsin)(inf A).

(63) Suppose $A \subseteq]-1, 1[$ and dom((the function arccos)$'_{]-1,1[}$) $=$ dom f_2 and for every x holds $x \in]-1, 1[$ and $f_2(x) = -\frac{1}{\sqrt{1-x^2}}$ and f_2 is continuous on A. Then $\int_A f_2(x)dx =$ (the function arccos)(sup A) $-$ (the function arccos)(inf A).

(64) Suppose that
(i) $A = [-\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}]$,
(ii) dom((the function arcsin)$'_{]-1,1[}$) $=$ dom f_2,

(iii) for every x holds $x \in]-1,1[$ and $f_2(x) = \frac{1}{\sqrt{1-x^2}}$, and

(iv) f_2 is continuous on A.

Then $\int\limits_A f_2(x)dx = \frac{\pi}{2}$.

(65) Suppose that

(i) $A = [-\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}]$,

(ii) $\operatorname{dom}((\text{the function arccos})'_{\restriction]-1,1[}) = \operatorname{dom} f_2$,

(iii) for every x holds $x \in]-1,1[$ and $f_2(x) = -\frac{1}{\sqrt{1-x^2}}$, and

(iv) f_2 is continuous on A.

Then $\int\limits_A f_2(x)dx = -\frac{\pi}{2}$.

(66) Suppose that f is differentiable on Z and g is differentiable on Z and $A \subseteq Z$ and $f'_{\restriction Z}$ is integrable on A and $f'_{\restriction Z}$ is bounded on A and $g'_{\restriction Z}$ is integrable on A and $g'_{\restriction Z}$ is bounded on A. Then $\int\limits_A (f'_{\restriction Z} + g'_{\restriction Z})(x)dx = ((f(\sup A) - f(\inf A)) + g(\sup A)) - g(\inf A)$.

(67) Suppose that f is differentiable on Z and g is differentiable on Z and $A \subseteq Z$ and $f'_{\restriction Z}$ is integrable on A and $f'_{\restriction Z}$ is bounded on A and $g'_{\restriction Z}$ is integrable on A and $g'_{\restriction Z}$ is bounded on A. Then $\int\limits_A (f'_{\restriction Z} - g'_{\restriction Z})(x)dx = f(\sup A) - f(\inf A) - (g(\sup A) - g(\inf A))$.

(68) Suppose f is differentiable on Z and $A \subseteq Z$ and $f'_{\restriction Z}$ is integrable on A and $f'_{\restriction Z}$ is bounded on A. Then $\int\limits_A (r\, f'_{\restriction Z})(x)dx = r \cdot f(\sup A) - r \cdot f(\inf A)$.

(69) $\int\limits_A ((\text{the function sin}) + (\text{the function cos}))(x)dx = (((-\text{the function cos})(\sup A) - (-\text{the function cos})(\inf A)) + (\text{the function sin})(\sup A)) - (\text{the function sin})(\inf A)$.

(70) If $A = [0, \frac{\pi}{2}]$, then $\int\limits_A ((\text{the function sin}) + (\text{the function cos}))(x)dx = 2$.

(71) If $A = [0, \pi]$, then $\int\limits_A ((\text{the function sin}) + (\text{the function cos}))(x)dx = 2$.

(72) If $A = [0, \frac{\pi \cdot 3}{2}]$, then $\int\limits_A ((\text{the function sin}) + (\text{the function cos}))(x)dx = 0$.

(73) If $A = [0, \pi \cdot 2]$, then $\int\limits_A ((\text{the function sin}) + (\text{the function cos}))(x)dx =$

0.

(74) If $A = [2 \cdot n \cdot \pi, (2 \cdot n + 1) \cdot \pi]$, then $\int_A ((\text{the function sin}) + (\text{the function cos}))(x)dx = 2$.

(75) If $A = [x + 2 \cdot n \cdot \pi, x + (2 \cdot n + 1) \cdot \pi]$, then $\int_A ((\text{the function sin}) + (\text{the function cos}))(x)dx = 2 \cdot \cos x - 2 \cdot \sin x$.

(76) $\int_A ((\text{the function sinh}) + (\text{the function cosh}))(x)dx = (((\text{the function cosh})(\sup A) - (\text{the function cosh})(\inf A)) + (\text{the function sinh})(\sup A)) - (\text{the function sinh})(\inf A)$.

(77) If $A = [0, 1]$, then $\int_A ((\text{the function sinh}) + (\text{the function cosh}))(x)dx = e - 1$.

(78) $\int_A ((\text{the function sin}) - (\text{the function cos}))(x)dx = (-\text{the function cos})(\sup A) - (-\text{the function cos})(\inf A) - ((\text{the function sin})(\sup A) - (\text{the function sin})(\inf A))$.

(79) If $A = [0, \frac{\pi}{2}]$, then $\int_A ((\text{the function sin}) - (\text{the function cos}))(x)dx = 0$.

(80) If $A = [0, \pi]$, then $\int_A ((\text{the function sin}) - (\text{the function cos}))(x)dx = 2$.

(81) If $A = [0, \frac{\pi \cdot 3}{2}]$, then $\int_A ((\text{the function sin}) - (\text{the function cos}))(x)dx = 2$.

(82) If $A = [0, \pi \cdot 2]$, then $\int_A ((\text{the function sin}) - (\text{the function cos}))(x)dx = 0$.

(83) If $A = [2 \cdot n \cdot \pi, (2 \cdot n + 1) \cdot \pi]$, then $\int_A ((\text{the function sin}) - (\text{the function cos}))(x)dx = 2$.

(84) If $A = [x + 2 \cdot n \cdot \pi, x + (2 \cdot n + 1) \cdot \pi]$, then $\int_A ((\text{the function sin}) - (\text{the function cos}))(x)dx = 2 \cdot \cos x + 2 \cdot \sin x$.

(85) $\int_A (r\,(\text{the function sin}))(x)dx = r \cdot (-\text{the function cos})(\sup A) - r \cdot (-\text{the function cos})(\inf A)$.

(86) $\int_A (r\,(\text{the function cos}))(x)dx = r \cdot (\text{the function sin})(\sup A) - r \cdot (\text{the function sin})(\inf A)$.

(87) $\int_A (r\,(\text{the function sinh}))(x)dx = r \cdot (\text{the function cosh})(\sup A) - r \cdot (\text{the function cosh})(\inf A)$.

(88) $\int_A (r\,(\text{the function cosh}))(x)dx = r \cdot (\text{the function sinh})(\sup A) - r \cdot (\text{the function sinh})(\inf A)$.

(89) $\int_A (r\,(\text{the function exp}))(x)dx = r \cdot (\text{the function exp})(\sup A) - r \cdot (\text{the function exp})(\inf A)$.

(90) $\int_A ((\text{the function sin})\,(\text{the function cos}))(x)dx = \frac{1}{2} \cdot ((\text{the function cos})(\inf A) \cdot (\text{the function cos})(\inf A) - (\text{the function cos})(\sup A) \cdot (\text{the function cos})(\sup A))$.

(91) If $A = [0, \frac{\pi}{2}]$, then $\int_A ((\text{the function sin})\,(\text{the function cos}))(x)dx = \frac{1}{2}$.

(92) If $A = [0, \pi]$, then $\int_A ((\text{the function sin})\,(\text{the function cos}))(x)dx = 0$.

(93) If $A = [0, \pi \cdot \frac{3}{2}]$, then $\int_A ((\text{the function sin})\,(\text{the function cos}))(x)dx = \frac{1}{2}$.

(94) If $A = [0, \pi \cdot 2]$, then $\int_A ((\text{the function sin})\,(\text{the function cos}))(x)dx = 0$.

(95) If $A = [2 \cdot n \cdot \pi, (2 \cdot n + 1) \cdot \pi]$, then $\int_A ((\text{the function sin})\,(\text{the function cos}))(x)dx = 0$.

(96) If $A = [x + 2 \cdot n \cdot \pi, x + (2 \cdot n + 1) \cdot \pi]$, then $\int_A ((\text{the function sin})\,(\text{the function cos}))(x)dx = 0$.

(97) $\int_A ((\text{the function sin})\,(\text{the function sin}))(x)dx = ((\text{the function cos})(\inf A) \cdot (\text{the function sin})(\inf A) - (\text{the function cos})(\sup A) \cdot (\text{the function sin})(\sup A)) + \int_A ((\text{the function cos})\,(\text{the function cos}))(x)dx$.

(98) $\int\limits_{A}((\text{the function sinh})\,(\text{the function sinh}))(x)dx = (\text{the function cosh})(\sup A)\cdot(\text{the function sinh})(\sup A) - (\text{the function cosh})(\inf A)\cdot(\text{the function sinh})(\inf A) - \int\limits_{A}((\text{the function cosh})\,(\text{the function cosh}))(x)dx$.

(99) $\int\limits_{A}((\text{the function sinh})\,(\text{the function cosh}))(x)dx = \frac{1}{2}\cdot((\text{the function cosh})(\sup A)\cdot(\text{the function cosh})(\sup A) - (\text{the function cosh})(\inf A)\cdot(\text{the function cosh})(\inf A))$.

(100) $\int\limits_{A}((\text{the function exp})\,(\text{the function exp}))(x)dx = \frac{1}{2}\cdot((\text{the function exp})(\sup A)^{\mathbf{2}} - (\text{the function exp})(\inf A)^{\mathbf{2}})$.

(101) $\int\limits_{A}((\text{the function exp})\,((\text{the function sin}) + (\text{the function cos})))(x)dx = ((\text{the function exp})\ (\text{the function sin}))(\sup A) - ((\text{the function exp})\ (\text{the function sin}))(\inf A)$.

(102) $\int\limits_{A}((\text{the function exp})\,((\text{the function cos}) - (\text{the function sin})))(x)dx = ((\text{the function exp})\ (\text{the function cos}))(\sup A) - ((\text{the function exp})\ (\text{the function cos}))(\inf A)$.

References

[1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(**1**):41–46, 1990.

[2] Czesław Byliński. The complex numbers. *Formalized Mathematics*, 1(**3**):507–513, 1990.

[3] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(**1**):55–65, 1990.

[4] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(**1**):153–164, 1990.

[5] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(**2**):357–367, 1990.

[6] Czesław Byliński and Piotr Rudnicki. Bounding boxes for compact sets in $\mathcal{E}^2$. *Formalized Mathematics*, 6(**3**):427–440, 1997.

[7] Noboru Endou and Artur Korniłowicz. The definition of the Riemann definite integral and some related lemmas. *Formalized Mathematics*, 8(**1**):93–102, 1999.

[8] Noboru Endou, Katsumi Wasaki, and Yasunari Shidama. Definition of integrability for partial functions from $\mathbb{R}$ to $\mathbb{R}$ and integrability for continuous functions. *Formalized Mathematics*, 9(**2**):281–284, 2001.

[9] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(**1**):35–40, 1990.

[10] Artur Korniłowicz and Yasunari Shidama. Inverse trigonometric functions arcsin and arccos. *Formalized Mathematics*, 13(**1**):73–79, 2005.

[11] Jarosław Kotowicz. Convergent sequences and the limit of sequences. *Formalized Mathematics*, 1(**2**):273–275, 1990.

[12] Jarosław Kotowicz. Partial functions from a domain to the set of real numbers. *Formalized Mathematics*, 1(**4**):703–709, 1990.

[13] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(**2**):269–272, 1990.
[14] Takashi Mitsuishi and Yuguang Yang. Properties of the trigonometric function. *Formalized Mathematics*, 8(**1**):103–106, 1999.
[15] Konrad Raczkowski and Andrzej Nędzusiak. Real exponents and logarithms. *Formalized Mathematics*, 2(**2**):213–216, 1991.
[16] Konrad Raczkowski and Paweł Sadowski. Real function continuity. *Formalized Mathematics*, 1(**4**):787–791, 1990.
[17] Konrad Raczkowski and Paweł Sadowski. Real function differentiability. *Formalized Mathematics*, 1(**4**):797–801, 1990.
[18] Konrad Raczkowski and Paweł Sadowski. Topological properties of subsets in real numbers. *Formalized Mathematics*, 1(**4**):777–780, 1990.
[19] Andrzej Trybulec. Subsets of complex numbers. *To appear in Formalized Mathematics*.
[20] Andrzej Trybulec. Binary operations applied to functions. *Formalized Mathematics*, 1(**2**):329–334, 1990.
[21] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(**1**):9–11, 1990.
[22] Andrzej Trybulec. On the sets inhabited by numbers. *Formalized Mathematics*, 11(**4**):341–347, 2003.
[23] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers. *Formalized Mathematics*, 1(**3**):445–449, 1990.
[24] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(**1**):67–71, 1990.
[25] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(**1**):73–83, 1990.
[26] Yuguang Yang and Yasunari Shidama. Trigonometric functions and existence of circle ratio. *Formalized Mathematics*, 7(**2**):255–263, 1998.

Received August 28, 2007