FORMALIZED MATHEMATICS
2006, Vol. **14**, No. **4**, Pages 153–159
DOI: 10.2478/v10037-006-0019-7

The Catalan Numbers. Part II[1]

Karol Pąk
Institute of Mathematics
University of Białystok
Akademicka 2, 15-267 Białystok, Poland

Summary. In this paper, we define sequence dominated by 0, in which every initial fragment contains more zeroes than ones. If $n \geq 2 \cdot m$ and $n > 0$, then the number of sequences dominated by 0 the length n including m of ones, is given by the formula

$$D(n,m) = \frac{n+1-2\cdot m}{n+1-m} \cdot \binom{n}{m}$$

and satisfies the recurrence relation

$$D(n,m) = D(n-1,m) \ + \ \sum_{i=0}^{m-1} D(2\cdot i, i) \cdot D(n-2\cdot(i+1), m-(i+1)).$$

Obviously, if $n = 2 \cdot m$, then we obtain the recurrence relation for the Catalan numbers (starting from 0)

$$C_{m+1} = \sum_{i=0}^{m-1} C_{i+1} \cdot C_{m-i}.$$

Using the above recurrence relation we can see that

$$\sum_{i=0}^{\infty} C_{i+1} \cdot x^i \ = \ 1 \ + \ \left(\sum_{i=0}^{\infty} C_{i+1} \cdot x^i \right)^2$$

where ($|x| < \frac{1}{4}$) and hence

$$\sum_{i=0}^{\infty} C_{i+1} \cdot x^i \ = \ \frac{1-\sqrt{1-4\cdot x}}{2\cdot x}.$$

MML identifier: `CATALAN2`, version: `7.8.03 4.75.958`

[1]This work has been partially supported by the KBN grant 4 T11C 039 24 and the FP6 IST grant TYPES No. 510996.

© 2006 University of Białystok
ISSN 1426–2630

The notation and terminology used here are introduced in the following papers: [2], [23], [7], [25], [19], [27], [5], [28], [9], [1], [26], [21], [6], [3], [14], [12], [16], [13], [20], [15], [8], [22], [11], [10], [18], [24], [17], and [4].

1. Preliminaries

For simplicity, we adopt the following convention: x, D denote sets, i, j, k, l, m, n denote elements of $\mathbb{N}$, p, q denote finite 0-sequences of $\mathbb{N}$, p', q' denote finite 0-sequences, and p_1, q_1 denote finite 0-sequences of D.

Next we state several propositions:

(1) $(p' \frown q')\restriction \operatorname{dom} p' = p'$.

(2) If $n \leq \operatorname{dom} p'$, then $(p' \frown q')\restriction n = p'\restriction n$.

(3) If $n = \operatorname{dom} p' + k$, then $(p' \frown q')\restriction n = p' \frown (q'\restriction k)$.

(4) There exists q' such that $p' = (p'\restriction n) \frown q'$.

(5) There exists q_1 such that $p_1 = (p_1\restriction n) \frown q_1$.

Let us consider p. We say that p is dominated by 0 if and only if:

(Def. 1) $\operatorname{rng} p \subseteq \{0, 1\}$ and for every k such that $k \leq \operatorname{dom} p$ holds $2 \cdot \sum(p\restriction k) \leq k$.

The following propositions are true:

(6) If p is dominated by 0, then $2 \cdot \sum(p\restriction k) \leq k$.

(7) If p is dominated by 0, then $p(0) = 0$.

Let us consider k, m. Then $k \longmapsto m$ is a finite 0-sequence of $\mathbb{N}$.

One can check that there exists a finite 0-sequence of $\mathbb{N}$ which is empty and dominated by 0 and there exists a finite 0-sequence of $\mathbb{N}$ which is non empty and dominated by 0.

The following propositions are true:

(8) $n \longmapsto 0$ is dominated by 0.

(9) If $n \geq m$, then $(n \longmapsto 0) \frown (m \longmapsto 1)$ is dominated by 0.

(10) If p is dominated by 0, then $p\restriction n$ is dominated by 0.

(11) If p is dominated by 0 and q is dominated by 0, then $p \frown q$ is dominated by 0.

(12) If p is dominated by 0, then $2 \cdot \sum(p\restriction(2 \cdot n + 1)) < 2 \cdot n + 1$.

(13) If p is dominated by 0 and $n \leq \operatorname{len} p - 2 \cdot \sum p$, then $p \frown (n \longmapsto 1)$ is dominated by 0.

(14) If p is dominated by 0 and $n \leq (k + \operatorname{len} p) - 2 \cdot \sum p$, then $(k \longmapsto 0) \frown p \frown (n \longmapsto 1)$ is dominated by 0.

(15) If p is dominated by 0 and $2 \cdot \sum(p\restriction k) = k$, then $k \leq \operatorname{len} p$ and $\operatorname{len}(p\restriction k) = k$.

(16) If p is dominated by 0 and $2 \cdot \sum(p\restriction k) = k$ and $p = (p\restriction k) \frown q$, then q is dominated by 0.

(17) If p is dominated by 0 and $2 \cdot \sum(p\restriction k) = k$ and $k = n + 1$, then $p\restriction k = (p\restriction n) \frown (1 \longmapsto 1)$.

(18) Let given m, p. Suppose $m = \min^*\{n : 2 \cdot \sum(p\restriction n) = n \ \wedge \ n > 0\}$ and $m > 0$ and p is dominated by 0. Then there exists q such that $p\restriction m = (1 \longmapsto 0) \frown q \frown (1 \longmapsto 1)$ and q is dominated by 0.

(19) Let given p. Suppose $\operatorname{rng} p \subseteq \{0, 1\}$ and p is not dominated by 0. Then there exists k such that $2 \cdot k + 1 = \min^*\{n : 2 \cdot \sum(p\restriction n) > n\}$ and $2 \cdot k + 1 \leq \operatorname{dom} p$ and $k = \sum(p\restriction(2 \cdot k))$ and $p(2 \cdot k) = 1$.

(20) Let given p, q, k. Suppose $p\restriction(2 \cdot k + \operatorname{len} q) = (k \longmapsto 0) \frown q \frown (k \longmapsto 1)$ and q is dominated by 0 and $2 \cdot \sum q = \operatorname{len} q$ and $k > 0$. Then $\min^*\{n : 2 \cdot \sum(p\restriction n) = n \ \wedge \ n > 0\} = 2 \cdot k + \operatorname{len} q$.

(21) Let given p. Suppose p is dominated by 0 and $\{i : 2 \cdot \sum(p\restriction i) = i \ \wedge \ i > 0\} = \emptyset$ and $\operatorname{len} p > 0$. Then there exists q such that $p = \langle 0\rangle \frown q$ and q is dominated by 0.

(22) If p is dominated by 0, then $\langle 0\rangle \frown p$ is dominated by 0 and $\{i : 2 \cdot \sum((\langle 0\rangle \frown p)\restriction i) = i \ \wedge \ i > 0\} = \emptyset$.

(23) If $\operatorname{rng} p \subseteq \{0, 1\}$ and p is not dominated by 0 and $2 \cdot k + 1 = \min^*\{n : 2 \cdot \sum(p\restriction n) > n\}$, then $p\restriction(2 \cdot k)$ is dominated by 0.

2. The Recurrence Relation for the Catalan Numbers

Let n, m be natural numbers. The functor $\operatorname{Domin}_0(n, m)$ yields a subset of $\{0, 1\}^\omega$ and is defined as follows:

(Def. 2) $x \in \operatorname{Domin}_0(n, m)$ iff there exists a finite 0-sequence p of $\mathbb{N}$ such that $p = x$ and p is dominated by 0 and $\operatorname{dom} p = n$ and $\sum p = m$.

Next we state two propositions:

(24) $p \in \operatorname{Domin}_0(n, m)$ iff p is dominated by 0 and $\operatorname{dom} p = n$ and $\sum p = m$.

(25) $\operatorname{Domin}_0(n, m) \subseteq \operatorname{Choose}(n, m, 1, 0)$.

Let us consider n, m. One can check that $\operatorname{Domin}_0(n, m)$ is finite.

One can prove the following propositions:

(26) $\operatorname{Domin}_0(n, m)$ is empty iff $2 \cdot m > n$.

(27) $\operatorname{Domin}_0(n, 0) = \{n \longmapsto 0\}$.

(28) $\operatorname{card} \operatorname{Domin}_0(n, 0) = 1$.

(29) Let given p, n. Suppose $\operatorname{rng} p \subseteq \{0, n\}$. Then there exists q such that $\operatorname{len} p = \operatorname{len} q$ and $\operatorname{rng} q \subseteq \{0, n\}$ and for every k such that $k \leq \operatorname{len} p$ holds $\sum(p\restriction k) + \sum(q\restriction k) = n \cdot k$ and for every k such that $k \in \operatorname{len} p$ holds $q(k) = n - p(k)$.

(30) If $m \leq n$, then $\binom{n}{m} > 0$.

(31) If $2\cdot(m+1)\leq n$, then $\operatorname{card}(\operatorname{Choose}(n,m+1,1,0)\setminus\operatorname{Domin}_0(n,m+1))=\operatorname{card}\operatorname{Choose}(n,m,1,0)$.

(32) If $2\cdot(m+1)\leq n$, then $\operatorname{card}\operatorname{Domin}_0(n,m+1)=\binom{n}{m+1}-\binom{n}{m}$.

(33) If $2\cdot m\leq n$, then $\operatorname{card}\operatorname{Domin}_0(n,m)=\frac{(n+1)-2\cdot m}{(n+1)-m}\cdot\binom{n}{m}$.

(34) $\operatorname{card}\operatorname{Domin}_0(2+k,1)=k+1$.

(35) $\operatorname{card}\operatorname{Domin}_0(4+k,2)=\frac{(k+1)\cdot(k+4)}{2}$.

(36) $\operatorname{card}\operatorname{Domin}_0(6+k,3)=\frac{(k+1)\cdot(k+5)\cdot(k+6)}{6}$.

(37) $\operatorname{card}\operatorname{Domin}_0(2\cdot n,n)=\frac{\binom{2\cdot n}{n}}{n+1}$.

(38) $\operatorname{card}\operatorname{Domin}_0(2\cdot n,n)=\operatorname{Catalan}(n+1)$.

Let us consider D. A functional non empty set is said to be a set of ω-sequences of D if:

(Def. 3) For every x such that $x\in$ it holds x is a finite 0-sequence of D.

Let us consider D. Then D^ω is a set of ω-sequences of D. Let F be a set of ω-sequences of D. We see that the element of F is a finite 0-sequence of D.

In the sequel p_2 denotes an element of $\mathbb{N}^\omega$.

We now state several propositions:

(39) $\overline{\overline{\{p_2:\operatorname{dom}p_2=2\cdot n\ \wedge\ p_2\text{ is dominated by }0\}}}=\binom{2\cdot n}{n}$.

(40) Let given n, m, k, j, l. Suppose $j=n-2\cdot(k+1)$ and $l=m-(k+1)$. Then $\overline{\overline{\{p_2:p_2\in\operatorname{Domin}_0(n,m)\ \wedge\ 2\cdot(k+1)=\min^*\{i:2\cdot\sum(p_2\restriction i)=i\ \wedge\ i>0\}\}}}=\operatorname{card}\operatorname{Domin}_0(2\cdot k,k)\cdot\operatorname{card}\operatorname{Domin}_0(j,l)$.

(41) Let given n, m. Suppose $2\cdot m\leq n$. Then there exists a finite 0-sequence C_1 of $\mathbb{N}$ such that $\overline{\overline{\{p_2:p_2\in\operatorname{Domin}_0(n,m)\ \wedge\ \{i:2\cdot\sum(p_2\restriction i)=i\ \wedge\ i>0\}\neq\emptyset\}}}=\sum C_1$ and $\operatorname{dom}C_1=m$ and for every j such that $j<m$ holds $C_1(j)=\operatorname{card}\operatorname{Domin}_0(2\cdot j,j)\cdot\operatorname{card}\operatorname{Domin}_0(n-'2\cdot(j+1),m-'(j+1))$.

(42) For every n such that $n>0$ holds $\operatorname{Domin}_0(2\cdot n,n)=\{p_2:p_2\in\operatorname{Domin}_0(2\cdot n,n)\ \wedge\ \{i:2\cdot\sum(p_2\restriction i)=i\ \wedge\ i>0\}\neq\emptyset\}$.

(43) Let given n. Suppose $n>0$. Then there exists a finite 0-sequence C_2 of $\mathbb{N}$ such that $\sum C_2=\operatorname{Catalan}(n+1)$ and $\operatorname{dom}C_2=n$ and for every j such that $j<n$ holds $C_2(j)=\operatorname{Catalan}(j+1)\cdot\operatorname{Catalan}(n-'j)$.

(44) $\overline{\overline{\{p_2:p_2\in\operatorname{Domin}_0(n+1,m)\ \wedge\ \{i:2\cdot\sum(p_2\restriction i)=i\ \wedge\ i>0\}=\emptyset\}}}=\operatorname{card}\operatorname{Domin}_0(n,m)$.

(45) Let given n, m. Suppose $2\cdot m\leq n$. Then there exists a finite 0-sequence C_1 of $\mathbb{N}$ such that $\operatorname{card}\operatorname{Domin}_0(n,m)=\sum C_1+\operatorname{card}\operatorname{Domin}_0(n-'1,m)$ and $\operatorname{dom}C_1=m$ and for every j such that $j<m$ holds $C_1(j)=\operatorname{card}\operatorname{Domin}_0(2\cdot j,j)\cdot\operatorname{card}\operatorname{Domin}_0(n-'2\cdot(j+1),m-'(j+1))$.

(46) For all n, k there exists p such that $\sum p=\operatorname{card}\operatorname{Domin}_0(2\cdot n+2+k,n+1)$ and $\operatorname{dom}p=k+1$ and for every i such that $i\leq k$ holds $p(i)=$

$\operatorname{card} \operatorname{Domin}_0(2 \cdot n + 1 + i, n)$.

3. Cauchy Product

We use the following convention: s_1, s_2, s_3 denote sequences of real numbers, r denotes a real number, and F_1, F_2, F_3 denote finite 0-sequences of $\mathbb{R}$.

Let us consider F_1. The functor $\sum F_1$ yields a real number and is defined as follows:

(Def. 4) $\sum F_1 = +_{\mathbb{R}} \odot F_1$.

Let us consider F_1, x. Then $F_1(x)$ is a real number.

Let s_1, s_2 be sequences of real numbers. The functor $s_1 (\#) s_2$ yields a sequence of real numbers and is defined by the condition (Def. 5).

(Def. 5) Let k be a natural number. Then there exists a finite 0-sequence F_1 of $\mathbb{R}$ such that $\operatorname{dom} F_1 = k + 1$ and for every n such that $n \in k + 1$ holds $F_1(n) = s_1(n) \cdot s_2(k -' n)$ and $\sum F_1 = (s_1 (\#) s_2)(k)$.

Let us notice that the functor $s_1 (\#) s_2$ is commutative.

One can prove the following propositions:

(47) For all F_1, n such that $n \in \operatorname{dom} F_1$ holds $\sum(F_1 \restriction n) + F_1(n) = \sum(F_1 \restriction (n + 1))$.

(48) For all F_2, F_3 such that $\operatorname{dom} F_2 = \operatorname{dom} F_3$ and for every n such that $n \in \operatorname{len} F_2$ holds $F_2(n) = F_3(\operatorname{len} F_2 -' (1 + n))$ holds $\sum F_2 = \sum F_3$.

(49) For all F_2, F_3 such that $\operatorname{dom} F_2 = \operatorname{dom} F_3$ and for every n such that $n \in \operatorname{len} F_2$ holds $F_2(n) = r \cdot F_3(n)$ holds $\sum F_2 = r \cdot \sum F_3$.

(50) $s_1 (\#) r\, s_2 = r\, (s_1 (\#) s_2)$.

(51) $s_1 (\#)(s_2 + s_3) = (s_1 (\#) s_2) + (s_1 (\#) s_3)$.

(52) $(s_1 (\#) s_2)(0) = s_1(0) \cdot s_2(0)$.

(53) For all s_1, s_2, n there exists F_1 such that $(\sum_{\alpha=0}^{\kappa}(s_1 (\#) s_2)(\alpha))_{\kappa \in \mathbb{N}}(n) = \sum F_1$ and $\operatorname{dom} F_1 = n + 1$ and for every i such that $i \in n + 1$ holds $F_1(i) = s_1(i) \cdot (\sum_{\alpha=0}^{\kappa}(s_2)(\alpha))_{\kappa \in \mathbb{N}}(n -' i)$.

(54) Let given s_1, s_2, n. Suppose s_2 is summable. Then there exists F_1 such that $(\sum_{\alpha=0}^{\kappa}(s_1 (\#) s_2)(\alpha))_{\kappa \in \mathbb{N}}(n) = \sum s_2 \cdot (\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa \in \mathbb{N}}(n) - \sum F_1$ and $\operatorname{dom} F_1 = n + 1$ and for every i such that $i \in n + 1$ holds $F_1(i) = s_1(i) \cdot \sum(s_2 \uparrow ((n -' i) + 1))$.

(55) For every F_1 there exists a finite 0-sequence a_1 of $\mathbb{R}$ such that $\operatorname{dom} a_1 = \operatorname{dom} F_1$ and $|\sum F_1| \leq \sum a_1$ and for every i such that $i \in \operatorname{dom} a_1$ holds $a_1(i) = |F_1(i)|$.

(56) For every s_1 such that s_1 is summable there exists r such that $0 < r$ and for every k holds $|\sum(s_1 \uparrow k)| < r$.

(57) For all s_1, n, m such that $n \leq m$ and for every i holds $s_1(i) \geq 0$ holds $(\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}}(n) \leq (\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}}(m)$.

(58) For all s_1, s_2 such that s_1 is absolutely summable and s_2 is summable holds $s_1\,(\#)\,s_2$ is summable and $\sum(s_1\,(\#)\,s_2) = \sum s_1 \cdot \sum s_2$.

(59) If $p = F_1$, then $\sum p = \sum F_1$.

4. The Generating Function for the Catalan Numbers

Next we state the proposition

(60) Let given r. Then there exists a sequence C_2 of real numbers such that
  (i) for every n holds $C_2(n) = \mathrm{Catalan}(n+1) \cdot r^n$, and
  (ii) if $|r| < \frac{1}{4}$, then C_2 is absolutely summable and $\sum C_2 = 1 + r \cdot (\sum C_2)^2$ and $\sum C_2 = \frac{2}{1+\sqrt{1-4\cdot r}}$ and if $r \neq 0$, then $\sum C_2 = \frac{1-\sqrt{1-4\cdot r}}{2\cdot r}$.

References

[1] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(**2**):377–382, 1990.
[2] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(**1**):91–96, 1990.
[3] Patrick Braselmann and Peter Koepke. Equivalences of inconsistency and Henkin models. *Formalized Mathematics*, 13(**1**):45–48, 2005.
[4] Czesław Byliński. The complex numbers. *Formalized Mathematics*, 1(**3**):507–513, 1990.
[5] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(**1**):55–65, 1990.
[6] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(**1**):153–164, 1990.
[7] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(**1**):47–53, 1990.
[8] Dorota Częstochowska and Adam Grabowski. Catalan numbers. *Formalized Mathematics*, 12(**3**):351–353, 2004.
[9] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(**1**):165–167, 1990.
[10] Jarosław Kotowicz. Monotone real sequences. Subsequences. *Formalized Mathematics*, 1(**3**):471–475, 1990.
[11] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(**2**):269–272, 1990.
[12] Rafał Kwiatek. Factorial and Newton coefficients. *Formalized Mathematics*, 1(**5**):887–890, 1990.
[13] Takaya Nishiyama and Yasuho Mizuhara. Binary arithmetics. *Formalized Mathematics*, 4(**1**):83–86, 1993.
[14] Library Committee of the Association of Mizar Users. Binary operations on numbers. *To appear in Formalized Mathematics*.
[15] Karol Pąk. Cardinal numbers and finite sets. *Formalized Mathematics*, 13(**3**):399–406, 2005.
[16] Karol Pąk. Stirling numbers of the second kind. *Formalized Mathematics*, 13(**2**):337–345, 2005.
[17] Konrad Raczkowski. Integer and rational exponents. *Formalized Mathematics*, 2(**1**):125–130, 1991.
[18] Konrad Raczkowski and Andrzej Nędzusiak. Series. *Formalized Mathematics*, 2(**4**):449–452, 1991.
[19] Andrzej Trybulec. Subsets of complex numbers. *To appear in Formalized Mathematics*.
[20] Andrzej Trybulec. Binary operations applied to functions. *Formalized Mathematics*, 1(**2**):329–334, 1990.
[21] Andrzej Trybulec. Domains and their Cartesian products. *Formalized Mathematics*, 1(**1**):115–122, 1990.

[22] Andrzej Trybulec. Function domains and Frænkel operator. *Formalized Mathematics*, 1(**3**):495–500, 1990.
[23] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(**1**):9–11, 1990.
[24] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers. *Formalized Mathematics*, 1(**3**):445–449, 1990.
[25] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(**1**):67–71, 1990.
[26] Tetsuya Tsunetou, Grzegorz Bancerek, and Yatsuka Nakamura. Zero-based finite sequences. *Formalized Mathematics*, 9(**4**):825–829, 2001.
[27] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(**1**):73–83, 1990.
[28] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(**1**):181–186, 1990.

Received October 31, 2006