FORMALIZED MATHEMATICS
Volume 13, Number 1, 2005
University of Białystok

# On the Characterization of Collineations of the Segre Product of Strongly Connected Partial Linear Spaces[1]

Adam Naumowicz
University of Białystok

**Summary.** In this paper we characterize the automorphisms (collineations) of the Segre product of partial linear spaces. In particular, we show that if all components of the product are strongly connected, then every collineation is determined by a set of isomorphisms between its components. The formalization follows the ideas presented in the *Journal of Geometry* paper [16] by Naumowicz and Prażmowski.

MML Identifier: PENCIL_3.

The articles [20], [10], [2], [23], [22], [6], [8], [9], [19], [24], [7], [1], [11], [5], [3], [17], [21], [12], [13], [18], [4], [15], and [14] provide the terminology and notation for this paper.

## 1. Preliminaries

The following propositions are true:

(1) Let $S$ be a non empty non void topological structure, $f$ be a collineation of $S$, and $p$, $q$ be points of $S$. If $p$, $q$ are collinear, then $f(p)$, $f(q)$ are collinear.

(2) Let $I$ be a non empty set, $i$ be an element of $I$, and $A$ be a non-Trivial-yielding 1-sorted yielding many sorted set indexed by $I$. Then $A(i)$ is non trivial.

[1]This work has been partially supported by the KBN grant 4 T11C 039 24 and the FP6 IST grant TYPES No. 510096.



© 2005 University of Białystok
ISSN 1426–2630

(3) Let $S$ be a non void identifying close blocks topological structure such that $S$ is strongly connected. Then $S$ has no isolated points.

(4) Let $S$ be a non empty non void identifying close blocks topological structure. If $S$ is strongly connected, then $S$ is connected.

(5) Let $S$ be a non void non degenerated topological structure and $L$ be a block of $S$. Then there exists a point $x$ of $S$ such that $x \notin L$.

(6) Let $I$ be a non empty set and $A$ be a nonempty TopStruct-yielding many sorted set indexed by $I$. Then every point of SegreProduct $A$ is an element of the support of $A$.

(7) Let $I$ be a non empty set, $A$ be a 1-sorted yielding many sorted set indexed by $I$, and $x$ be an element of $I$. Then (the support of $A$)$(x) = \Omega_{A(x)}$.

(8) Let $I$ be a non empty set, $i$ be an element of $I$, and $A$ be a non-Trivial-yielding TopStruct-yielding many sorted set indexed by $I$. Then there exists a Segre-like non trivial-yielding many sorted subset $L$ indexed by the support of $A$ such that $\mathrm{index}(L) = i$ and $\prod L$ is a Segre coset of $A$.

(9) Let $I$ be a non empty set, $i$ be an element of $I$, $A$ be a non-Trivial-yielding TopStruct-yielding many sorted set indexed by $I$, and $p$ be a point of SegreProduct $A$. Then there exists a Segre-like non trivial-yielding many sorted subset $L$ indexed by the support of $A$ such that $\mathrm{index}(L) = i$ and $\prod L$ is a Segre coset of $A$ and $p \in \prod L$.

(10) Let $I$ be a non empty set, $A$ be a non-Trivial-yielding TopStruct-yielding many sorted set indexed by $I$, and $b$ be a Segre-like non trivial-yielding many sorted subset indexed by the support of $A$. If $\prod b$ is a Segre coset of $A$, then $b(\mathrm{index}(b)) = \Omega_{A(\mathrm{index}(b))}$.

(11) Let $I$ be a non empty set, $A$ be a non-Trivial-yielding TopStruct-yielding many sorted set indexed by $I$, and $L_1$, $L_2$ be Segre-like non trivial-yielding many sorted subsets indexed by the support of $A$. Suppose $\prod L_1$ is a Segre coset of $A$ and $\prod L_2$ is a Segre coset of $A$ and $\mathrm{index}(L_1) = \mathrm{index}(L_2)$ and $\prod L_1$ meets $\prod L_2$. Then $L_1 = L_2$.

(12) Let $I$ be a non empty set, $A$ be a PLS-yielding many sorted set indexed by $I$, $L$ be a Segre-like non trivial-yielding many sorted subset indexed by the support of $A$, and $B$ be a block of $A(\mathrm{index}(L))$. Then $\prod(L +\cdot (\mathrm{index}(L), B))$ is a block of SegreProduct $A$.

(13) Let $I$ be a non empty set, $A$ be a PLS-yielding many sorted set indexed by $I$, $i$ be an element of $I$, $p$ be a point of $A(i)$, and $L$ be a Segre-like non trivial-yielding many sorted subset indexed by the support of $A$. Suppose $i \neq \mathrm{index}(L)$. Then $L +\cdot (i, \{p\})$ is a Segre-like non trivial-yielding many sorted subset indexed by the support of $A$.

(14) Let $I$ be a non empty set, $A$ be a PLS-yielding many sorted set indexed

by $I$, $i$ be an element of $I$, $S$ be a subset of $A(i)$, and $L$ be a Segre-like non trivial-yielding many sorted subset indexed by the support of $A$. Then $\prod(L +\cdot (i, S))$ is a subset of SegreProduct $A$.

(15) Let $I$ be a non empty set, $P$ be a many sorted set indexed by $I$, and $i$ be an element of $I$. Then $\{P\}(i)$ is non empty and trivial.

(16) Let $I$ be a non empty set, $i$ be an element of $I$, $A$ be a PLS-yielding many sorted set indexed by $I$, $B$ be a block of $A(i)$, and $P$ be an element of the support of $A$. Then $\prod(\{P\} +\cdot (i, B))$ is a block of SegreProduct $A$.

(17) Let $I$ be a non empty set, $A$ be a PLS-yielding many sorted set indexed by $I$, and $p$, $q$ be points of SegreProduct $A$. Suppose $p \neq q$. Then $p$, $q$ are collinear if and only if for all many sorted sets $p_1$, $q_1$ indexed by $I$ such that $p_1 = p$ and $q_1 = q$ there exists an element $i$ of $I$ such that for all points $a$, $b$ of $A(i)$ such that $a = p_1(i)$ and $b = q_1(i)$ holds $a \neq b$ and $a$, $b$ are collinear and for every element $j$ of $I$ such that $j \neq i$ holds $p_1(j) = q_1(j)$.

(18) Let $I$ be a non empty set, $A$ be a PLS-yielding many sorted set indexed by $I$, $b$ be a Segre-like non trivial-yielding many sorted subset indexed by the support of $A$, and $x$ be a point of $A(\mathrm{index}(b))$. Then there exists a many sorted set $p$ indexed by $I$ such that $p \in \prod b$ and $\{(p +\cdot (\mathrm{index}(b), x)$ **qua** set$)\} = \prod(b +\cdot (\mathrm{index}(b), \{x\}))$.

Let $I$ be a finite non empty set and let $b_1$, $b_2$ be many sorted sets indexed by $I$. The functor ${b_1}'(b_2)$ yields a natural number and is defined by:

(Def. 1) ${b_1}'(b_2) = \overline{\overline{\{i; i \text{ ranges over elements of } I\colon b_1(i) \neq b_2(i)\}}}$.

One can prove the following proposition

(19) Let $I$ be a finite non empty set, $b_1$, $b_2$ be many sorted sets indexed by $I$, and $i$ be an element of $I$. If $b_1(i) \neq b_2(i)$, then ${b_1}'(b_2) = {b_1}'(b_2 +\cdot (i, b_1(i))) + 1$.

## 2. The Adherence of Segre Cosets

Let $I$ be a non empty set, let $A$ be a PLS-yielding many sorted set indexed by $I$, and let $B_1$, $B_2$ be Segre cosets of $A$. The predicate $B_1||B_2$ is defined as follows:

(Def. 2) For every point $x$ of SegreProduct $A$ such that $x \in B_1$ there exists a point $y$ of SegreProduct $A$ such that $y \in B_2$ and $x$, $y$ are collinear.

Next we state several propositions:

(20) Let $I$ be a non empty set, $A$ be a PLS-yielding many sorted set indexed by $I$, and $B_1$, $B_2$ be Segre cosets of $A$. Suppose $B_1||B_2$. Let $f$ be a collineation of SegreProduct $A$ and $C_1$, $C_2$ be Segre cosets of $A$. If $C_1 = f^\circ B_1$ and $C_2 = f^\circ B_2$, then $C_1||C_2$.

(21) Let $I$ be a non empty set, $A$ be a PLS-yielding many sorted set indexed by $I$, and $B_1$, $B_2$ be Segre cosets of $A$. Suppose $B_1$ misses $B_2$. Then $B_1||B_2$ if and only if for all Segre-like non trivial-yielding many sorted subsets $b_1$, $b_2$ indexed by the support of $A$ such that $B_1 = \prod b_1$ and $B_2 = \prod b_2$ holds $\mathrm{index}(b_1) = \mathrm{index}(b_2)$ and there exists an element $r$ of $I$ such that $r \neq \mathrm{index}(b_1)$ and for every element $i$ of $I$ such that $i \neq r$ holds $b_1(i) = b_2(i)$ and for all points $c_1$, $c_2$ of $A(r)$ such that $b_1(r) = \{c_1\}$ and $b_2(r) = \{c_2\}$ holds $c_1$, $c_2$ are collinear.

(22) Let $I$ be a finite non empty set and $A$ be a PLS-yielding many sorted set indexed by $I$. Suppose that for every element $i$ of $I$ holds $A(i)$ is connected. Let $i$ be an element of $I$, $p$ be a point of $A(i)$, and $b_1$, $b_2$ be Segre-like non trivial-yielding many sorted subsets indexed by the support of $A$. Suppose $\prod b_1$ is a Segre coset of $A$ and $\prod b_2$ is a Segre coset of $A$ and $b_1 = b_2 +\cdot (i, \{p\})$ and $p \notin b_2(i)$. Then there exists a finite sequence $D$ of elements of $2^{\text{the carrier of } \mathrm{SegreProduct}\, A}$ such that

(i) $D(1) = \prod b_1$,

(ii) $D(\mathrm{len}\, D) = \prod b_2$,

(iii) for every natural number $i$ such that $i \in \mathrm{dom}\, D$ holds $D(i)$ is a Segre coset of $A$, and

(iv) for every natural number $i$ such that $1 \leq i$ and $i < \mathrm{len}\, D$ and for all Segre cosets $D_1$, $D_2$ of $A$ such that $D_1 = D(i)$ and $D_2 = D(i+1)$ holds $D_1$ misses $D_2$ and $D_1||D_2$.

(23) Let $I$ be a finite non empty set and $A$ be a PLS-yielding many sorted set indexed by $I$. Suppose that for every element $i$ of $I$ holds $A(i)$ is connected. Let $B_1$, $B_2$ be Segre cosets of $A$. Suppose $B_1$ misses $B_2$. Let $b_1$, $b_2$ be Segre-like non trivial-yielding many sorted subsets indexed by the support of $A$. Suppose $B_1 = \prod b_1$ and $B_2 = \prod b_2$. Then $\mathrm{index}(b_1) = \mathrm{index}(b_2)$ if and only if there exists a finite sequence $D$ of elements of $2^{\text{the carrier of } \mathrm{SegreProduct}\, A}$ such that $D(1) = B_1$ and $D(\mathrm{len}\, D) = B_2$ and for every natural number $i$ such that $i \in \mathrm{dom}\, D$ holds $D(i)$ is a Segre coset of $A$ and for every natural number $i$ such that $1 \leq i$ and $i < \mathrm{len}\, D$ and for all Segre cosets $D_1$, $D_2$ of $A$ such that $D_1 = D(i)$ and $D_2 = D(i+1)$ holds $D_1$ misses $D_2$ and $D_1||D_2$.

(24) Let $I$ be a finite non empty set and $A$ be a PLS-yielding many sorted set indexed by $I$. Suppose that for every element $i$ of $I$ holds $A(i)$ is strongly connected. Let $f$ be a collineation of $\mathrm{SegreProduct}\, A$, $B_1$, $B_2$ be Segre cosets of $A$, and $b_1$, $b_2$, $b_3$, $b_4$ be Segre-like non trivial-yielding many sorted subsets indexed by the support of $A$. If $B_1 = \prod b_1$ and $B_2 = \prod b_2$ and $f^\circ B_1 = \prod b_3$ and $f^\circ B_2 = \prod b_4$, then if $\mathrm{index}(b_1) = \mathrm{index}(b_2)$, then $\mathrm{index}(b_3) = \mathrm{index}(b_4)$.

(25) Let $I$ be a finite non empty set and $A$ be a PLS-yielding many sorted set

indexed by $I$. Suppose that for every element $i$ of $I$ holds $A(i)$ is strongly connected. Let $f$ be a collineation of SegreProduct $A$. Then there exists a permutation $s$ of $I$ such that for all elements $i$, $j$ of $I$ holds $s(i) = j$ if and only if for every Segre coset $B_1$ of $A$ and for all Segre-like non trivial-yielding many sorted subsets $b_1$, $b_2$ indexed by the support of $A$ such that $B_1 = \prod b_1$ and $f^\circ B_1 = \prod b_2$ holds if index$(b_1) = i$, then index$(b_2) = j$.

Let $I$ be a finite non empty set and let $A$ be a PLS-yielding many sorted set indexed by $I$. Let us assume that for every element $i$ of $I$ holds $A(i)$ is strongly connected. Let $f$ be a collineation of SegreProduct $A$. The functor IndPerm$(f)$ yields a permutation of $I$ and is defined by the condition (Def. 3).

(Def. 3) Let $i$, $j$ be elements of $I$. Then (IndPerm$(f)$)$(i) = j$ if and only if for every Segre coset $B_1$ of $A$ and for all Segre-like non trivial-yielding many sorted subsets $b_1$, $b_2$ indexed by the support of $A$ such that $B_1 = \prod b_1$ and $f^\circ B_1 = \prod b_2$ holds if index$(b_1) = i$, then index$(b_2) = j$.

## 3. Canonical Embeddings and Automorphisms of Segre Product

Let $I$ be a finite non empty set and let $A$ be a PLS-yielding many sorted set indexed by $I$. Let us assume that for every element $i$ of $I$ holds $A(i)$ is strongly connected. Let $f$ be a collineation of SegreProduct $A$ and let $b_1$ be a Segre-like non trivial-yielding many sorted subset indexed by the support of $A$. Let us assume that $\prod b_1$ is a Segre coset of $A$. The functor CanEmb$(f, b_1)$ yields a map from $A$(index$(b_1)$) into $A$((IndPerm$(f)$)(index$(b_1)$)) and is defined by the conditions (Def. 4).

(Def. 4)(i) CanEmb$(f, b_1)$ is isomorphic, and

(ii) for every many sorted set $a$ indexed by $I$ such that $a$ is a point of SegreProduct $A$ and $a \in \prod b_1$ and for every many sorted set $b$ indexed by $I$ such that $b = f(a)$ holds $b$((IndPerm$(f)$)(index$(b_1)$)) = (CanEmb$(f, b_1)$)$(a$(index$(b_1)$)).

Next we state two propositions:

(26) Let $I$ be a finite non empty set and $A$ be a PLS-yielding many sorted set indexed by $I$. Suppose that for every element $i$ of $I$ holds $A(i)$ is strongly connected. Let $f$ be a collineation of SegreProduct $A$ and $B_1$, $B_2$ be Segre cosets of $A$. Suppose $B_1$ misses $B_2$ and $B_1 || B_2$. Let $b_1$, $b_2$ be Segre-like non trivial-yielding many sorted subsets indexed by the support of $A$. If $\prod b_1 = B_1$ and $\prod b_2 = B_2$, then CanEmb$(f, b_1)$ = CanEmb$(f, b_2)$.

(27) Let $I$ be a finite non empty set and $A$ be a PLS-yielding many sorted set indexed by $I$. Suppose that for every element $i$ of $I$ holds $A(i)$ is strongly connected. Let $f$ be a collineation of SegreProduct $A$ and $b_1$, $b_2$ be Segre-like non trivial-yielding many sorted subsets indexed by the support of $A$.

Suppose $\prod b_1$ is a Segre coset of $A$ and $\prod b_2$ is a Segre coset of $A$ and $\mathrm{index}(b_1) = \mathrm{index}(b_2)$. Then $\mathrm{CanEmb}(f, b_1) = \mathrm{CanEmb}(f, b_2)$.

Let $I$ be a finite non empty set and let $A$ be a PLS-yielding many sorted set indexed by $I$. Let us assume that for every element $i$ of $I$ holds $A(i)$ is strongly connected. Let $f$ be a collineation of SegreProduct $A$ and let $i$ be an element of $I$. The functor $\mathrm{CanEmb}(f, i)$ yields a map from $A(i)$ into $A((\mathrm{IndPerm}(f))(i))$ and is defined by the condition (Def. 5).

(Def. 5) Let $b$ be a Segre-like non trivial-yielding many sorted subset indexed by the support of $A$. If $\prod b$ is a Segre coset of $A$ and $\mathrm{index}(b) = i$, then $\mathrm{CanEmb}(f, i) = \mathrm{CanEmb}(f, b)$.

Next we state the proposition

(28) Let $I$ be a finite non empty set and $A$ be a PLS-yielding many sorted set indexed by $I$. Suppose that for every element $i$ of $I$ holds $A(i)$ is strongly connected. Let $f$ be a collineation of SegreProduct $A$. Then there exists a permutation $s$ of $I$ and there exists a function yielding many sorted set $B$ indexed by $I$ such that for every element $i$ of $I$ holds

(i) $B(i)$ is a map from $A(i)$ into $A(s(i))$,

(ii) for every map $m$ from $A(i)$ into $A(s(i))$ such that $m = B(i)$ holds $m$ is isomorphic, and

(iii) for every point $p$ of SegreProduct $A$ and for every many sorted set $a$ indexed by $I$ such that $a = p$ and for every many sorted set $b$ indexed by $I$ such that $b = f(p)$ and for every map $m$ from $A(i)$ into $A(s(i))$ such that $m = B(i)$ holds $b(s(i)) = m(a(i))$.

## References


[1] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(**2**):377–382, 1990.
[2] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(**1**):41–46, 1990.
[3] Grzegorz Bancerek. König's theorem. *Formalized Mathematics*, 1(**3**):589–593, 1990.
[4] Grzegorz Bancerek. The "way-below" relation. *Formalized Mathematics*, 6(**1**):169–176, 1997.
[5] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(**1**):107–114, 1990.
[6] Józef Białas. Group and field definitions. *Formalized Mathematics*, 1(**3**):433–439, 1990.
[7] Józef Białas. Properties of fields. *Formalized Mathematics*, 1(**5**):807–812, 1990.
[8] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(**1**):55–65, 1990.
[9] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(**1**):153–164, 1990.
[10] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(**1**):47–53, 1990.
[11] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(**1**):165–167, 1990.
[12] Artur Korniłowicz. Some basic properties of many sorted sets. *Formalized Mathematics*, 5(**3**):395–399, 1996.
[13] Beata Madras. Product of family of universal algebras. *Formalized Mathematics*, 4(**1**):103–108, 1993.
[14] Adam Naumowicz. On cosets in Segre's product of partial linear spaces. *Formalized Mathematics*, 9(**4**):795–800, 2001.

[15] Adam Naumowicz. On Segre's product of partial line spaces. *Formalized Mathematics*, 9(**2**):383–390, 2001.
[16] Adam Naumowicz and Krzysztof Prażmowski. On Segre's product of partial line spaces and spaces of pencils. *Journal of Geometry*, 71(**1**):128–143, 2001.
[17] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(**1**):223–230, 1990.
[18] Piotr Rudnicki and Andrzej Trybulec. Multivariate polynomials with arbitrary number of variables. *Formalized Mathematics*, 9(**1**):95–110, 2001.
[19] Andrzej Trybulec. Binary operations applied to functions. *Formalized Mathematics*, 1(**2**):329–334, 1990.
[20] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(**1**):9–11, 1990.
[21] Andrzej Trybulec. Many-sorted sets. *Formalized Mathematics*, 4(**1**):15–22, 1993.
[22] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(**1**):67–71, 1990.
[23] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(**1**):73–83, 1990.
[24] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(**1**):181–186, 1990.

*Received October 18, 2004*