FORMALIZED MATHEMATICS
Volume 13, Number 1, 2005
University of Białystok

A Theory of Matrices of Complex Elements

Wenpai Chang
Shinshu University
Nagano

Hiroshi Yamazaki
Shinshu University
Nagano

Yatsuka Nakamura
Shinshu University
Nagano

Summary. A concept of "Matrix of Complex" is defined here. Addition, subtraction, scalar multiplication and product are introduced using correspondent definitions of "Matrix of Field". Many equations for such operations consist of a case of "Matrix of Field". A calculation method of product of matrices is shown using a finite sequence of Complex in the last theorem.

MML Identifier: MATRIX_5.

The articles [11], [14], [1], [4], [2], [15], [6], [10], [9], [3], [8], [7], [13], [12], and [5] provide the terminology and notation for this paper.

The following two propositions are true:

(1) $1 = 1_{\mathbb{C}_{\mathrm{F}}}$.

(2) $0_{\mathbb{C}_{\mathrm{F}}} = 0$.

Let A be a matrix over $\mathbb{C}$. The functor $A_{\mathbb{C}_{\mathrm{F}}}$ yields a matrix over $\mathbb{C}_{\mathrm{F}}$ and is defined by:

(Def. 1) $A_{\mathbb{C}_{\mathrm{F}}} = A$.

Let A be a matrix over $\mathbb{C}_{\mathrm{F}}$. The functor $A_{\mathbb{C}}$ yielding a matrix over $\mathbb{C}$ is defined by:

(Def. 2) $A_{\mathbb{C}} = A$.

We now state four propositions:

(3) For all matrices A, B over $\mathbb{C}$ such that $A_{\mathbb{C}_{\mathrm{F}}} = B_{\mathbb{C}_{\mathrm{F}}}$ holds $A = B$.

(4) For all matrices A, B over $\mathbb{C}_{\mathrm{F}}$ such that $A_{\mathbb{C}} = B_{\mathbb{C}}$ holds $A = B$.

(5) For every matrix A over $\mathbb{C}$ holds $A = (A_{\mathbb{C}_{\mathrm{F}}})_{\mathbb{C}}$.

(6) For every matrix A over $\mathbb{C}_{\mathrm{F}}$ holds $A = (A_{\mathbb{C}})_{\mathbb{C}_{\mathrm{F}}}$.

Let A, B be matrices over $\mathbb{C}$. The functor $A + B$ yielding a matrix over $\mathbb{C}$ is defined as follows:

© 2005 University of Białystok
ISSN 1426–2630

(Def. 3) $A + B = (A_{\mathbb{C}_\mathrm{F}} + B_{\mathbb{C}_\mathrm{F}})_{\mathbb{C}}$.

Let A be a matrix over $\mathbb{C}$. The functor $-A$ yielding a matrix over $\mathbb{C}$ is defined as follows:

(Def. 4) $-A = (-A_{\mathbb{C}_\mathrm{F}})_{\mathbb{C}}$.

Let A, B be matrices over $\mathbb{C}$. The functor $A - B$ yields a matrix over $\mathbb{C}$ and is defined as follows:

(Def. 5) $A - B = (A_{\mathbb{C}_\mathrm{F}} - B_{\mathbb{C}_\mathrm{F}})_{\mathbb{C}}$.

Let A, B be matrices over $\mathbb{C}$. The functor $A \cdot B$ yielding a matrix over $\mathbb{C}$ is defined as follows:

(Def. 6) $A \cdot B = (A_{\mathbb{C}_\mathrm{F}} \cdot B_{\mathbb{C}_\mathrm{F}})_{\mathbb{C}}$.

Let x be a complex number and let A be a matrix over $\mathbb{C}$. The functor $x \cdot A$ yielding a matrix over $\mathbb{C}$ is defined as follows:

(Def. 7) For every element e_1 of $\mathbb{C}_\mathrm{F}$ such that $e_1 = x$ holds $x \cdot A = (e_1 \cdot A_{\mathbb{C}_\mathrm{F}})_{\mathbb{C}}$.

One can prove the following propositions:

(7) For every matrix A over $\mathbb{C}$ holds $\operatorname{len} A = \operatorname{len}(A_{\mathbb{C}_\mathrm{F}})$ and $\operatorname{width} A = \operatorname{width}(A_{\mathbb{C}_\mathrm{F}})$.

(8) For every matrix A over $\mathbb{C}_\mathrm{F}$ holds $\operatorname{len} A = \operatorname{len}(A_{\mathbb{C}})$ and $\operatorname{width} A = \operatorname{width}(A_{\mathbb{C}})$.

(9) For every matrix M over $\mathbb{C}$ such that $\operatorname{len} M > 0$ holds $--M = M$.

(10) For every field K and for every matrix M over K holds $1_K \cdot M = M$.

(11) For every matrix M over $\mathbb{C}$ holds $1 \cdot M = M$.

(12) For every field K and for all elements a, b of K and for every matrix M over K holds $a \cdot (b \cdot M) = (a \cdot b) \cdot M$.

(13) For every field K and for all elements a, b of K and for every matrix M over K holds $(a + b) \cdot M = a \cdot M + b \cdot M$.

(14) For every matrix M over $\mathbb{C}$ holds $M + M = 2 \cdot M$.

(15) For every matrix M over $\mathbb{C}$ holds $M + M + M = 3 \cdot M$.

Let n, m be natural numbers. The functor $\begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}^{n \times m}_{\mathbb{C}}$ yields a matrix over $\mathbb{C}$ and is defined by:

(Def. 8) $\begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}^{n \times m}_{\mathbb{C}} = \left(\begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}^{n \times m}_{\mathbb{C}_\mathrm{F}}\right)_{\mathbb{C}}$.

One can prove the following propositions:

(16) For all natural numbers n, m holds $\begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}^{n\times m}_{\mathbb{C}\quad \mathbb{C}_{\mathrm{F}}} = \begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}^{n\times m}_{\mathbb{C}_{\mathrm{F}}}$.

(17) For every matrix M over $\mathbb{C}$ such that $\operatorname{len} M > 0$ holds $M + -M = \begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}^{(\operatorname{len} M)\times(\operatorname{width} M)}_{\mathbb{C}}$.

(18) For every matrix M over $\mathbb{C}$ such that $\operatorname{len} M > 0$ holds $M - M = \begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}^{(\operatorname{len} M)\times(\operatorname{width} M)}_{\mathbb{C}}$.

(19) For all matrices M_1, M_2, M_3 over $\mathbb{C}$ such that $\operatorname{len} M_1 = \operatorname{len} M_2$ and $\operatorname{len} M_2 = \operatorname{len} M_3$ and $\operatorname{width} M_1 = \operatorname{width} M_2$ and $\operatorname{width} M_2 = \operatorname{width} M_3$ and $\operatorname{len} M_1 > 0$ and $M_1 + M_3 = M_2 + M_3$ holds $M_1 = M_2$.

(20) For all matrices M_1, M_2 over $\mathbb{C}$ such that $\operatorname{len} M_2 > 0$ holds $M_1 - -M_2 = M_1 + M_2$.

(21) For all matrices M_1, M_2 over $\mathbb{C}$ such that $\operatorname{len} M_1 = \operatorname{len} M_2$ and $\operatorname{width} M_1 = \operatorname{width} M_2$ and $\operatorname{len} M_1 > 0$ and $M_1 = M_1 + M_2$ holds $M_2 = \begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}^{(\operatorname{len} M_1)\times(\operatorname{width} M_1)}_{\mathbb{C}}$.

(22) For all matrices M_1, M_2 over $\mathbb{C}$ such that $\operatorname{len} M_1 = \operatorname{len} M_2$ and $\operatorname{width} M_1 = \operatorname{width} M_2$ and $\operatorname{len} M_1 > 0$ and $M_1 - M_2 = \begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}^{(\operatorname{len} M_1)\times(\operatorname{width} M_1)}_{\mathbb{C}}$ holds $M_1 = M_2$.

(23) For all matrices M_1, M_2 over $\mathbb{C}$ such that $\operatorname{len} M_1 = \operatorname{len} M_2$ and $\operatorname{width} M_1 = \operatorname{width} M_2$ and $\operatorname{len} M_1 > 0$ and $M_1 + M_2 = \begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}^{(\operatorname{len} M_1)\times(\operatorname{width} M_1)}_{\mathbb{C}}$ holds $M_2 = -M_1$.

(24) For all natural numbers n, m such that $n > 0$ holds

$$-\begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}_{\mathbb{C}}^{n\times m} = \begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}_{\mathbb{C}}^{n\times m}.$$

(25) For all matrices M_1, M_2 over $\mathbb{C}$ such that $\operatorname{len} M_1 = \operatorname{len} M_2$ and $\operatorname{width} M_1 = \operatorname{width} M_2$ and $\operatorname{len} M_1 > 0$ and $M_2 - M_1 = M_2$ holds $M_1 = \begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}_{\mathbb{C}}^{(\operatorname{len} M_1)\times(\operatorname{width} M_1)}$.

(26) For all matrices M_1, M_2 over $\mathbb{C}$ such that $\operatorname{len} M_1 = \operatorname{len} M_2$ and $\operatorname{width} M_1 = \operatorname{width} M_2$ and $\operatorname{len} M_1 > 0$ holds $M_1 = M_1 - (M_2 - M_2)$.

(27) For all matrices M_1, M_2 over $\mathbb{C}$ such that $\operatorname{len} M_1 = \operatorname{len} M_2$ and $\operatorname{width} M_1 = \operatorname{width} M_2$ and $\operatorname{len} M_1 > 0$ holds $-(M_1 + M_2) = -M_1 + -M_2$.

(28) For all matrices M_1, M_2 over $\mathbb{C}$ such that $\operatorname{len} M_1 = \operatorname{len} M_2$ and $\operatorname{width} M_1 = \operatorname{width} M_2$ and $\operatorname{len} M_1 > 0$ holds $M_1 - (M_1 - M_2) = M_2$.

(29) For all matrices M_1, M_2, M_3 over $\mathbb{C}$ such that $\operatorname{len} M_1 = \operatorname{len} M_2$ and $\operatorname{len} M_2 = \operatorname{len} M_3$ and $\operatorname{width} M_1 = \operatorname{width} M_2$ and $\operatorname{width} M_2 = \operatorname{width} M_3$ and $\operatorname{len} M_1 > 0$ and $M_1 - M_3 = M_2 - M_3$ holds $M_1 = M_2$.

(30) For all matrices M_1, M_2, M_3 over $\mathbb{C}$ such that $\operatorname{len} M_1 = \operatorname{len} M_2$ and $\operatorname{len} M_2 = \operatorname{len} M_3$ and $\operatorname{width} M_1 = \operatorname{width} M_2$ and $\operatorname{width} M_2 = \operatorname{width} M_3$ and $\operatorname{len} M_1 > 0$ and $M_3 - M_1 = M_3 - M_2$ holds $M_1 = M_2$.

(31) For all matrices M_1, M_2, M_3 over $\mathbb{C}$ such that $\operatorname{len} M_2 = \operatorname{len} M_3$ and $\operatorname{width} M_2 = \operatorname{width} M_3$ and $\operatorname{width} M_1 = \operatorname{len} M_2$ and $\operatorname{len} M_1 > 0$ and $\operatorname{len} M_2 > 0$ holds $M_1 \cdot (M_2 + M_3) = M_1 \cdot M_2 + M_1 \cdot M_3$.

(32) For all matrices M_1, M_2, M_3 over $\mathbb{C}$ such that $\operatorname{len} M_2 = \operatorname{len} M_3$ and $\operatorname{width} M_2 = \operatorname{width} M_3$ and $\operatorname{len} M_1 = \operatorname{width} M_2$ and $\operatorname{len} M_2 > 0$ and $\operatorname{len} M_1 > 0$ holds $(M_2 + M_3) \cdot M_1 = M_2 \cdot M_1 + M_3 \cdot M_1$.

(33) For all matrices M_1, M_2 over $\mathbb{C}$ such that $\operatorname{len} M_1 = \operatorname{len} M_2$ and $\operatorname{width} M_1 = \operatorname{width} M_2$ holds $M_1 + M_2 = M_2 + M_1$.

(34) For all matrices M_1, M_2, M_3 over $\mathbb{C}$ such that $\operatorname{len} M_1 = \operatorname{len} M_2$ and $\operatorname{len} M_1 = \operatorname{len} M_3$ and $\operatorname{width} M_1 = \operatorname{width} M_2$ and $\operatorname{width} M_1 = \operatorname{width} M_3$ holds $(M_1 + M_2) + M_3 = M_1 + (M_2 + M_3)$.

(35) For every matrix M over $\mathbb{C}$ such that $\operatorname{len} M > 0$ holds $M + \begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}_{\mathbb{C}}^{(\operatorname{len} M)\times(\operatorname{width} M)} = M$.

(36) Let K be a field, b be an element of K, and M_1, M_2 be matrices over K. If $\operatorname{len} M_1 = \operatorname{len} M_2$ and $\operatorname{width} M_1 = \operatorname{width} M_2$ and $\operatorname{len} M_1 > 0$, then $b \cdot (M_1 + M_2) = b \cdot M_1 + b \cdot M_2$.

(37) Let M_1, M_2 be matrices over $\mathbb{C}$ and a be a complex number. If $\operatorname{len} M_1 = \operatorname{len} M_2$ and $\operatorname{width} M_1 = \operatorname{width} M_2$ and $\operatorname{len} M_1 > 0$, then $a \cdot (M_1 + M_2) = a \cdot M_1 + a \cdot M_2$.

(38) For every field K and for all matrices M_1, M_2 over K such that $\operatorname{width} M_1 = \operatorname{len} M_2$ and $\operatorname{len} M_1 > 0$ and $\operatorname{len} M_2 > 0$ holds $\begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}_K^{(\operatorname{len} M_1)\times(\operatorname{width} M_1)} \cdot M_2 = \begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}_K^{(\operatorname{len} M_1)\times(\operatorname{width} M_2)}$.

(39) For all matrices M_1, M_2 over $\mathbb{C}$ such that $\operatorname{width} M_1 = \operatorname{len} M_2$ and $\operatorname{len} M_1 > 0$ and $\operatorname{len} M_2 > 0$ holds $\begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}_{\mathbb{C}}^{(\operatorname{len} M_1)\times(\operatorname{width} M_1)} \cdot M_2 = \begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}_{\mathbb{C}}^{(\operatorname{len} M_1)\times(\operatorname{width} M_2)}$.

(40) For every field K and for every matrix M_1 over K such that $\operatorname{len} M_1 > 0$ holds $0_K \cdot M_1 = \begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}_K^{(\operatorname{len} M_1)\times(\operatorname{width} M_1)}$.

(41) For every matrix M_1 over $\mathbb{C}$ such that $\operatorname{len} M_1 > 0$ holds $0 \cdot M_1 = \begin{pmatrix} 0 & \ldots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \ldots & 0 \end{pmatrix}_{\mathbb{C}}^{(\operatorname{len} M_1)\times(\operatorname{width} M_1)}$.

Let s be a finite sequence of elements of $\mathbb{C}$ and let k be a natural number. Then $s(k)$ is an element of $\mathbb{C}$.

We now state the proposition

(42) Let i, j be natural numbers and M_1, M_2 be matrices over $\mathbb{C}$. Suppose $\operatorname{len} M_1 > 0$ and $\operatorname{len} M_2 > 0$ and $\operatorname{width} M_1 = \operatorname{len} M_2$ and $1 \leq i$ and $i \leq \operatorname{len} M_1$ and $1 \leq j$ and $j \leq \operatorname{width} M_2$. Then there exists a finite sequence s of elements of $\mathbb{C}$ such that $\operatorname{len} s = \operatorname{len} M_2$ and $s(1) = (M_1 \circ (i, 1)) \cdot (M_2 \circ (1, j))$ and for every natural number k such that $1 \leq k$ and $k < \operatorname{len} M_2$ holds $s(k+1) = s(k) + (M_1 \circ (i, k+1)) \cdot (M_2 \circ (k+1, j))$.

References

[1] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(**1**):91–96, 1990.

[2] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(**1**):107–114, 1990.

[3] Czesław Byliński. The complex numbers. *Formalized Mathematics*, 1(**3**):507–513, 1990.

[4] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(**1**):55–65, 1990.
[5] Library Committee. Binary operations on numbers. *To appear in Formalized Mathematics*.
[6] Katarzyna Jankowska. Matrices. Abelian group of matrices. *Formalized Mathematics*, 2(**4**):475–480, 1991.
[7] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Formalized Mathematics*, 1(**2**):335–342, 1990.
[8] Anna Justyna Milewska. The field of complex numbers. *Formalized Mathematics*, 9(**2**):265–269, 2001.
[9] Yatsuka Nakamura and Hiroshi Yamazaki. Calculation of matrices of field elements. Part I. *Formalized Mathematics*, 11(**4**):385–391, 2003.
[10] Andrzej Trybulec. Subsets of complex numbers. *To appear in Formalized Mathematics*.
[11] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(**1**):9–11, 1990.
[12] Wojciech A. Trybulec. Groups. *Formalized Mathematics*, 1(**5**):821–827, 1990.
[13] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(**2**):291–296, 1990.
[14] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(**1**):67–71, 1990.
[15] Katarzyna Zawadzka. The product and the determinant of matrices with entries in a field. *Formalized Mathematics*, 4(**1**):1–8, 1993.

Received December 10, 2004