FORMALIZED MATHEMATICS
Volume 13, Number 2, Pages 199–219
University of Białystok, 2005

The Fashoda Meet Theorem for Rectangles

Yatsuka Nakamura
Shinshu University
Nagano, Japan

Andrzej Trybulec
Institute of Computer Science
University of Białystok
Sosnowa 64, 15-887 Białystok
Poland

Summary. Here, the so called Fashoda Meet Theorem is proven in the case of rectangles. All cases of proper location of arcs are listed up, and it is shown that the theorem is valid in each case. Such a list of cases will be useful when one wants to apply the theorem.

MML identifier: JGRAPH_7, version: 7.5.01 4.39.921

The articles [1], [6], [15], [17], [5], [2], [3], [16], [7], [14], [13], [10], [11], [8], [4], [9], and [12] provide the notation and terminology for this paper.

One can prove the following propositions:

(1) For all real numbers a, b, d and for every point p of $\mathcal{E}^2_{\mathrm{T}}$ such that $a < b$ and $p_{\mathbf{2}} = d$ and $a \leq p_{\mathbf{1}}$ and $p_{\mathbf{1}} \leq b$ holds $p \in \mathcal{L}([a, d], [b, d])$.

(2) Let n be a natural number, P be a subset of $\mathcal{E}^n_{\mathrm{T}}$, and p_1, p_2 be points of $\mathcal{E}^n_{\mathrm{T}}$. Suppose P is an arc from p_1 to p_2. Then there exists a map f from $\mathbb{I}$ into $\mathcal{E}^n_{\mathrm{T}}$ such that f is continuous and one-to-one and $\operatorname{rng} f = P$ and $f(0) = p_1$ and $f(1) = p_2$.

(3) Let p_1, p_2 be points of $\mathcal{E}^2_{\mathrm{T}}$ and b, c, d be real numbers. If $(p_1)_{\mathbf{1}} < b$ and $(p_1)_{\mathbf{1}} = (p_2)_{\mathbf{1}}$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$, then $p_1 \leq_{\mathrm{Rectangle}((p_1)_{\mathbf{1}},b,c,d)} p_2$.

(4) Let p_1, p_2 be points of $\mathcal{E}^2_{\mathrm{T}}$ and b, c be real numbers. Suppose $(p_1)_{\mathbf{1}} < b$ and $c < (p_2)_{\mathbf{2}}$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq (p_2)_{\mathbf{2}}$ and $(p_1)_{\mathbf{1}} \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$. Then $p_1 \leq_{\mathrm{Rectangle}((p_1)_{\mathbf{1}},b,c,(p_2)_{\mathbf{2}})} p_2$.

(5) Let p_1, p_2 be points of $\mathcal{E}^2_{\mathrm{T}}$ and c, d be real numbers. Suppose $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $c < d$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $c \leq (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$. Then $p_1 \leq_{\mathrm{Rectangle}((p_1)_{\mathbf{1}},(p_2)_{\mathbf{1}},c,d)} p_2$.

© 2005 University of Białystok
ISSN 1426–2630

(6) Let p_1, p_2 be points of $\mathcal{E}^2_{\mathrm{T}}$ and b, d be real numbers. If $(p_2)_{\mathbf{2}} < d$ and $(p_2)_{\mathbf{2}} \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$, then $p_1 \leq_{\mathrm{Rectangle}((p_1)_{\mathbf{1}},b,(p_2)_{\mathbf{2}},d)} p_2$.

(7) Let p_1, p_2 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$. Then $p_1 \leq_{\mathrm{Rectangle}(a,b,c,d)} p_2$.

(8) Let p_1, p_2 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{1}} = b$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} \leq b$ and $c \leq (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$. Then $p_1 \leq_{\mathrm{Rectangle}(a,b,c,d)} p_2$.

(9) Let p_1, p_2 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} \leq b$ and $a < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$. Then $p_1 \leq_{\mathrm{Rectangle}(a,b,c,d)} p_2$.

(10) Let p_1, p_2 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = b$ and $(p_2)_{\mathbf{1}} = b$ and $c \leq (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} < (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$. Then $p_1 \leq_{\mathrm{Rectangle}(a,b,c,d)} p_2$.

(11) Let p_1, p_2 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = b$ and $(p_2)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$. Then $p_1 \leq_{\mathrm{Rectangle}(a,b,c,d)} p_2$.

(12) Let p_1, p_2 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = c$ and $(p_2)_{\mathbf{2}} = c$ and $a < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} \leq b$. Then $p_1 \leq_{\mathrm{Rectangle}(a,b,c,d)} p_2$.

(13) Let p_1, p_2 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{1}} = b$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} \leq b$ and $c \leq (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$. Then $p_1 \leq_{\mathrm{Rectangle}(a,b,c,d)} p_2$.

(14) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{1}} = a$ and $(p_4)_{\mathbf{1}} = a$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} < (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \leq d$. Then p_1, p_2, p_3, p_4 are in this order on $\mathrm{Rectangle}(a, b, c, d)$.

(15) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{1}} = a$ and $(p_4)_{\mathbf{2}} = d$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $a \leq (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on $\mathrm{Rectangle}(a, b, c, d)$.

(16) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{1}} = a$ and $(p_4)_{\mathbf{1}} = b$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \leq d$. Then p_1, p_2, p_3, p_4 are in this order on $\mathrm{Rectangle}(a, b, c, d)$.

(17) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{1}} = a$ and

$(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(18) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{2}} = d$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a \leq (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(19) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{1}} = b$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a \leq (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \leq d$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(20) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a \leq (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(21) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{1}} = b$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(22) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $c \leq (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(23) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(24) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{2}} = d$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(25) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{1}} = b$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and

$(p_3)_{\mathbf{1}} \leq b$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \leq d$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(26) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(27) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{1}} = b$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(28) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$. Suppose $(p_1)_{\mathbf{1}} \neq (p_3)_{\mathbf{1}}$ and $(p_4)_{\mathbf{2}} \neq (p_2)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq (p_2)_{\mathbf{2}}$ and $(p_1)_{\mathbf{1}} \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq (p_3)_{\mathbf{1}}$ and $(p_4)_{\mathbf{2}} \leq (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq (p_2)_{\mathbf{2}}$ and $(p_1)_{\mathbf{1}} < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq (p_3)_{\mathbf{1}}$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle$((p_1)_{\mathbf{1}}, (p_3)_{\mathbf{1}}, (p_4)_{\mathbf{2}}, (p_2)_{\mathbf{2}})$.

(29) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(30) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{1}} = b$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $d \geq (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} > (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} > (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \geq c$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(31) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $d \geq (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} > (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \geq c$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(32) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $c \leq (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(33) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = c$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on

Rectangle(a, b, c, d).

(34) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{2}} = d$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(35) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{1}} = b$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \leq d$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(36) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(37) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{1}} = b$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(38) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $c \leq (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(39) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(40) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{1}} = b$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} \leq b$ and $d \geq (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} > (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} > (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \geq c$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(41) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} \leq b$ and $d \geq (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} > (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \geq c$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on Rectangle(a, b, c, d).

(42) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} \leq b$ and $c \leq (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on $\mathrm{Rectangle}(a, b, c, d)$.

(43) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = c$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} \leq b$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on $\mathrm{Rectangle}(a, b, c, d)$.

(44) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = b$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{1}} = b$ and $d \geq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} > (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} > (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} > (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \geq c$. Then p_1, p_2, p_3, p_4 are in this order on $\mathrm{Rectangle}(a, b, c, d)$.

(45) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = b$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{2}} = c$ and $d \geq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} > (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} > (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \geq c$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$. Then p_1, p_2, p_3, p_4 are in this order on $\mathrm{Rectangle}(a, b, c, d)$.

(46) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = b$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $d \geq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} > (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \geq c$ and $b \geq (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} > (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} > a$. Then p_1, p_2, p_3, p_4 are in this order on $\mathrm{Rectangle}(a, b, c, d)$.

(47) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = b$ and $(p_2)_{\mathbf{2}} = c$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $b \geq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} > (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} > (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} > a$. Then p_1, p_2, p_3, p_4 are in this order on $\mathrm{Rectangle}(a, b, c, d)$.

(48) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$ and a, b, c, d be real numbers. Suppose $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = c$ and $(p_2)_{\mathbf{2}} = c$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $b \geq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} > (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} > (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} > (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} > a$. Then p_1, p_2, p_3, p_4 are in this order on $\mathrm{Rectangle}(a, b, c, d)$.

(49) Let A, B, C, D be real numbers and h, g be maps from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose $A > 0$ and $C > 0$ and $h = \mathrm{AffineMap}(A, B, C, D)$ and $g = \mathrm{AffineMap}(\frac{1}{A}, -\frac{B}{A}, \frac{1}{C}, -\frac{D}{C})$. Then $g = h^{-1}$ and $h = g^{-1}$.

(50) Let A, B, C, D be real numbers and h be a map from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose $A > 0$ and $C > 0$ and $h = \mathrm{AffineMap}(A, B, C, D)$. Then h is a homeomorphism and for all points p_1, p_2 of $\mathcal{E}^2_{\mathrm{T}}$ such that $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ holds $h(p_1)_{\mathbf{1}} < h(p_2)_{\mathbf{1}}$.

(51) Let A, B, C, D be real numbers and h be a map from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose $A > 0$ and $C > 0$ and $h = \mathrm{AffineMap}(A, B, C, D)$. Then h is a homeomorphism and for all points p_1, p_2 of $\mathcal{E}^2_{\mathrm{T}}$ such that $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ holds $h(p_1)_{\mathbf{2}} < h(p_2)_{\mathbf{2}}$.

(52) Let a, b, c, d be real numbers, h be a map from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$, and f be a map from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose $a < b$ and $c < d$ and $h = \mathrm{AffineMap}(\frac{2}{b-a}, -\frac{b+a}{b-a}, \frac{2}{d-c}, -\frac{d+c}{d-c})$ and $\mathrm{rng}\, f \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$. Then $\mathrm{rng}(h \cdot f) \subseteq \mathrm{ClosedInsideOfRectangle}(-1, 1, -1, 1)$.

(53) Let a, b, c, d be real numbers, h be a map from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$, and f be a map from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose $a < b$ and $c < d$ and $h = \mathrm{AffineMap}(\frac{2}{b-a}, -\frac{b+a}{b-a}, \frac{2}{d-c}, -\frac{d+c}{d-c})$ and f is continuous and one-to-one. Then $h \cdot f$ is continuous and one-to-one.

(54) Let a, b, c, d be real numbers, h be a map from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$, f be a map from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$, and O be a point of $\mathbb{I}$. Suppose $a < b$ and $c < d$ and $h = \mathrm{AffineMap}(\frac{2}{b-a}, -\frac{b+a}{b-a}, \frac{2}{d-c}, -\frac{d+c}{d-c})$ and $f(O)_{\mathbf{1}} = a$. Then $(h \cdot f)(O)_{\mathbf{1}} = -1$.

(55) Let a, b, c, d be real numbers, h be a map from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$, f be a map from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$, and I be a point of $\mathbb{I}$. Suppose $a < b$ and $c < d$ and $h = \mathrm{AffineMap}(\frac{2}{b-a}, -\frac{b+a}{b-a}, \frac{2}{d-c}, -\frac{d+c}{d-c})$ and $f(I)_{\mathbf{2}} = d$. Then $(h \cdot f)(I)_{\mathbf{2}} = 1$.

(56) Let a, b, c, d be real numbers, h be a map from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$, f be a map from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$, and I be a point of $\mathbb{I}$. Suppose $a < b$ and $c < d$ and $h = \mathrm{AffineMap}(\frac{2}{b-a}, -\frac{b+a}{b-a}, \frac{2}{d-c}, -\frac{d+c}{d-c})$ and $f(I)_{\mathbf{1}} = b$. Then $(h \cdot f)(I)_{\mathbf{1}} = 1$.

(57) Let a, b, c, d be real numbers, h be a map from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$, f be a map from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$, and I be a point of $\mathbb{I}$. Suppose $a < b$ and $c < d$ and $h = \mathrm{AffineMap}(\frac{2}{b-a}, -\frac{b+a}{b-a}, \frac{2}{d-c}, -\frac{d+c}{d-c})$ and $f(I)_{\mathbf{2}} = c$. Then $(h \cdot f)(I)_{\mathbf{2}} = -1$.

(58) Let a, b, c, d be real numbers, h be a map from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$, f be a map from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$, and O, I be points of $\mathbb{I}$. Suppose $a < b$ and $c < d$ and $h = \mathrm{AffineMap}(\frac{2}{b-a}, -\frac{b+a}{b-a}, \frac{2}{d-c}, -\frac{d+c}{d-c})$ and $c \leq f(O)_{\mathbf{2}}$ and $f(O)_{\mathbf{2}} < f(I)_{\mathbf{2}}$ and $f(I)_{\mathbf{2}} \leq d$. Then $-1 \leq (h \cdot f)(O)_{\mathbf{2}}$ and $(h \cdot f)(O)_{\mathbf{2}} < (h \cdot f)(I)_{\mathbf{2}}$ and $(h \cdot f)(I)_{\mathbf{2}} \leq 1$.

(59) Let a, b, c, d be real numbers, h be a map from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$, f be a map from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$, and O, I be points of $\mathbb{I}$. Suppose $a < b$ and $c < d$ and $h = \mathrm{AffineMap}(\frac{2}{b-a}, -\frac{b+a}{b-a}, \frac{2}{d-c}, -\frac{d+c}{d-c})$ and $c \leq f(O)_{\mathbf{2}}$ and $f(O)_{\mathbf{2}} \leq d$ and $a \leq f(I)_{\mathbf{1}}$ and $f(I)_{\mathbf{1}} \leq b$. Then $-1 \leq (h \cdot f)(O)_{\mathbf{2}}$ and $(h \cdot f)(O)_{\mathbf{2}} \leq 1$ and $-1 \leq (h \cdot f)(I)_{\mathbf{1}}$ and $(h \cdot f)(I)_{\mathbf{1}} \leq 1$.

(60) Let a, b, c, d be real numbers, h be a map from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$, f be a map from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$, and O, I be points of $\mathbb{I}$. Suppose $a < b$ and $c < d$ and $h = \mathrm{AffineMap}(\frac{2}{b-a}, -\frac{b+a}{b-a}, \frac{2}{d-c}, -\frac{d+c}{d-c})$ and $c \leq f(O)_{\mathbf{2}}$ and $f(O)_{\mathbf{2}} \leq d$ and $c \leq f(I)_{\mathbf{2}}$ and $f(I)_{\mathbf{2}} \leq d$. Then $-1 \leq (h \cdot f)(O)_{\mathbf{2}}$ and $(h \cdot f)(O)_{\mathbf{2}} \leq 1$ and $-1 \leq (h \cdot f)(I)_{\mathbf{2}}$ and $(h \cdot f)(I)_{\mathbf{2}} \leq 1$.

(61) Let a, b, c, d be real numbers, h be a map from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$, f be a map from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$, and O, I be points of $\mathbb{I}$. Suppose $a < b$ and $c < d$ and $h = \mathrm{AffineMap}(\frac{2}{b-a}, -\frac{b+a}{b-a}, \frac{2}{d-c}, -\frac{d+c}{d-c})$ and $c \leq f(O)_{\mathbf{2}}$ and $f(O)_{\mathbf{2}} \leq d$ and $a < f(I)_{\mathbf{1}}$ and $f(I)_{\mathbf{1}} \leq b$. Then $-1 \leq (h \cdot f)(O)_{\mathbf{2}}$ and $(h \cdot f)(O)_{\mathbf{2}} \leq 1$ and $-1 < (h \cdot f)(I)_{\mathbf{1}}$ and $(h \cdot f)(I)_{\mathbf{1}} \leq 1$.

(62) Let a, b, c, d be real numbers, h be a map from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$, f be a map from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$, and O, I be points of $\mathbb{I}$. Suppose $a < b$ and $c < d$ and $h = \mathrm{AffineMap}(\frac{2}{b-a}, -\frac{b+a}{b-a}, \frac{2}{d-c}, -\frac{d+c}{d-c})$ and $a \leq f(O)_{\mathbf{1}}$ and $f(O)_{\mathbf{1}} < f(I)_{\mathbf{1}}$ and $f(I)_{\mathbf{1}} \leq b$. Then $-1 \leq (h \cdot f)(O)_{\mathbf{1}}$ and $(h \cdot f)(O)_{\mathbf{1}} < (h \cdot f)(I)_{\mathbf{1}}$ and $(h \cdot f)(I)_{\mathbf{1}} \leq 1$.

(63) Let a, b, c, d be real numbers, h be a map from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$, f be a map from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$, and O, I be points of $\mathbb{I}$. Suppose $a < b$ and $c < d$ and $h = \mathrm{AffineMap}(\frac{2}{b-a}, -\frac{b+a}{b-a}, \frac{2}{d-c}, -\frac{d+c}{d-c})$ and $a \leq f(O)_{\mathbf{1}}$ and $f(O)_{\mathbf{1}} \leq b$ and $c \leq f(I)_{\mathbf{2}}$ and $f(I)_{\mathbf{2}} \leq d$. Then $-1 \leq (h \cdot f)(O)_{\mathbf{1}}$ and $(h \cdot f)(O)_{\mathbf{1}} \leq 1$ and $-1 \leq (h \cdot f)(I)_{\mathbf{2}}$ and $(h \cdot f)(I)_{\mathbf{2}} \leq 1$.

(64) Let a, b, c, d be real numbers, h be a map from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$, f be a map from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$, and O, I be points of $\mathbb{I}$. Suppose $a < b$ and $c < d$ and $h = \mathrm{AffineMap}(\frac{2}{b-a}, -\frac{b+a}{b-a}, \frac{2}{d-c}, -\frac{d+c}{d-c})$ and $a \leq f(O)_{\mathbf{1}}$ and $f(O)_{\mathbf{1}} \leq b$ and $a < f(I)_{\mathbf{1}}$ and $f(I)_{\mathbf{1}} \leq b$. Then $-1 \leq (h \cdot f)(O)_{\mathbf{1}}$ and $(h \cdot f)(O)_{\mathbf{1}} \leq 1$ and $-1 < (h \cdot f)(I)_{\mathbf{1}}$ and $(h \cdot f)(I)_{\mathbf{1}} \leq 1$.

(65) Let a, b, c, d be real numbers, h be a map from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$, f be a map from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$, and O, I be points of $\mathbb{I}$. Suppose $a < b$ and $c < d$ and $h = \mathrm{AffineMap}(\frac{2}{b-a}, -\frac{b+a}{b-a}, \frac{2}{d-c}, -\frac{d+c}{d-c})$ and $d \geq f(O)_{\mathbf{2}}$ and $f(O)_{\mathbf{2}} > f(I)_{\mathbf{2}}$ and $f(I)_{\mathbf{2}} \geq c$. Then $1 \geq (h \cdot f)(O)_{\mathbf{2}}$ and $(h \cdot f)(O)_{\mathbf{2}} > (h \cdot f)(I)_{\mathbf{2}}$ and $(h \cdot f)(I)_{\mathbf{2}} \geq -1$.

(66) Let a, b, c, d be real numbers, h be a map from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$, f be a map from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$, and O, I be points of $\mathbb{I}$. Suppose $a < b$ and $c < d$ and $h = \mathrm{AffineMap}(\frac{2}{b-a}, -\frac{b+a}{b-a}, \frac{2}{d-c}, -\frac{d+c}{d-c})$ and $c \leq f(O)_{\mathbf{2}}$ and $f(O)_{\mathbf{2}} \leq d$ and $a < f(I)_{\mathbf{1}}$ and $f(I)_{\mathbf{1}} \leq b$. Then $-1 \leq (h \cdot f)(O)_{\mathbf{2}}$ and $(h \cdot f)(O)_{\mathbf{2}} \leq 1$ and $-1 < (h \cdot f)(I)_{\mathbf{1}}$ and $(h \cdot f)(I)_{\mathbf{1}} \leq 1$.

(67) Let a, b, c, d be real numbers, h be a map from $\mathcal{E}^2_{\mathrm{T}}$ into $\mathcal{E}^2_{\mathrm{T}}$, f be a map from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$, and O, I be points of $\mathbb{I}$. Suppose $a < b$ and $c < d$ and $h = \mathrm{AffineMap}(\frac{2}{b-a}, -\frac{b+a}{b-a}, \frac{2}{d-c}, -\frac{d+c}{d-c})$ and $a < f(I)_{\mathbf{1}}$ and $f(I)_{\mathbf{1}} < f(O)_{\mathbf{1}}$ and $f(O)_{\mathbf{1}} \leq b$. Then $-1 < (h \cdot f)(I)_{\mathbf{1}}$ and $(h \cdot f)(I)_{\mathbf{1}} < (h \cdot f)(O)_{\mathbf{1}}$ and $(h \cdot f)(O)_{\mathbf{1}} \leq 1$.

One can prove the following propositions:

(68) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{1}} = a$ and $(p_4)_{\mathbf{1}} = a$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} < (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \leq d$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is

continuous and one-to-one and $\operatorname{rng} f \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$ and $\operatorname{rng} g \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$. Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(69) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{1}} = a$ and $(p_4)_{\mathbf{1}} = a$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} < (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \leq d$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$ and $Q \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$. Then P meets Q.

(70) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{1}} = a$ and $(p_4)_{\mathbf{2}} = d$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $a \leq (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$ and $\operatorname{rng} g \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$. Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(71) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{1}} = a$ and $(p_4)_{\mathbf{2}} = d$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $a \leq (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$ and $Q \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$. Then P meets Q.

(72) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{1}} = a$ and $(p_4)_{\mathbf{1}} = b$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \leq d$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$ and $\operatorname{rng} g \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$. Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(73) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{1}} = a$ and $(p_4)_{\mathbf{1}} = b$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \leq d$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$ and $Q \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$. Then P meets Q.

(74) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{1}} = a$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$

and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(75) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{1}} = a$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(76) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{2}} = d$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a \leq (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(77) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{2}} = d$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a \leq (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(78) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{1}} = b$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a \leq (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \leq d$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(79) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{1}} = b$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a \leq (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \leq d$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(80) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a \leq (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq \operatorname{ClosedInsideOfRectangle}(a, b, c, d)$ and $\operatorname{rng} g \subseteq \operatorname{ClosedInsideOfRectangle}(a, b, c, d)$. Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(81) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a \leq (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq \operatorname{ClosedInsideOfRectangle}(a, b, c, d)$ and $Q \subseteq \operatorname{ClosedInsideOfRectangle}(a, b, c, d)$. Then P meets Q.

(82) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{1}} = b$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq \operatorname{ClosedInsideOfRectangle}(a, b, c, d)$ and $\operatorname{rng} g \subseteq \operatorname{ClosedInsideOfRectangle}(a, b, c, d)$. Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(83) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{1}} = b$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq \operatorname{ClosedInsideOfRectangle}(a, b, c, d)$ and $Q \subseteq \operatorname{ClosedInsideOfRectangle}(a, b, c, d)$. Then P meets Q.

(84) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $c \leq (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq \operatorname{ClosedInsideOfRectangle}(a, b, c, d)$ and $\operatorname{rng} g \subseteq \operatorname{ClosedInsideOfRectangle}(a, b, c, d)$. Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(85) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $c \leq (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and

$a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(86) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(87) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = a$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(88) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{2}} = d$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(89) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{2}} = d$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(90) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{1}} = b$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \leq d$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(91) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P,

Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{1}} = b$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \leq d$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$ and $Q \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$. Then P meets Q.

(92) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$ and $\operatorname{rng} g \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$. Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(93) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$ and $Q \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$. Then P meets Q.

(94) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{1}} = b$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$ and $\operatorname{rng} g \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$. Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(95) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{1}} = b$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$ and $Q \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$. Then P meets Q.

(96) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $c \leq (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and

$g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(97) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $c \leq (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(98) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(99) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a \leq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(100) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{1}} = b$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(101) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{1}} = b$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(102) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and

$(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $c \leq (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$ and $\operatorname{rng} g \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$. Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(103) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $c \leq (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} < (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$ and $Q \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$. Then P meets Q.

(104) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $c \leq (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$ and $\operatorname{rng} g \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$. Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(105) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $c \leq (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$ and $Q \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$. Then P meets Q.

(106) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = c$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$ and $\operatorname{rng} g \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$. Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(107) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = a$ and $(p_2)_{\mathbf{2}} = c$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$

and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(108) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{2}} = d$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(109) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{2}} = d$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(110) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{1}} = b$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \leq d$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(111) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{1}} = b$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \leq d$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(112) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(113) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = d$ and $(p_4)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and

$(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(114) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{1}} = b$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(115) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{1}} = b$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $c \leq (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} < (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(116) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $c \leq (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(117) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $c \leq (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(118) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(119) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = d$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$ and $Q \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$. Then P meets Q.

(120) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{1}} = b$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} \leq b$ and $d \geq (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} > (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} > (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \geq c$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\mathrm{rng}\, f \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$ and $\mathrm{rng}\, g \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$. Then $\mathrm{rng}\, f$ meets $\mathrm{rng}\, g$.

(121) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{1}} = b$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} \leq b$ and $d \geq (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} > (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} > (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \geq c$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$ and $Q \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$. Then P meets Q.

(122) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} \leq b$ and $d \geq (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} > (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \geq c$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\mathrm{rng}\, f \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$ and $\mathrm{rng}\, g \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$. Then $\mathrm{rng}\, f$ meets $\mathrm{rng}\, g$.

(123) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} \leq b$ and $d \geq (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} > (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \geq c$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$ and $Q \subseteq \mathrm{ClosedInsideOfRectangle}(a, b, c, d)$. Then P meets Q.

(124) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} \leq b$ and $c \leq (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continu-

ous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(125) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} \leq b$ and $c \leq (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \leq d$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(126) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = c$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} \leq b$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(127) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = d$ and $(p_2)_{\mathbf{2}} = c$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $a \leq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} \leq b$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} < (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} < (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(128) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = b$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{1}} = b$ and $d \geq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} > (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} > (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} > (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \geq c$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(129) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = b$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{1}} = b$ and $d \geq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} > (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} > (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} > (p_4)_{\mathbf{2}}$ and $(p_4)_{\mathbf{2}} \geq c$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(130) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = b$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{2}} = c$ and $d \geq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} > (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} > (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \geq c$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and

$g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(131) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = b$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{1}} = b$ and $(p_4)_{\mathbf{2}} = c$ and $d \geq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} > (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} > (p_3)_{\mathbf{2}}$ and $(p_3)_{\mathbf{2}} \geq c$ and $a < (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} \leq b$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(132) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = b$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $d \geq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} > (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \geq c$ and $b \geq (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} > (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} > a$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(133) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = b$ and $(p_2)_{\mathbf{1}} = b$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $d \geq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} > (p_2)_{\mathbf{2}}$ and $(p_2)_{\mathbf{2}} \geq c$ and $b \geq (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} > (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} > a$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(134) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = b$ and $(p_2)_{\mathbf{2}} = c$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $b \geq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} > (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} > (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} > a$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $\operatorname{rng} g \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(135) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{1}} = b$ and $(p_2)_{\mathbf{2}} = c$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $c \leq (p_1)_{\mathbf{2}}$ and $(p_1)_{\mathbf{2}} \leq d$ and $b \geq (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} > (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} > (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} > a$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq$ ClosedInsideOfRectangle(a, b, c, d) and $Q \subseteq$ ClosedInsideOfRectangle(a, b, c, d). Then P meets Q.

(136) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and f, g be maps from $\mathbb{I}$ into $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = c$ and $(p_2)_{\mathbf{2}} = c$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $b \geq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} > (p_2)_{\mathbf{1}}$ and

$(p_2)_{\mathbf{1}} > (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} > (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} > a$ and $f(0) = p_1$ and $f(1) = p_3$ and $g(0) = p_2$ and $g(1) = p_4$ and f is continuous and one-to-one and g is continuous and one-to-one and $\operatorname{rng} f \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$ and $\operatorname{rng} g \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$. Then $\operatorname{rng} f$ meets $\operatorname{rng} g$.

(137) Let p_1, p_2, p_3, p_4 be points of $\mathcal{E}^2_{\mathrm{T}}$, a, b, c, d be real numbers, and P, Q be subsets of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that $a < b$ and $c < d$ and $(p_1)_{\mathbf{2}} = c$ and $(p_2)_{\mathbf{2}} = c$ and $(p_3)_{\mathbf{2}} = c$ and $(p_4)_{\mathbf{2}} = c$ and $b \geq (p_1)_{\mathbf{1}}$ and $(p_1)_{\mathbf{1}} > (p_2)_{\mathbf{1}}$ and $(p_2)_{\mathbf{1}} > (p_3)_{\mathbf{1}}$ and $(p_3)_{\mathbf{1}} > (p_4)_{\mathbf{1}}$ and $(p_4)_{\mathbf{1}} > a$ and P is an arc from p_1 to p_3 and Q is an arc from p_2 to p_4 and $P \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$ and $Q \subseteq \text{ClosedInsideOfRectangle}(a, b, c, d)$. Then P meets Q.

References

[1] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(**1**):91–96, 1990.
[2] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(**1**):55–65, 1990.
[3] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(**1**):153–164, 1990.
[4] Czesław Byliński and Piotr Rudnicki. Bounding boxes for compact sets in $\mathcal{E}^2$. *Formalized Mathematics*, 6(**3**):427–440, 1997.
[5] Agata Darmochwał. Families of subsets, subspaces and mappings in topological spaces. *Formalized Mathematics*, 1(**2**):257–261, 1990.
[6] Agata Darmochwał. The Euclidean space. *Formalized Mathematics*, 2(**4**):599–603, 1991.
[7] Agata Darmochwał and Yatsuka Nakamura. Metric spaces as topological spaces – fundamental concepts. *Formalized Mathematics*, 2(**4**):605–608, 1991.
[8] Agata Darmochwał and Yatsuka Nakamura. The topological space $\mathcal{E}^2_{\mathrm{T}}$. Arcs, line segments and special polygonal arcs. *Formalized Mathematics*, 2(**5**):617–621, 1991.
[9] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(**1**):35–40, 1990.
[10] Artur Korniłowicz. The ordering of points on a curve. Part III. *Formalized Mathematics*, 10(**3**):169–171, 2002.
[11] Yatsuka Nakamura. On Outside Fashoda Meet Theorem. *Formalized Mathematics*, 9(**4**):697–704, 2001.
[12] Yatsuka Nakamura. General Fashoda meet theorem for unit circle and square. *Formalized Mathematics*, 11(**3**):213–224, 2003.
[13] Yatsuka Nakamura and Andrzej Trybulec. A decomposition of a simple closed curves and the order of their points. *Formalized Mathematics*, 6(**4**):563–572, 1997.
[14] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(**1**):223–230, 1990.
[15] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(**1**):9–11, 1990.
[16] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(**1**):67–71, 1990.
[17] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(**1**):73–83, 1990.

Received January 3, 2005