FORMALIZED MATHEMATICS
Volume 12, Number 1, 2004
University of Białystok

Little Bezout Theorem (Factor Theorem)[1]

Piotr Rudnicki
University of Alberta
Edmonton

Summary. We present a formalization of the factor theorem for univariate polynomials, also called the (little) Bezout theorem: Let r belong to a commutative ring L and $p(x)$ be a polynomial over L. Then $x-r$ divides $p(x)$ iff $p(r)=0$. We also prove some consequences of this theorem like that any non zero polynomial of degree n over an algebraically closed integral domain has n (non necessarily distinct) roots.

MML Identifier: UPROOTS.

The articles [28], [37], [26], [10], [2], [27], [36], [15], [20], [38], [7], [8], [3], [6], [35], [32], [24], [23], [11], [21], [16], [19], [17], [18], [1], [12], [33], [29], [22], [9], [34], [4], [25], [39], [13], [30], [14], [31], and [5] provide the notation and terminology for this paper.

1. Preliminaries

One can prove the following propositions:

(1) For every natural number n holds n is non empty iff $n=1$ or $n>1$.

(2) Let f be a finite sequence of elements of $\mathbb{N}$. Suppose that for every natural number i such that $i \in \operatorname{dom} f$ holds $f(i) \neq 0$. Then $\sum f = \operatorname{len} f$ if and only if $f = \operatorname{len} f \mapsto 1$.

The scheme *IndFinSeq0* deals with a finite sequence $\mathcal{A}$ and a binary predicate $\mathcal{P}$, and states that:

For every natural number i such that $1 \leqslant i$ and $i \leqslant \operatorname{len} \mathcal{A}$ holds $\mathcal{P}[i, \mathcal{A}(i)]$

[1] This work has been supported by NSERC Grant OGP9207.

© 2004 University of Białystok
ISSN 1426–2630

provided the parameters meet the following requirements:

- $\mathcal{P}[1, \mathcal{A}(1)]$, and
- For every natural number i such that $1 \leqslant i$ and $i < \operatorname{len} \mathcal{A}$ holds if $\mathcal{P}[i, \mathcal{A}(i)]$, then $\mathcal{P}[i+1, \mathcal{A}(i+1)]$.

We now state the proposition

(3) Let L be an add-associative right zeroed right complementable non empty loop structure and r be a finite sequence of elements of L. Suppose $\operatorname{len} r \geqslant 2$ and for every natural number k such that $2 < k$ and $k \in \operatorname{dom} r$ holds $r(k) = 0_L$. Then $\sum r = r_1 + r_2$.

2. Canonical Ordering of a Finite Set

Let A be a finite set. The functor $\operatorname{CFS}(A)$ yielding a finite sequence of elements of A is defined by the conditions (Def. 1).

(Def. 1)(i) $\operatorname{len} \operatorname{CFS}(A) = \operatorname{card} A$, and

(ii) there exists a finite sequence f such that $\operatorname{len} f = \operatorname{card} A$ and $f(1) = \langle \operatorname{choose}(A), A \setminus \{\operatorname{choose}(A)\} \rangle$ or $\operatorname{card} A = 0$ and for every natural number i such that $1 \leqslant i$ and $i < \operatorname{card} A$ and for every set x such that $f(i) = x$ holds $f(i+1) = \langle \operatorname{choose}(x_{\mathbf{2}}), x_{\mathbf{2}} \setminus \{\operatorname{choose}(x_{\mathbf{2}})\} \rangle$ and for every natural number i such that $i \in \operatorname{dom} \operatorname{CFS}(A)$ holds $(\operatorname{CFS}(A))(i) = f(i)_{\mathbf{1}}$.

The following four propositions are true:

(4) For every finite set A holds $\operatorname{CFS}(A)$ is one-to-one.

(5) For every finite set A holds $\operatorname{rng} \operatorname{CFS}(A) = A$.

(6) For every set a holds $\operatorname{CFS}(\{a\}) = \langle a \rangle$.

(7) For every finite set A holds $(\operatorname{CFS}(A))^{-1}$ is a function from A into $\operatorname{Seg} \operatorname{card} A$.

3. More about Bags

Let X be a set, let S be a finite subset of X, and let n be a natural number. The functor (S, n)-bag yields an element of $\operatorname{Bags} X$ and is defined by:

(Def. 2) (S, n)-bag $= \operatorname{EmptyBag} X + \cdot (S \longmapsto n)$.

We now state several propositions:

(8) Let X be a set, S be a finite subset of X, n be a natural number, and i be a set. If $i \notin S$, then $((S, n)\text{-bag})(i) = 0$.

(9) Let X be a set, S be a finite subset of X, n be a natural number, and i be a set. If $i \in S$, then $((S, n)\text{-bag})(i) = n$.

(10) For every set X and for every finite subset S of X and for every natural number n such that $n \neq 0$ holds $\operatorname{support}(S, n)\text{-bag} = S$.

(11) Let X be a set, S be a finite subset of X, and n be a natural number. If S is empty or $n = 0$, then (S, n)-bag $=$ EmptyBag X.

(12) Let X be a set, S, T be finite subsets of X, and n be a natural number. If S misses T, then $(S \cup T, n)$-bag $= (S, n)$-bag $+ (T, n)$-bag.

Let A be a set and let b be a bag of A. The functor degree(b) yielding a natural number is defined as follows:

(Def. 3) There exists a finite sequence f of elements of $\mathbb{N}$ such that degree(b) $= \sum f$ and $f = b \cdot \mathrm{CFS}(\mathrm{support}\, b)$.

We now state several propositions:

(13) For every set A and for every bag b of A holds $b =$ EmptyBag A iff degree(b) $= 0$.

(14) Let A be a set, S be a finite subset of A, and b be a bag of A. Then $S =$ support b and degree(b) $=$ card S if and only if $b = (S, 1)$-bag.

(15) Let A be a set, S be a finite subset of A, and b be a bag of A. Suppose support $b \subseteq S$. Then there exists a finite sequence f of elements of $\mathbb{N}$ such that $f = b \cdot \mathrm{CFS}(S)$ and degree(b) $= \sum f$.

(16) For every set A and for all bags b, b_1, b_2 of A such that $b = b_1 + b_2$ holds degree(b) $=$ degree(b_1) $+$ degree(b_2).

(17) Let L be an associative commutative unital non empty groupoid, f, g be finite sequences of elements of L, and p be a permutation of dom f. If $g = f \cdot p$, then $\prod g = \prod f$.

4. More on Polynomials

Let L be a non empty zero structure and let p be a polynomial of L. We say that p is non-zero if and only if:

(Def. 4) $p \neq \mathbf{0}.L$.

One can prove the following proposition

(18) For every non empty zero structure L and for every polynomial p of L holds p is non-zero iff len $p > 0$.

Let L be a non trivial non empty zero structure. Note that there exists a polynomial of L which is non-zero.

Let L be a non degenerated non empty multiplicative loop with zero structure and let x be an element of L. Note that $\langle x, \mathbf{1}_L \rangle$ is non-zero.

Next we state three propositions:

(19) For every non empty zero structure L and for every polynomial p of L such that len $p > 0$ holds $p(\mathrm{len}\, p -' 1) \neq 0_L$.

(20) Let L be a non empty zero structure and p be an algebraic sequence of L. If len $p = 1$, then $p = \langle p(0) \rangle$ and $p(0) \neq 0_L$.

(21) Let L be an add-associative right zeroed right complementable right distributive non empty double loop structure and p be a polynomial of L. Then $p * \mathbf{0}.L = \mathbf{0}.L$.

Let us mention that there exists a well unital non empty double loop structure which is algebraic-closed, add-associative, right zeroed, right complementable, Abelian, commutative, associative, distributive, integral domain-like, and non degenerated.

We now state the proposition

(22) Let L be an add-associative right zeroed right complementable distributive integral domain-like non empty double loop structure and p, q be polynomials of L. If $p * q = \mathbf{0}.L$, then $p = \mathbf{0}.L$ or $q = \mathbf{0}.L$.

Let L be an add-associative right zeroed right complementable distributive integral domain-like non empty double loop structure. Observe that Polynom-Ring L is integral domain-like.

Let L be an integral domain and let p, q be non-zero polynomials of L. One can check that $p * q$ is non-zero.

We now state a number of propositions:

(23) For every non degenerated commutative ring L and for all polynomials p, q of L holds $\operatorname{Roots} p \cup \operatorname{Roots} q \subseteq \operatorname{Roots}(p * q)$.

(24) For every integral domain L and for all polynomials p, q of L holds $\operatorname{Roots}(p * q) = \operatorname{Roots} p \cup \operatorname{Roots} q$.

(25) Let L be an add-associative right zeroed right complementable distributive non empty double loop structure, p be a polynomial of L, and p_1 be an element of Polynom-Ring L. If $p = p_1$, then $-p = -p_1$.

(26) Let L be an add-associative right zeroed right complementable distributive non empty double loop structure, p, q be polynomials of L, and p_1, q_1 be elements of Polynom-Ring L. If $p = p_1$ and $q = q_1$, then $p - q = p_1 - q_1$.

(27) Let L be an Abelian add-associative right zeroed right complementable distributive non empty double loop structure and p, q, r be polynomials of L. Then $p * q - p * r = p * (q - r)$.

(28) Let L be an add-associative right zeroed right complementable distributive non empty double loop structure and p, q be polynomials of L. If $p - q = \mathbf{0}.L$, then $p = q$.

(29) Let L be an Abelian add-associative right zeroed right complementable distributive integral domain-like non empty double loop structure and p, q, r be polynomials of L. If $p \neq \mathbf{0}.L$ and $p * q = p * r$, then $q = r$.

(30) Let L be an integral domain, n be a natural number, and p be a polynomial of L. If $p \neq \mathbf{0}.L$, then $p^n \neq \mathbf{0}.L$.

(31) For every commutative ring L and for all natural numbers i, j and for every polynomial p of L holds $p^i * p^j = p^{i+j}$.

(32) For every non empty multiplicative loop with zero structure L holds $\mathbf{1}.L = \langle \mathbf{1}_L \rangle$.

(33) Let L be an add-associative right zeroed right complementable right unital right distributive non empty double loop structure and p be a polynomial of L. Then $p * \langle \mathbf{1}_L \rangle = p$.

(34) Let L be an add-associative right zeroed right complementable distributive non empty double loop structure and p, q be polynomials of L. If $\operatorname{len} p = 0$ or $\operatorname{len} q = 0$, then $\operatorname{len}(p * q) = 0$.

(35) Let L be an add-associative right zeroed right complementable distributive non empty double loop structure and p, q be polynomials of L. If $p * q$ is non-zero, then p is non-zero and q is non-zero.

(36) Let L be an add-associative right zeroed right complementable distributive commutative associative left unital non empty double loop structure and p, q be polynomials of L. If $p(\operatorname{len} p -' 1) \cdot q(\operatorname{len} q -' 1) \neq 0_L$, then $0 < \operatorname{len}(p * q)$.

(37) Let L be an add-associative right zeroed right complementable distributive commutative associative left unital integral domain-like non empty double loop structure and p, q be polynomials of L. If $1 < \operatorname{len} p$ and $1 < \operatorname{len} q$, then $\operatorname{len} p < \operatorname{len}(p * q)$.

(38) Let L be an add-associative right zeroed right complementable left distributive non empty double loop structure, a, b be elements of L, and p be a polynomial of L. Then $(\langle a, b \rangle * p)(0) = a \cdot p(0)$ and for every natural number i holds $(\langle a, b \rangle * p)(i + 1) = a \cdot p(i + 1) + b \cdot p(i)$.

(39) Let L be an add-associative right zeroed right complementable distributive well unital commutative associative non degenerated non empty double loop structure, r be an element of L, and q be a non-zero polynomial of L. Then $\operatorname{len}(\langle r, \mathbf{1}_L \rangle * q) = \operatorname{len} q + 1$.

(40) Let L be a non degenerated commutative ring, x be an element of L, and i be a natural number. Then $\operatorname{len}(\langle x, \mathbf{1}_L \rangle^i) = i + 1$.

Let L be a non degenerated commutative ring, let x be an element of L, and let n be a natural number. Note that $\langle x, \mathbf{1}_L \rangle^n$ is non-zero.

Next we state two propositions:

(41) Let L be a non degenerated commutative ring, x be an element of L, q be a non-zero polynomial of L, and i be a natural number. Then $\operatorname{len}(\langle x, \mathbf{1}_L \rangle^i * q) = i + \operatorname{len} q$.

(42) Let L be an add-associative right zeroed right complementable distributive well unital commutative associative non degenerated non empty double loop structure, r be an element of L, and p, q be polynomials of L. If $p = \langle r, \mathbf{1}_L \rangle * q$ and $p(\operatorname{len} p -' 1) = \mathbf{1}_L$, then $q(\operatorname{len} q -' 1) = \mathbf{1}_L$.

5. Little Bezout Theorem

Let L be a non empty zero structure, let p be a polynomial of L, and let n be a natural number. The functor poly_shift(p, n) yields a polynomial of L and is defined by:

(Def. 5) For every natural number i holds $(\text{poly_shift}(p, n))(i) = p(n + i)$.

We now state several propositions:

(43) For every non empty zero structure L and for every polynomial p of L holds $\text{poly_shift}(p, 0) = p$.

(44) Let L be a non empty zero structure, n be a natural number, and p be a polynomial of L. If $n \geqslant \operatorname{len} p$, then $\text{poly_shift}(p, n) = \mathbf{0}.L$.

(45) Let L be a non degenerated non empty multiplicative loop with zero structure, n be a natural number, and p be a polynomial of L. If $n \leqslant \operatorname{len} p$, then $\operatorname{len} \text{poly_shift}(p, n) + n = \operatorname{len} p$.

(46) Let L be a non degenerated commutative ring, x be an element of L, n be a natural number, and p be a polynomial of L. If $n < \operatorname{len} p$, then $\text{eval}(\text{poly_shift}(p, n), x) = x \cdot \text{eval}(\text{poly_shift}(p, n + 1), x) + p(n)$.

(47) For every non degenerated commutative ring L and for every polynomial p of L such that $\operatorname{len} p = 1$ holds $\operatorname{Roots} p = \emptyset$.

Let L be a non degenerated commutative ring, let r be an element of L, and let p be a polynomial of L. Let us assume that r is a root of p. The functor poly_quotient(p, r) yielding a polynomial of L is defined as follows:

(Def. 6)(i) $\operatorname{len} \text{poly_quotient}(p, r) + 1 = \operatorname{len} p$ and for every natural number i holds $(\text{poly_quotient}(p, r))(i) = \text{eval}(\text{poly_shift}(p, i + 1), r)$ if $\operatorname{len} p > 0$,

(ii) $\text{poly_quotient}(p, r) = \mathbf{0}.L$, otherwise.

Next we state several propositions:

(48) Let L be a non degenerated commutative ring, r be an element of L, and p be a non-zero polynomial of L. If r is a root of p, then $\operatorname{len} \text{poly_quotient}(p, r) > 0$.

(49) Let L be an add-associative right zeroed right complementable left distributive well unital non empty double loop structure and x be an element of L. Then $\operatorname{Roots}\langle -x, \mathbf{1}_L \rangle = \{x\}$.

(50) Let L be a non trivial commutative ring, x be an element of L, and p, q be polynomials of L. If $p = \langle -x, \mathbf{1}_L \rangle * q$, then x is a root of p.

(51) Let L be a non degenerated commutative ring, r be an element of L, and p be a polynomial of L. If r is a root of p, then $p = \langle -r, \mathbf{1}_L \rangle * \text{poly_quotient}(p, r)$.

(52) Let L be a non degenerated commutative ring, r be an element of L, and p, q be polynomials of L. If $p = \langle -r, \mathbf{1}_L \rangle * q$, then r is a root of p.

6. Polynomials Defined by Roots

Let L be an integral domain and let p be a non-zero polynomial of L. One can verify that $\mathrm{Roots}\, p$ is finite.

Let L be a non degenerated commutative ring, let x be an element of L, and let p be a non-zero polynomial of L. The functor multiplicity(p, x) yields a natural number and is defined by the condition (Def. 7).

(Def. 7) There exists a finite non empty subset F of $\mathbb{N}$ such that $F = \{k; k$ ranges over natural numbers: $\bigvee_{q\,:\,\text{polynomial of } L}\ p = \langle -x, \mathbf{1}_L\rangle^k * q\}$ and multiplicity$(p, x) = \max F$.

Next we state two propositions:

(53) Let L be a non degenerated commutative ring, p be a non-zero polynomial of L, and x be an element of L. Then x is a root of p if and only if multiplicity$(p, x) \geqslant 1$.

(54) For every non degenerated commutative ring L and for every element x of L holds multiplicity$(\langle -x, \mathbf{1}_L\rangle, x) = 1$.

Let L be an integral domain and let p be a non-zero polynomial of L. The functor BRoots(p) yields a bag of the carrier of L and is defined as follows:

(Def. 8) $\mathrm{support}\, \mathrm{BRoots}(p) = \mathrm{Roots}\, p$ and for every element x of L holds $(\mathrm{BRoots}(p))(x) = \mathrm{multiplicity}(p, x)$.

Next we state several propositions:

(55) For every integral domain L and for every element x of L holds $\mathrm{BRoots}(\langle -x, \mathbf{1}_L\rangle) = (\{x\}, 1)\text{-bag}$.

(56) Let L be an integral domain, x be an element of L, and p, q be non-zero polynomials of L. Then multiplicity$(p * q, x) =$ multiplicity$(p, x) +$ multiplicity(q, x).

(57) For every integral domain L and for all non-zero polynomials p, q of L holds $\mathrm{BRoots}(p * q) = \mathrm{BRoots}(p) + \mathrm{BRoots}(q)$.

(58) For every integral domain L and for every non-zero polynomial p of L such that $\mathrm{len}\, p = 1$ holds degree$(\mathrm{BRoots}(p)) = 0$.

(59) For every integral domain L and for every element x of L and for every natural number n holds $\mathrm{degree}(\mathrm{BRoots}(\langle -x, \mathbf{1}_L\rangle^n)) = n$.

(60) For every algebraic-closed integral domain L and for every non-zero polynomial p of L holds $\mathrm{degree}(\mathrm{BRoots}(p)) = \mathrm{len}\, p -' 1$.

Let L be an add-associative right zeroed right complementable distributive non empty double loop structure, let c be an element of L, and let n be a natural number. The functor fpoly_mult_root(c, n) yielding a finite sequence of elements of Polynom-Ring L is defined as follows:

(Def. 9) $\mathrm{len}\, \mathrm{fpoly_mult_root}(c, n) = n$ and for every natural number i such that $i \in \mathrm{dom}\, \mathrm{fpoly_mult_root}(c, n)$ holds $(\mathrm{fpoly_mult_root}(c, n))(i) = \langle -c, \mathbf{1}_L\rangle$.

Let L be an add-associative right zeroed right complementable distributive non empty double loop structure and let b be a bag of the carrier of L. The functor poly_with_roots(b) yields a polynomial of L and is defined by the condition (Def. 10).

(Def. 10) There exists a finite sequence f of elements of (the carrier of Polynom-Ring L)* and there exists a finite sequence s of elements of L such that $\operatorname{len} f = \operatorname{card} \operatorname{support} b$ and $s = \operatorname{CFS}(\operatorname{support} b)$ and for every natural number i such that $i \in \operatorname{dom} f$ holds $f(i) = \operatorname{fpoly_mult_root}(s_i, b(s_i))$ and $\operatorname{poly_with_roots}(b) = \prod \operatorname{Flat}(f)$.

The following propositions are true:

(61) Let L be an Abelian add-associative right zeroed right complementable commutative distributive right unital non empty double loop structure. Then poly_with_roots(EmptyBag (the carrier of L)) $= \langle \mathbf{1}_L \rangle$.

(62) Let L be an add-associative right zeroed right complementable distributive non empty double loop structure and c be an element of L. Then poly_with_roots($(\{c\}, 1)$ -bag) $= \langle -c, \mathbf{1}_L \rangle$.

(63) Let L be an add-associative right zeroed right complementable distributive non empty double loop structure, b be a bag of the carrier of L, f be a finite sequence of elements of (the carrier of Polynom-Ring L)*, and s be a finite sequence of elements of L. Suppose $\operatorname{len} f = \operatorname{card} \operatorname{support} b$ and $s = \operatorname{CFS}(\operatorname{support} b)$ and for every natural number i such that $i \in \operatorname{dom} f$ holds $f(i) = \operatorname{fpoly_mult_root}(s_i, b(s_i))$. Then $\operatorname{len} \operatorname{Flat}(f) = \operatorname{degree}(b)$.

(64) Let L be an add-associative right zeroed right complementable distributive non empty double loop structure, b be a bag of the carrier of L, f be a finite sequence of elements of (the carrier of Polynom-Ring L)*, s be a finite sequence of elements of L, and c be an element of L such that $\operatorname{len} f = \operatorname{card} \operatorname{support} b$ and $s = \operatorname{CFS}(\operatorname{support} b)$ and for every natural number i such that $i \in \operatorname{dom} f$ holds $f(i) = \operatorname{fpoly_mult_root}(s_i, b(s_i))$. Then

(i) if $c \in \operatorname{support} b$, then $\operatorname{card}(\operatorname{Flat}(f)^{-1}(\{\langle -c, \mathbf{1}_L \rangle\})) = b(c)$, and

(ii) if $c \notin \operatorname{support} b$, then $\operatorname{card}(\operatorname{Flat}(f)^{-1}(\{\langle -c, \mathbf{1}_L \rangle\})) = 0$.

(65) For every commutative ring L and for all bags b_1, b_2 of the carrier of L holds $\operatorname{poly_with_roots}(b_1 + b_2) = \operatorname{poly_with_roots}(b_1) * \operatorname{poly_with_roots}(b_2)$.

(66) Let L be an algebraic-closed integral domain and p be a non-zero polynomial of L. If $p(\operatorname{len} p -' 1) = \mathbf{1}_L$, then $p = \operatorname{poly_with_roots}(\operatorname{BRoots}(p))$.

(67) Let L be a commutative ring, s be a non empty finite subset of L, and f be a finite sequence of elements of Polynom-Ring L. Suppose $\operatorname{len} f = \operatorname{card} s$ and for every natural number i and for every element c of L such that $i \in \operatorname{dom} f$ and $c = (\operatorname{CFS}(s))(i)$ holds $f(i) = \langle -c, \mathbf{1}_L \rangle$. Then poly_with_roots($(s, 1)$ -bag) $= \prod f$.

(68) Let L be a non trivial commutative ring, s be a non empty finite subset

of L, x be an element of L, and f be a finite sequence of elements of L. Suppose $\operatorname{len} f = \operatorname{card} s$ and for every natural number i and for every element c of L such that $i \in \operatorname{dom} f$ and $c = (\mathrm{CFS}(s))(i)$ holds $f(i) = \operatorname{eval}(\langle -c, \mathbf{1}_L \rangle, x)$. Then $\operatorname{eval}(\operatorname{poly_with_roots}((s,1)\text{-bag}), x) = \prod f$.

References

[1] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(**2**):377–382, 1990.
[2] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(**1**):41–46, 1990.
[3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(**1**):107–114, 1990.
[4] Grzegorz Bancerek and Piotr Rudnicki. On defining functions on trees. *Formalized Mathematics*, 4(**1**):91–101, 1993.
[5] Józef Białas. Group and field definitions. *Formalized Mathematics*, 1(**3**):433–439, 1990.
[6] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(**3**):529–536, 1990.
[7] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(**1**):55–65, 1990.
[8] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(**1**):153–164, 1990.
[9] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Formalized Mathematics*, 1(**3**):521–527, 1990.
[10] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(**1**):47–53, 1990.
[11] Czesław Byliński. The sum and product of finite sequences of real numbers. *Formalized Mathematics*, 1(**4**):661–668, 1990.
[12] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(**1**):165–167, 1990.
[13] Andrzej Kondracki. The Chinese Remainder Theorem. *Formalized Mathematics*, 6(**4**):573–577, 1997.
[14] Jarosław Kotowicz. Monotone real sequences. Subsequences. *Formalized Mathematics*, 1(**3**):471–475, 1990.
[15] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Formalized Mathematics*, 1(**2**):335–342, 1990.
[16] Anna Justyna Milewska. The field of complex numbers. *Formalized Mathematics*, 9(**2**):265–269, 2001.
[17] Robert Milewski. The evaluation of polynomials. *Formalized Mathematics*, 9(**2**):391–395, 2001.
[18] Robert Milewski. Fundamental theorem of algebra. *Formalized Mathematics*, 9(**3**):461–470, 2001.
[19] Robert Milewski. The ring of polynomials. *Formalized Mathematics*, 9(**2**):339–346, 2001.
[20] Michał Muzalewski. Construction of rings and left-, right-, and bi-modules over a ring. *Formalized Mathematics*, 2(**1**):3–11, 1991.
[21] Michał Muzalewski and Lesław W. Szczerba. Construction of finite sequences over ring and left-, right-, and bi-modules over a ring. *Formalized Mathematics*, 2(**1**):97–104, 1991.
[22] Yatsuka Nakamura, Piotr Rudnicki, Andrzej Trybulec, and Pauline N. Kawamoto. Preliminaries to circuits, I. *Formalized Mathematics*, 5(**2**):167–172, 1996.
[23] Takaya Nishiyama and Yasuho Mizuhara. Binary arithmetics. *Formalized Mathematics*, 4(**1**):83–86, 1993.
[24] Jan Popiołek. Real normed space. *Formalized Mathematics*, 2(**1**):111–115, 1991.
[25] Piotr Rudnicki and Andrzej Trybulec. Multivariate polynomials with arbitrary number of variables. *Formalized Mathematics*, 9(**1**):95–110, 2001.
[26] Andrzej Trybulec. Subsets of complex numbers. *To appear in Formalized Mathematics*.
[27] Andrzej Trybulec. Semilattice operations on finite subsets. *Formalized Mathematics*, 1(**2**):369–376, 1990.
[28] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(**1**):9–11, 1990.

[29] Andrzej Trybulec. Tuples, projections and Cartesian products. *Formalized Mathematics*, 1(**1**):97–105, 1990.
[30] Andrzej Trybulec. Many-sorted sets. *Formalized Mathematics*, 4(**1**):15–22, 1993.
[31] Andrzej Trybulec. On the sets inhabited by numbers. *Formalized Mathematics*, 11(**4**):341–347, 2003.
[32] Wojciech A. Trybulec. Binary operations on finite sequences. *Formalized Mathematics*, 1(**5**):979–981, 1990.
[33] Wojciech A. Trybulec. Groups. *Formalized Mathematics*, 1(**5**):821–827, 1990.
[34] Wojciech A. Trybulec. Linear combinations in real linear space. *Formalized Mathematics*, 1(**3**):581–588, 1990.
[35] Wojciech A. Trybulec. Pigeon hole principle. *Formalized Mathematics*, 1(**3**):575–579, 1990.
[36] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(**2**):291–296, 1990.
[37] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(**1**):67–71, 1990.
[38] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(**1**):73–83, 1990.
[39] Katarzyna Zawadzka. The sum and product of finite sequences of elements of a field. *Formalized Mathematics*, 3(**2**):205–211, 1992.

Received December 30, 2003