FORMALIZED MATHEMATICS
Volume 13, Number 2, Pages 347–352
University of Białystok, 2005

Limit of Sequence of Subsets

Bo Zhang
Shinshu University
Nagano, Japan

Hiroshi Yamazaki
Shinshu University
Nagano, Japan

Yatsuka Nakamura
Shinshu University
Nagano, Japan

Summary. A concept of "limit of sequence of subsets" is defined here. This article contains the following items: 1. definition of the superior sequence and the inferior sequence of sets, 2. definition of the superior limit and the inferior limit of sets, and additional properties for the sigma-field of sets, 3. definition of the limit value of a convergent sequence of sets, and additional properties for the sigma-field of sets.

MML identifier: `SETLIM_1`, version: `7.5.01 4.39.921`

The notation and terminology used here are introduced in the following papers: [9], [1], [13], [2], [10], [6], [11], [4], [12], [14], [8], [7], [3], and [5].

For simplicity, we adopt the following rules: n, m, k, k_1, k_2 denote natural numbers, x, X, Y, Z denote sets, A denotes a subset of X, B, A_1, A_2, A_3 denote sequences of subsets of X, S_1 denotes a σ-field of subsets of X, and S, S_2, S_3, S_4 denote sequences of subsets of S_1.

Next we state a number of propositions:

(1) For every function f from $\mathbb{N}$ into Y and for every n holds $\{f(k) : n \leq k\} \neq \emptyset$.

(2) For every function f from $\mathbb{N}$ into Y holds $f(n+m) \in \{f(k) : n \leq k\}$.

(3) For every function f from $\mathbb{N}$ into Y holds $\{f(k_1) : n \leq k_1\} = \{f(k_2) : n+1 \leq k_2\} \cup \{f(n)\}$.

(4) Let f be a function from $\mathbb{N}$ into Y. Then for every k_1 holds $x \in f(n+k_1)$ if and only if for every Z such that $Z \in \{f(k_2) : n \leq k_2\}$ holds $x \in Z$.

(5) For every non empty set Y and for every function f from $\mathbb{N}$ into Y holds $x \in \operatorname{rng} f$ iff there exists n such that $x = f(n)$.

(6) For every non empty set Y and for every function f from $\mathbb{N}$ into Y holds $\operatorname{rng} f = \{f(k)\}$.

© 2005 University of Białystok
ISSN 1426–2630

(7) For every non empty set Y and for every function f from $\mathbb{N}$ into Y holds $\operatorname{rng}(f \upharpoonright k) = \{f(n) : k \leq n\}$.

(8) $x \in \bigcap \operatorname{rng} B$ iff for every n holds $x \in B(n)$.

(9) Intersection $B = \bigcap \operatorname{rng} B$.

(10) Intersection $B \subseteq \bigcup B$.

(11) If for every n holds $B(n) = A$, then $\bigcup B = A$.

(12) If for every n holds $B(n) = A$, then Intersection $B = A$.

(13) If B is constant, then $\bigcup B =$ Intersection B.

(14) If B is constant and the value of $B = A$, then for every n holds $\bigcup\{B(k) : n \leq k\} = A$.

(15) If B is constant and the value of $B = A$, then for every n holds $\bigcap\{B(k) : n \leq k\} = A$.

(16) Let given X, B and f be a function. Suppose $\operatorname{dom} f = \mathbb{N}$ and for every n holds $f(n) = \bigcap\{B(k) : n \leq k\}$. Then f is a sequence of subsets of X.

(17) Let X be a set, B be a sequence of subsets of X, and f be a function. Suppose $\operatorname{dom} f = \mathbb{N}$ and for every n holds $f(n) = \bigcup\{B(k) : n \leq k\}$. Then f is a function from $\mathbb{N}$ into 2^X.

Let us consider X, B. We say that B is monotone if and only if:

(Def. 1) B is non-decreasing or non-increasing.

Let B be a function. The inferior setsequence B yields a function and is defined by the conditions (Def. 2).

(Def. 2)(i) dom (the inferior setsequence B) $= \mathbb{N}$, and

(ii) for every n holds (the inferior setsequence B)$(n) = \bigcap\{B(k) : n \leq k\}$.

Let X be a set and let B be a sequence of subsets of X. Then the inferior setsequence B is a sequence of subsets of X.

Let B be a function. The superior setsequence B yields a function and is defined by the conditions (Def. 3).

(Def. 3)(i) dom (the superior setsequence B) $= \mathbb{N}$, and

(ii) for every n holds (the superior setsequence B)$(n) = \bigcup\{B(k) : n \leq k\}$.

Let X be a set and let B be a sequence of subsets of X. Then the superior setsequence B is a sequence of subsets of X.

Next we state several propositions:

(18) (The inferior setsequence B)$(0) =$ Intersection B.

(19) (The superior setsequence B)$(0) = \bigcup B$.

(20) $x \in$ (the inferior setsequence B)(n) iff for every k holds $x \in B(n + k)$.

(21) $x \in$ (the superior setsequence B)(n) iff there exists k such that $x \in B(n + k)$.

(22) (The inferior setsequence B)$(n) =$ (the inferior setsequence B)$(n + 1) \cap B(n)$.

(23) (The superior setsequence B)(n) = (the superior setsequence B)$(n+1) \cup B(n)$.

(24) The inferior setsequence B is non-decreasing.

(25) The superior setsequence B is non-increasing.

(26) The inferior setsequence B is monotone and the superior setsequence B is monotone.

Let X be a set and let A be a sequence of subsets of X. Observe that the inferior setsequence A is non-decreasing.

Let X be a set and let A be a sequence of subsets of X. Observe that the superior setsequence A is non-increasing.

The following propositions are true:

(27) Intersection $B \subseteq$ (the inferior setsequence B)(n).

(28) (The superior setsequence B)$(n) \subseteq \bigcup B$.

(29) For all B, n holds $\{B(k) : n \leq k\}$ is a family of subsets of X.

(30) $\bigcup B =$ (Intersection Complement B)$^{\mathrm{c}}$.

(31) (The inferior setsequence B)(n) = (the superior setsequence Complement B)$(n)^{\mathrm{c}}$.

(32) (The superior setsequence B)(n) = (the inferior setsequence Complement B)$(n)^{\mathrm{c}}$.

(33) Complement (the inferior setsequence B) = the superior setsequence Complement B.

(34) Complement (the superior setsequence B) = the inferior setsequence Complement B.

(35) Suppose that for every n holds $A_3(n) = A_1(n) \cup A_2(n)$. Let given n. Then (the inferior setsequence B)$(n) \cup$ (the inferior setsequence A_2)$(n) \subseteq$ (the inferior setsequence A_3)(n).

(36) Suppose that for every n holds $A_3(n) = A_1(n) \cap A_2(n)$. Let given n. Then (the inferior setsequence A_3)$(n) =$ (the inferior setsequence A_1)$(n) \cap$ (the inferior setsequence A_2)(n).

(37) Suppose that for every n holds $A_3(n) = A_1(n) \cup A_2(n)$. Let given n. Then (the superior setsequence A_3)$(n) =$ (the superior setsequence A_1)$(n) \cup$ (the superior setsequence A_2)(n).

(38) Suppose that for every n holds $A_3(n) = A_1(n) \cap A_2(n)$. Let given n. Then (the superior setsequence A_3)$(n) \subseteq$ (the superior setsequence A_1)$(n) \cap$ (the superior setsequence A_2)(n).

(39) If B is constant and the value of $B = A$, then for every n holds (the inferior setsequence B)$(n) = A$.

(40) If B is constant and the value of $B = A$, then for every n holds (the superior setsequence B)$(n) = A$.

(41) If B is non-decreasing, then $B(n) \subseteq$ (the superior setsequence B)$(n+1)$.

(42) If B is non-decreasing, then (the superior setsequence B)$(n) =$ (the superior setsequence B)$(n+1)$.

(43) If B is non-decreasing, then (the superior setsequence B)$(n) = \bigcup B$.

(44) If B is non-decreasing, then Intersection (the superior setsequence B) $= \bigcup B$.

(45) If B is non-decreasing, then $B(n) \subseteq$ (the inferior setsequence B)$(n+1)$.

(46) If B is non-decreasing, then (the inferior setsequence B)$(n) = B(n)$.

(47) If B is non-decreasing, then the inferior setsequence $B = B$.

(48) If B is non-increasing, then (the superior setsequence B)$(n+1) \subseteq B(n)$.

(49) If B is non-increasing, then (the superior setsequence B)$(n) = B(n)$.

(50) If B is non-increasing, then the superior setsequence $B = B$.

(51) If B is non-increasing, then (the inferior setsequence B)$(n+1) \subseteq B(n)$.

(52) If B is non-increasing, then (the inferior setsequence B)$(n) =$ (the inferior setsequence B)$(n+1)$.

(53) If B is non-increasing, then (the inferior setsequence B)$(n) =$ Intersection B.

(54) If B is non-increasing, then $\bigcup$(the inferior setsequence B) $=$ Intersection B.

Let X be a set and let B be a sequence of subsets of X. Then $\liminf B$ can be characterized by the condition:

(Def. 4) $\liminf B = \bigcup$(the inferior setsequence B).

Let X be a set and let B be a sequence of subsets of X. Then $\limsup B$ can be characterized by the condition:

(Def. 5) $\limsup B =$ Intersection (the superior setsequence B).

Let X be a set and let B be a sequence of subsets of X. We introduce $\lim B$ as a synonym of $\limsup B$.

Next we state a number of propositions:

(55) Intersection $B \subseteq \liminf B$.

(56) $\liminf B = \lim$ (the inferior setsequence B).

(57) $\limsup B = \lim$ (the superior setsequence B).

(58) $\limsup B = (\liminf$ Complement $B)^{\mathrm{c}}$.

(59) If B is constant and the value of $B = A$, then B is convergent and $\lim B = A$ and $\liminf B = A$ and $\limsup B = A$.

(60) If B is non-decreasing, then $\limsup B = \bigcup B$.

(61) If B is non-decreasing, then $\liminf B = \bigcup B$.

(62) If B is non-increasing, then $\limsup B =$ Intersection B.

(63) If B is non-increasing, then $\liminf B =$ Intersection B.

(64) If B is non-decreasing, then B is convergent and $\lim B = \bigcup B$.

(65) If B is non-increasing, then B is convergent and $\lim B = \operatorname{Intersection} B$.

(66) If B is monotone, then B is convergent.

Let X be a set, let S_1 be a σ-field of subsets of X, and let S be a sequence of subsets of S_1. Let us observe that S is constant if and only if:

(Def. 6) There exists an element A of S_1 such that for every n holds $S(n) = A$.

Let X be a set, let S_1 be a σ-field of subsets of X, and let S be a sequence of subsets of S_1. Then the inferior setsequence S is a sequence of subsets of S_1.

Let X be a set, let S_1 be a σ-field of subsets of X, and let S be a sequence of subsets of S_1. Then the superior setsequence S is a sequence of subsets of S_1.

The following propositions are true:

(67) $x \in \limsup S$ iff for every n there exists k such that $x \in S(n+k)$.

(68) $x \in \liminf S$ iff there exists n such that for every k holds $x \in S(n+k)$.

(69) $\operatorname{Intersection} S \subseteq \liminf S$.

(70) $\limsup S \subseteq \bigcup S$.

(71) $\liminf S \subseteq \limsup S$.

Let X be a set, let S_1 be a σ-field of subsets of X, and let S be a sequence of subsets of S_1. The functor $S^{\mathbf{c}}$ yields a sequence of subsets of S_1 and is defined by:

(Def. 7) $S^{\mathbf{c}} = \operatorname{Complement} S$.

Next we state a number of propositions:

(72) $\liminf S = (\limsup(S^{\mathbf{c}}))^{\mathrm{c}}$.

(73) $\limsup S = (\liminf(S^{\mathbf{c}}))^{\mathrm{c}}$.

(74) If for every n holds $S_4(n) = S_2(n) \cup S_3(n)$, then $\liminf S_2 \cup \liminf S_3 \subseteq \liminf S_4$.

(75) If for every n holds $S_4(n) = S_2(n) \cap S_3(n)$, then $\liminf S_4 = \liminf S_2 \cap \liminf S_3$.

(76) If for every n holds $S_4(n) = S_2(n) \cup S_3(n)$, then $\limsup S_4 = \limsup S_2 \cup \limsup S_3$.

(77) If for every n holds $S_4(n) = S_2(n) \cap S_3(n)$, then $\limsup S_4 \subseteq \limsup S_2 \cap \limsup S_3$.

(78) If S is constant and the value of $S = A$, then S is convergent and $\lim S = A$ and $\liminf S = A$ and $\limsup S = A$.

(79) If S is non-decreasing, then $\limsup S = \bigcup S$.

(80) If S is non-decreasing, then $\liminf S = \bigcup S$.

(81) If S is non-decreasing, then S is convergent and $\lim S = \bigcup S$.

(82) If S is non-increasing, then $\limsup S = \operatorname{Intersection} S$.

(83) If S is non-increasing, then $\liminf S = \operatorname{Intersection} S$.

(84) If S is non-increasing, then S is convergent and $\lim S = \text{Intersection } S$.

(85) If S is monotone, then S is convergent.

References

[1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(**1**):41–46, 1990.
[2] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(**1**):55–65, 1990.
[3] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(**1**):153–164, 1990.
[4] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(**1**):47–53, 1990.
[5] Adam Grabowski. On the Kuratowski limit operators. *Formalized Mathematics*, 11(**4**):399–409, 2003.
[6] Jarosław Kotowicz. Monotone real sequences. Subsequences. *Formalized Mathematics*, 1(**3**):471–475, 1990.
[7] Andrzej Nędzusiak. σ-fields and probability. *Formalized Mathematics*, 1(**2**):401–407, 1990.
[8] Beata Padlewska. Families of sets. *Formalized Mathematics*, 1(**1**):147–152, 1990.
[9] Andrzej Trybulec. Subsets of complex numbers. *To appear in Formalized Mathematics.*
[10] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(**1**):9–11, 1990.
[11] Andrzej Trybulec. Moore-Smith convergence. *Formalized Mathematics*, 6(**2**):213–225, 1997.
[12] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(**1**):67–71, 1990.
[13] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(**1**):73–83, 1990.
[14] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(**1**):181–186, 1990.

Received March 15, 2005