FORMALIZED MATHEMATICS
Volume 12, Number 2, 2004
University of Białystok

The Class of Series-Parallel Graphs. Part III

Krzysztof Retel
University of Bialystok

Summary. This paper contains some facts and theorems relating to the following operations on graphs: union, sum, complement and "embeds". We also introduce connected graphs to prove that a finite irreflexive symmetric N-free graph is a finite series-parallel graph. This article continues the formalization of [22].

MML Identifier: NECKLA_3.

The papers [25], [24], [28], [12], [29], [31], [30], [2], [13], [1], [27], [18], [17], [8], [14], [16], [20], [23], [7], [10], [26], [11], [4], [6], [19], [15], [5], [21], [3], and [9] provide the notation and terminology for this paper.

1. Preliminaries

In this paper A, B, a, b, c, d, e, f, g, h denote sets.

One can prove the following three propositions:

(1) $\mathrm{id}_A \restriction B = \mathrm{id}_A \cap [\!: B, B :]$.

(2) $\mathrm{id}_{\{a,b,c,d\}} = \{\langle a, a\rangle, \langle b, b\rangle, \langle c, c\rangle, \langle d, d\rangle\}$.

(3) $[\!: \{a, b, c, d\}, \{e, f, g, h\} :] = \{\langle a, e\rangle, \langle a, f\rangle, \langle b, e\rangle, \langle b, f\rangle, \langle a, g\rangle, \langle a, h\rangle, \langle b, g\rangle, \langle b, h\rangle\} \cup \{\langle c, e\rangle, \langle c, f\rangle, \langle d, e\rangle, \langle d, f\rangle, \langle c, g\rangle, \langle c, h\rangle, \langle d, g\rangle, \langle d, h\rangle\}$.

Let X, Y be trivial sets. Observe that every relation between X and Y is trivial.

We now state the proposition

(4) For every trivial set X and for every binary relation R on X such that R is non empty there exists a set x such that $R = \{\langle x, x\rangle\}$.

© 2004 University of Białystok
ISSN 1426–2630

Let X be a trivial set. Observe that every binary relation on X is trivial, reflexive, symmetric, transitive, and strongly connected.

We now state the proposition

(5) For every non empty trivial set X holds every binary relation on X is symmetric in X.

One can verify that there exists a relational structure which is non empty, strict, finite, irreflexive, and symmetric.

Let L be an irreflexive relational structure. Observe that every full relational substructure of L is irreflexive.

Let L be a symmetric relational structure. Note that every full relational substructure of L is symmetric.

One can prove the following proposition

(6) Let R be an irreflexive symmetric relational structure. Suppose $\overline{\overline{\text{the carrier of } R}} = 2$. Then there exist sets a, b such that the carrier of $R = \{a, b\}$ but the internal relation of $R = \{\langle a, b\rangle, \langle b, a\rangle\}$ or the internal relation of $R = \emptyset$.

2. Some Facts about Operations "UnionOf" and "SumOf"

Let R be a non empty relational structure and let S be a relational structure. Note that $\mathrm{UnionOf}(R, S)$ is non empty and $\mathrm{SumOf}(R, S)$ is non empty.

Let R be a relational structure and let S be a non empty relational structure. Observe that $\mathrm{UnionOf}(R, S)$ is non empty and $\mathrm{SumOf}(R, S)$ is non empty.

Let R, S be finite relational structures. One can check that $\mathrm{UnionOf}(R, S)$ is finite and $\mathrm{SumOf}(R, S)$ is finite.

Let R, S be symmetric relational structures. One can check that $\mathrm{UnionOf}(R, S)$ is symmetric and $\mathrm{SumOf}(R, S)$ is symmetric.

Let R, S be irreflexive relational structures. Observe that $\mathrm{UnionOf}(R, S)$ is irreflexive.

The following four propositions are true:

(7) Let R, S be irreflexive relational structures. Suppose the carrier of R misses the carrier of S. Then $\mathrm{SumOf}(R, S)$ is irreflexive.

(8) For all relational structures R_1, R_2 holds $\mathrm{UnionOf}(R_1, R_2) = \mathrm{UnionOf}(R_2, R_1)$ and $\mathrm{SumOf}(R_1, R_2) = \mathrm{SumOf}(R_2, R_1)$.

(9) Let G be an irreflexive relational structure and G_1, G_2 be relational structures. If $G = \mathrm{UnionOf}(G_1, G_2)$ or $G = \mathrm{SumOf}(G_1, G_2)$, then G_1 is irreflexive and G_2 is irreflexive.

(10) Let G be a non empty relational structure and H_1, H_2 be relational structures. Suppose that

(i) the carrier of H_1 misses the carrier of H_2, and

(ii) the relational structure of $G = \mathrm{UnionOf}(H_1, H_2)$ or the relational structure of $G = \mathrm{SumOf}(H_1, H_2)$.

Then H_1 is a full relational substructure of G and H_2 is a full relational substructure of G.

3. Theorems Relating to the Complement of Relational Structure

One can prove the following proposition

(11) The internal relation of $\mathrm{ComplRelStr}\,\mathrm{Necklace}\,4 = \{\langle 0, 2\rangle, \langle 2, 0\rangle, \langle 0, 3\rangle, \langle 3, 0\rangle, \langle 1, 3\rangle, \langle 3, 1\rangle\}$.

Let R be a relational structure. Note that $\mathrm{ComplRelStr}\,R$ is irreflexive.

Let R be a symmetric relational structure. Note that $\mathrm{ComplRelStr}\,R$ is symmetric.

Next we state several propositions:

(12) For every relational structure R holds the internal relation of R misses the internal relation of $\mathrm{ComplRelStr}\,R$.

(13) For every relational structure R holds $\mathrm{id}_{\text{the carrier of } R}$ misses the internal relation of $\mathrm{ComplRelStr}\,R$.

(14) Let G be a relational structure. Then $[:$the carrier of G, the carrier of $G:] = \mathrm{id}_{\text{the carrier of } G} \cup$ the internal relation of $G \cup$ the internal relation of $\mathrm{ComplRelStr}\,G$.

(15) For every strict irreflexive relational structure G such that G is trivial holds $\mathrm{ComplRelStr}\,G = G$.

(16) For every strict irreflexive relational structure G holds $\mathrm{ComplRelStr}\,\mathrm{ComplRelStr}\,G = G$.

(17) For all relational structures G_1, G_2 such that the carrier of G_1 misses the carrier of G_2 holds $\mathrm{ComplRelStr}\,\mathrm{UnionOf}(G_1, G_2) = \mathrm{SumOf}(\mathrm{ComplRelStr}\,G_1, \mathrm{ComplRelStr}\,G_2)$.

(18) For all relational structures G_1, G_2 such that the carrier of G_1 misses the carrier of G_2 holds $\mathrm{ComplRelStr}\,\mathrm{SumOf}(G_1, G_2) = \mathrm{UnionOf}(\mathrm{ComplRelStr}\,G_1, \mathrm{ComplRelStr}\,G_2)$.

(19) Let G be a relational structure and H be a full relational substructure of G. Then the internal relation of $\mathrm{ComplRelStr}\,H =$ (the internal relation of $\mathrm{ComplRelStr}\,G$) $|^2$ (the carrier of $\mathrm{ComplRelStr}\,H$).

(20) Let G be a non empty irreflexive relational structure, x be an element of the carrier of G, and x' be an element of the carrier of $\mathrm{ComplRelStr}\,G$. If $x = x'$, then $\mathrm{ComplRelStr}\,\mathrm{sub}(\Omega_G \setminus \{x\}) = \mathrm{sub}(\Omega_{\mathrm{ComplRelStr}\,G} \setminus \{x'\})$.

4. Another Facts Relating to Operation "embeds"

Let us observe that every non empty relational structure which is trivial and strict is also N-free.

The following propositions are true:

(21) Let R be a reflexive antisymmetric relational structure and S be a relational structure. Then there exists a map f from R into S such that for all elements x, y of the carrier of R holds $\langle x, y\rangle \in$ the internal relation of R iff $\langle f(x), f(y)\rangle \in$ the internal relation of S if and only if S embeds R.

(22) Let G be a non empty relational structure and H be a non empty full relational substructure of G. Then G embeds H.

(23) Let G be a non empty relational structure and H be a non empty full relational substructure of G. If G is N-free, then H is N-free.

(24) For every non empty irreflexive relational structure G holds G embeds Necklace 4 iff ComplRelStr G embeds Necklace 4.

(25) For every non empty irreflexive relational structure G holds G is N-free iff ComplRelStr G is N-free.

5. Connected Graphs

Let R be a relational structure. A path of R is a reduction sequence w.r.t. the internal relation of R.

Let R be a relational structure. We say that R is path-connected if and only if the condition (Def. 1) is satisfied.

(Def. 1) Let x, y be sets. Suppose $x \in$ the carrier of R and $y \in$ the carrier of R and $x \neq y$. Then the internal relation of R reduces x to y or the internal relation of R reduces y to x.

One can check that every relational structure which is empty is also path-connected.

One can check that every non empty relational structure which is connected is also path-connected.

We now state the proposition

(26) Let R be a non empty transitive reflexive relational structure and x, y be elements of R. Suppose the internal relation of R reduces x to y. Then $\langle x, y\rangle \in$ the internal relation of R.

One can check that every non empty transitive reflexive relational structure which is path-connected is also connected.

Next we state the proposition

(27) Let R be a symmetric relational structure and x, y be sets. Suppose $x \in$ the carrier of R and $y \in$ the carrier of R. Suppose the internal relation of R reduces x to y. Then the internal relation of R reduces y to x.

Let R be a symmetric relational structure. Let us observe that R is path-connected if and only if the condition (Def. 2) is satisfied.

(Def. 2) Let x, y be sets. Suppose $x \in$ the carrier of R and $y \in$ the carrier of R and $x \neq y$. Then the internal relation of R reduces x to y.

Let R be a relational structure and let x be an element of R. The functor component(x) yielding a subset of R is defined as follows:

(Def. 3) $\text{component}(x) = [x]_{\text{EqCl(the internal relation of } R)}$.

Next we state the proposition

(28) For every non empty relational structure R and for every element x of R holds $x \in \text{component}(x)$.

Let R be a non empty relational structure and let x be an element of R. Note that component(x) is non empty.

Next we state a number of propositions:

(29) Let R be a relational structure, x be an element of R, and y be a set. If $y \in \text{component}(x)$, then $\langle x, y \rangle \in \text{EqCl(the internal relation of } R)$.

(30) Let R be a relational structure, x be an element of R, and A be a set. Then $A = \text{component}(x)$ if and only if for every set y holds $y \in A$ iff $\langle x, y \rangle \in \text{EqCl(the internal relation of } R)$.

(31) Let R be a non empty irreflexive symmetric relational structure. Suppose R is not path-connected. Then there exist non empty strict irreflexive symmetric relational structures G_1, G_2 such that the carrier of G_1 misses the carrier of G_2 and the relational structure of $R = \text{UnionOf}(G_1, G_2)$.

(32) Let R be a non empty irreflexive symmetric relational structure. Suppose $\text{ComplRelStr}\, R$ is not path-connected. Then there exist non empty strict irreflexive symmetric relational structures G_1, G_2 such that the carrier of G_1 misses the carrier of G_2 and the relational structure of $R = \text{SumOf}(G_1, G_2)$.

(33) For every irreflexive relational structure G such that $G \in \text{FinRelStrSp}$ holds $\text{ComplRelStr}\, G \in \text{FinRelStrSp}$.

(34) Let R be an irreflexive symmetric relational structure. Suppose $\overline{\overline{\text{the carrier of } R}} = 2$ and the carrier of $R \in \mathbf{U}_0$. Then the relational structure of $R \in \text{FinRelStrSp}$.

(35) For every relational structure R such that $R \in \text{FinRelStrSp}$ holds R is symmetric.

(36) Let G be a relational structure, H_1, H_2 be non empty relational structures, x be an element of the carrier of H_1, and y be an element of the

carrier of H_2. Suppose $G = \mathrm{UnionOf}(H_1, H_2)$ and the carrier of H_1 misses the carrier of H_2. Then $\langle x, y\rangle \notin$ the internal relation of G.

(37) Let G be a relational structure, H_1, H_2 be non empty relational structures, x be an element of the carrier of H_1, and y be an element of the carrier of H_2. If $G = \mathrm{SumOf}(H_1, H_2)$, then $\langle x, y\rangle \notin$ the internal relation of $\mathrm{ComplRelStr}\, G$.

(38) Let G be a non empty symmetric relational structure, x be an element of the carrier of G, and R_1, R_2 be non empty relational structures. Suppose the carrier of R_1 misses the carrier of R_2 and $\mathrm{sub}(\Omega_G \setminus \{x\}) = \mathrm{UnionOf}(R_1, R_2)$ and G is path-connected. Then there exists an element b of the carrier of R_1 such that $\langle b, x\rangle \in$ the internal relation of G.

(39) Let G be a non empty symmetric irreflexive relational structure, a, b, c, d be elements of the carrier of G, and Z be a subset of the carrier of G. Suppose that $Z = \{a, b, c, d\}$ and a, b, c, d are mutually different and $\langle a, b\rangle \in$ the internal relation of G and $\langle b, c\rangle \in$ the internal relation of G and $\langle c, d\rangle \in$ the internal relation of G and $\langle a, c\rangle \notin$ the internal relation of G and $\langle a, d\rangle \notin$ the internal relation of G and $\langle b, d\rangle \notin$ the internal relation of G. Then $\mathrm{sub}(Z)$ embeds Necklace 4.

(40) Let G be a non empty irreflexive symmetric relational structure, x be an element of the carrier of G, and R_1, R_2 be non empty relational structures. Suppose that

(i) the carrier of R_1 misses the carrier of R_2,

(ii) $\mathrm{sub}(\Omega_G \setminus \{x\}) = \mathrm{UnionOf}(R_1, R_2)$,

(iii) G is non trivial and path-connected, and

(iv) $\mathrm{ComplRelStr}\, G$ is path-connected.

Then G embeds Necklace 4.

(41) Let G be a non empty strict finite irreflexive symmetric relational structure. Suppose G is N-free and the carrier of $G \in \mathbf{U}_0$. Then the relational structure of $G \in$ FinRelStrSp .

Acknowledgments

The author would like to thank Professor Grzegorz Bancerek for many helpful conversations during the preparation of this paper, which allowed me to publish it and for aquainting me with new search engine in MML (MML Query). I am also very grateful to Professor Andrzej Trybulec, Adam Naumowicz, and Adam Grabowski for their valuable advice.

References

[1] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(**2**):377–382, 1990.

[2] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(**1**):91–96, 1990.

[3] Grzegorz Bancerek. The well ordering relations. *Formalized Mathematics*, 1(**1**):123–129, 1990.
[4] Grzegorz Bancerek. Reduction relations. *Formalized Mathematics*, 5(**4**):469–478, 1996.
[5] Grzegorz Bancerek. Bounds in posets and relational substructures. *Formalized Mathematics*, 6(**1**):81–91, 1997.
[6] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(**1**):107–114, 1990.
[7] Józef Białas. Group and field definitions. *Formalized Mathematics*, 1(**3**):433–439, 1990.
[8] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(**1**):55–65, 1990.
[9] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(**1**):153–164, 1990.
[10] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Formalized Mathematics*, 1(**3**):521–527, 1990.
[11] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(**2**):357–367, 1990.
[12] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(**1**):47–53, 1990.
[13] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(**1**):165–167, 1990.
[14] Wojciech Leończuk and Krzysztof Prażmowski. Incidence projective spaces. *Formalized Mathematics*, 2(**2**):225–232, 1991.
[15] Robert Milewski. Lattice of congruences in many sorted algebra. *Formalized Mathematics*, 5(**4**):479–483, 1996.
[16] Bogdan Nowak and Grzegorz Bancerek. Universal classes. *Formalized Mathematics*, 1(**3**):595–600, 1990.
[17] Beata Padlewska. Families of sets. *Formalized Mathematics*, 1(**1**):147–152, 1990.
[18] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(**1**):223–230, 1990.
[19] Konrad Raczkowski and Paweł Sadowski. Equivalence relations and classes of abstraction. *Formalized Mathematics*, 1(**3**):441–444, 1990.
[20] Krzysztof Retel. The class of series – parallel graphs. Part I. *Formalized Mathematics*, 11(**1**):99–103, 2003.
[21] Krzysztof Retel. The class of series-parallel graphs. Part II. *Formalized Mathematics*, 11(**3**):289–291, 2003.
[22] Stephan Thomasse. On better-quasi-ordering countable series-parallel orders. *Transactions of the American Mathematical Society*, 352(6):2491–2505, 2000.
[23] Andrzej Trybulec. Binary operations applied to functions. *Formalized Mathematics*, 1(**2**):329–334, 1990.
[24] Andrzej Trybulec. Enumerated sets. *Formalized Mathematics*, 1(**1**):25–34, 1990.
[25] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(**1**):9–11, 1990.
[26] Wojciech A. Trybulec. Groups. *Formalized Mathematics*, 1(**5**):821–827, 1990.
[27] Wojciech A. Trybulec. Partially ordered sets. *Formalized Mathematics*, 1(**2**):313–319, 1990.
[28] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(**1**):67–71, 1990.
[29] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(**1**):73–83, 1990.
[30] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(**1**):181–186, 1990.
[31] Edmund Woronowicz and Anna Zalewska. Properties of binary relations. *Formalized Mathematics*, 1(**1**):85–89, 1990.

Received February 3, 2004