FORMALIZED MATHEMATICS
Volume 12, Number 2, 2004
University of Białystok

The Exponential Function on Banach Algebra

Yasunari Shidama
Shinshu University
Nagano

Summary. In this article, the basic properties of the exponential function on Banach algebra are described.

MML Identifier: LOPBAN_4.

The notation and terminology used here are introduced in the following papers: [17], [19], [20], [3], [4], [2], [16], [5], [1], [18], [9], [11], [12], [8], [6], [7], [13], [10], [21], [14], and [15].

For simplicity, we use the following convention: X denotes a Banach algebra, p denotes a real number, w, z, z_1, z_2 denote elements of X, k, l, m, n denote natural numbers, s_1, s_2, s_3, s, s' denote sequences of X, and r_1 denotes a sequence of real numbers.

Let X be a non empty normed algebra structure and let x, y be elements of X. We say that x, y are commutative if and only if:

(Def. 1) $x \cdot y = y \cdot x$.

Let us note that the predicate x, y are commutative is symmetric.

Next we state a number of propositions:

(1) If s_2 is convergent and s_3 is convergent and $\lim(s_2 - s_3) = 0_X$, then $\lim s_2 = \lim s_3$.

(2) For every z such that for every natural number n holds $s(n) = z$ holds $\lim s = z$.

(3) If s is convergent and s' is convergent, then $s \cdot s'$ is convergent.

(4) If s is convergent, then $z \cdot s$ is convergent.

(5) If s is convergent, then $s \cdot z$ is convergent.

© 2004 University of Białystok
ISSN 1426–2630

(6) If s is convergent, then $\lim(z \cdot s) = z \cdot \lim s$.

(7) If s is convergent, then $\lim(s \cdot z) = \lim s \cdot z$.

(8) If s is convergent and s' is convergent, then $\lim(s \cdot s') = \lim s \cdot \lim s'$.

(9) $(\sum_{\alpha=0}^{\kappa}(z \cdot s_1)(\alpha))_{\kappa \in \mathbb{N}} = z \cdot (\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa \in \mathbb{N}}$ and $(\sum_{\alpha=0}^{\kappa}(s_1 \cdot z)(\alpha))_{\kappa \in \mathbb{N}} = (\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa \in \mathbb{N}} \cdot z$.

(10) $\|(\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa \in \mathbb{N}}(k)\| \leqslant (\sum_{\alpha=0}^{\kappa}\|s_1\|(\alpha))_{\kappa \in \mathbb{N}}(k)$.

(11) If for every n such that $n \leqslant m$ holds $s_2(n) = s_3(n)$, then $(\sum_{\alpha=0}^{\kappa}(s_2)(\alpha))_{\kappa \in \mathbb{N}}(m) = (\sum_{\alpha=0}^{\kappa}(s_3)(\alpha))_{\kappa \in \mathbb{N}}(m)$.

(12) If for every n holds $\|s_1(n)\| \leqslant r_1(n)$ and r_1 is convergent and $\lim r_1 = 0$, then s_1 is convergent and $\lim s_1 = 0_X$.

Let us consider X and let z be an element of X. The functor $z\,\mathrm{ExpSeq}$ yielding a sequence of X is defined as follows:

(Def. 2) For every n holds $z\,\mathrm{ExpSeq}(n) = \frac{1}{n!} \cdot z_{\mathbb{N}}^{n}$.

The scheme *ExNormSpace CASE* deals with a non empty Banach algebra $\mathcal{A}$ and a binary functor $\mathcal{F}$ yielding a point of $\mathcal{A}$, and states that:

> For every k there exists a sequence s_1 of $\mathcal{A}$ such that for every n holds if $n \leqslant k$, then $s_1(n) = \mathcal{F}(k, n)$ and if $n > k$, then $s_1(n) = 0_{\mathcal{A}}$

for all values of the parameters.

Next we state the proposition

(13) For every k such that $0 < k$ holds $(k -' 1)! \cdot k = k!$ and for all m, k such that $k \leqslant m$ holds $(m -' k)! \cdot ((m + 1) - k) = ((m + 1) -' k)!$.

Let n be a natural number. The functor $\mathrm{Coef}\,n$ yields a sequence of real numbers and is defined by:

(Def. 3) For every natural number k holds if $k \leqslant n$, then $(\mathrm{Coef}\,n)(k) = \frac{n!}{k! \cdot (n -' k)!}$ and if $k > n$, then $(\mathrm{Coef}\,n)(k) = 0$.

Let n be a natural number. The functor $\mathrm{Coef_e}\,n$ yielding a sequence of real numbers is defined by:

(Def. 4) For every natural number k holds if $k \leqslant n$, then $(\mathrm{Coef_e}\,n)(k) = \frac{1}{k! \cdot (n -' k)!}$ and if $k > n$, then $(\mathrm{Coef_e}\,n)(k) = 0$.

Let us consider X, s_1. The functor $\mathrm{Shift}\,s_1$ yielding a sequence of X is defined as follows:

(Def. 5) $(\mathrm{Shift}\,s_1)(0) = 0_X$ and for every natural number k holds $(\mathrm{Shift}\,s_1)(k + 1) = s_1(k)$.

Let us consider n, let us consider X, and let z, w be elements of X. The functor $\mathrm{Expan}(n, z, w)$ yields a sequence of X and is defined by:

(Def. 6) For every natural number k holds if $k \leqslant n$, then $(\mathrm{Expan}(n, z, w))(k) = (\mathrm{Coef}\,n)(k) \cdot z_{\mathbb{N}}^{k} \cdot w_{\mathbb{N}}^{n -' k}$ and if $n < k$, then $(\mathrm{Expan}(n, z, w))(k) = 0_X$.

Let us consider n, let us consider X, and let z, w be elements of X. The functor $\mathrm{Expan_e}(n, z, w)$ yields a sequence of X and is defined as follows:

(Def. 7) For every natural number k holds if $k \leqslant n$, then $(\mathrm{Expan_e}(n, z, w))(k) = (\mathrm{Coef_e}\, n)(k) \cdot z_{\mathbb{N}}^{k} \cdot w_{\mathbb{N}}^{n-'k}$ and if $n < k$, then $(\mathrm{Expan_e}(n, z, w))(k) = 0_X$.

Let us consider n, let us consider X, and let z, w be elements of X. The functor $\mathrm{Alfa}(n, z, w)$ yields a sequence of X and is defined as follows:

(Def. 8) For every natural number k holds if $k \leqslant n$, then $(\mathrm{Alfa}(n, z, w))(k) = z\,\mathrm{ExpSeq}(k) \cdot (\sum_{\alpha=0}^{\kappa} w\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(n -' k)$ and if $n < k$, then $(\mathrm{Alfa}(n, z, w))(k) = 0_X$.

Let us consider X, let z, w be elements of X, and let n be a natural number. The functor $\mathrm{Conj}(n, z, w)$ yields a sequence of X and is defined by:

(Def. 9) For every natural number k holds if $k \leqslant n$, then $(\mathrm{Conj}(n, z, w))(k) = z\,\mathrm{ExpSeq}(k) \cdot ((\sum_{\alpha=0}^{\kappa} w\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(n) - (\sum_{\alpha=0}^{\kappa} w\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(n -' k))$ and if $n < k$, then $(\mathrm{Conj}(n, z, w))(k) = 0_X$.

One can prove the following propositions:

(14) $z\,\mathrm{ExpSeq}(n+1) = \frac{1}{n+1} \cdot z \cdot z\,\mathrm{ExpSeq}(n)$ and $z\,\mathrm{ExpSeq}(0) = \mathbf{1}_X$ and $\|z\,\mathrm{ExpSeq}(n)\| \leqslant \|z\|\,\mathrm{ExpSeq}(n)$.

(15) If $0 < k$, then $(\mathrm{Shift}\, s_1)(k) = s_1(k -' 1)$.

(16) $(\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}}(k) = (\sum_{\alpha=0}^{\kappa}(\mathrm{Shift}\, s_1)(\alpha))_{\kappa\in\mathbb{N}}(k) + s_1(k)$.

(17) For all z, w such that z, w are commutative holds $(z + w)_{\mathbb{N}}^{n} = (\sum_{\alpha=0}^{\kappa}(\mathrm{Expan}(n, z, w))(\alpha))_{\kappa\in\mathbb{N}}(n)$.

(18) $\mathrm{Expan_e}(n, z, w) = \frac{1}{n!} \cdot \mathrm{Expan}(n, z, w)$.

(19) For all z, w such that z, w are commutative holds $\frac{1}{n!} \cdot (z + w)_{\mathbb{N}}^{n} = (\sum_{\alpha=0}^{\kappa}(\mathrm{Expan_e}(n, z, w))(\alpha))_{\kappa\in\mathbb{N}}(n)$.

(20) $0_X\,\mathrm{ExpSeq}$ is norm-summable and $\sum(0_X\,\mathrm{ExpSeq}) = \mathbf{1}_X$.

Let us consider X and let z be an element of X. Observe that $z\,\mathrm{ExpSeq}$ is norm-summable.

Next we state a number of propositions:

(21) $z\,\mathrm{ExpSeq}(0) = \mathbf{1}_X$ and $(\mathrm{Expan}(0, z, w))(0) = \mathbf{1}_X$.

(22) If $l \leqslant k$, then $(\mathrm{Alfa}(k+1, z, w))(l) = (\mathrm{Alfa}(k, z, w))(l) + (\mathrm{Expan_e}(k+1, z, w))(l)$.

(23) $(\sum_{\alpha=0}^{\kappa}(\mathrm{Alfa}(k+1, z, w))(\alpha))_{\kappa\in\mathbb{N}}(k) = (\sum_{\alpha=0}^{\kappa}(\mathrm{Alfa}(k, z, w))(\alpha))_{\kappa\in\mathbb{N}}(k) + (\sum_{\alpha=0}^{\kappa}(\mathrm{Expan_e}(k+1, z, w))(\alpha))_{\kappa\in\mathbb{N}}(k)$.

(24) $z\,\mathrm{ExpSeq}(k) = (\mathrm{Expan_e}(k, z, w))(k)$.

(25) For all z, w such that z, w are commutative holds $(\sum_{\alpha=0}^{\kappa} z + w\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(n) = (\sum_{\alpha=0}^{\kappa}(\mathrm{Alfa}(n, z, w))(\alpha))_{\kappa\in\mathbb{N}}(n)$.

(26) For all z, w such that z, w are commutative holds $(\sum_{\alpha=0}^{\kappa} z\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(k) \cdot (\sum_{\alpha=0}^{\kappa} w\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(k) - (\sum_{\alpha=0}^{\kappa} z + w\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(k) = (\sum_{\alpha=0}^{\kappa}(\mathrm{Conj}(k, z, w))(\alpha))_{\kappa\in\mathbb{N}}(k)$.

(27) $0 \leqslant \|z\|\,\mathrm{ExpSeq}(n)$.

(28) $\|(\sum_{\alpha=0}^{\kappa} z\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(k)\| \leqslant (\sum_{\alpha=0}^{\kappa}\|z\|\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(k)$ and $(\sum_{\alpha=0}^{\kappa}\|z\|\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(k) \leqslant \sum(\|z\|\,\mathrm{ExpSeq})$ and $\|(\sum_{\alpha=0}^{\kappa} z\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(k)\| \leqslant \sum(\|z\|\,\mathrm{ExpSeq})$.

(29) $1 \leqslant \sum(\|z\|\,\mathrm{ExpSeq})$.

(30) $|(\sum_{\alpha=0}^{\kappa}\|z\|\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(n)| = (\sum_{\alpha=0}^{\kappa}\|z\|\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(n)$ and if $n \leqslant m$, then $|(\sum_{\alpha=0}^{\kappa}\|z\|\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(m) - (\sum_{\alpha=0}^{\kappa}\|z\|\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(n)| = (\sum_{\alpha=0}^{\kappa}\|z\|\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(m) - (\sum_{\alpha=0}^{\kappa}\|z\|\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(n)$.

(31) $|(\sum_{\alpha=0}^{\kappa}\|\mathrm{Conj}(k,z,w)\|(\alpha))_{\kappa\in\mathbb{N}}(n)| = (\sum_{\alpha=0}^{\kappa}\|\mathrm{Conj}(k,z,w)\|(\alpha))_{\kappa\in\mathbb{N}}(n)$.

(32) For every real number p such that $p > 0$ there exists n such that for every k such that $n \leqslant k$ holds $|(\sum_{\alpha=0}^{\kappa}\|\mathrm{Conj}(k,z,w)\|(\alpha))_{\kappa\in\mathbb{N}}(k)| < p$.

(33) For every s_1 such that for every k holds $s_1(k) = (\sum_{\alpha=0}^{\kappa}(\mathrm{Conj}(k,z,w))(\alpha))_{\kappa\in\mathbb{N}}(k)$ holds s_1 is convergent and $\lim s_1 = 0_X$.

Let X be a Banach algebra. The functor $\exp X$ yielding a function from the carrier of X into the carrier of X is defined by:

(Def. 10) For every element z of the carrier of X holds $(\exp X)(z) = \sum(z\,\mathrm{ExpSeq})$.

Let us consider X, z. The functor $\exp z$ yields an element of X and is defined by:

(Def. 11) $\exp z = (\exp X)(z)$.

One can prove the following propositions:

(34) For every z holds $\exp z = \sum(z\,\mathrm{ExpSeq})$.

(35) Let given z_1, z_2. Suppose z_1, z_2 are commutative. Then $\exp(z_1 + z_2) = \exp z_1 \cdot \exp z_2$ and $\exp(z_2 + z_1) = \exp z_2 \cdot \exp z_1$ and $\exp(z_1 + z_2) = \exp(z_2 + z_1)$ and $\exp z_1$, $\exp z_2$ are commutative.

(36) For all z_1, z_2 such that z_1, z_2 are commutative holds $z_1 \cdot \exp z_2 = \exp z_2 \cdot z_1$.

(37) $\exp(0_X) = \mathbf{1}_X$.

(38) $\exp z \cdot \exp(-z) = \mathbf{1}_X$ and $\exp(-z) \cdot \exp z = \mathbf{1}_X$.

(39) $\exp z$ is invertible and $(\exp z)^{-1} = \exp(-z)$ and $\exp(-z)$ is invertible and $(\exp(-z))^{-1} = \exp z$.

(40) For every z and for all real numbers s, t holds $s \cdot z$, $t \cdot z$ are commutative.

(41) Let given z and s, t be real numbers. Then $\exp(s \cdot z) \cdot \exp(t \cdot z) = \exp((s+t) \cdot z)$ and $\exp(t \cdot z) \cdot \exp(s \cdot z) = \exp((t+s) \cdot z)$ and $\exp((s+t) \cdot z) = \exp((t+s) \cdot z)$ and $\exp(s \cdot z)$, $\exp(t \cdot z)$ are commutative.

References

[1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(**1**):41–46, 1990.

[2] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(**1**):91–96, 1990.

[3] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(**1**):55–65, 1990.

[4] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(**1**):153–164, 1990.

[5] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(**1**):35–40, 1990.
[6] Jarosław Kotowicz. Convergent sequences and the limit of sequences. *Formalized Mathematics*, 1(**2**):273–275, 1990.
[7] Jarosław Kotowicz. Monotone real sequences. Subsequences. *Formalized Mathematics*, 1(**3**):471–475, 1990.
[8] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(**2**):269–272, 1990.
[9] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Formalized Mathematics*, 1(**2**):335–342, 1990.
[10] Takaya Nishiyama and Yasuho Mizuhara. Binary arithmetics. *Formalized Mathematics*, 4(**1**):83–86, 1993.
[11] Jan Popiołek. Some properties of functions modul and signum. *Formalized Mathematics*, 1(**2**):263–264, 1990.
[12] Jan Popiołek. Real normed space. *Formalized Mathematics*, 2(**1**):111–115, 1991.
[13] Konrad Raczkowski and Andrzej Nędzusiak. Series. *Formalized Mathematics*, 2(**4**):449–452, 1991.
[14] Yasunari Shidama. The Banach algebra of bounded linear operators. *Formalized Mathematics*, 12(**2**):103–108, 2004.
[15] Yasunari Shidama. The series on Banach algebra. *Formalized Mathematics*, 12(**2**):131–138, 2004.
[16] Andrzej Trybulec. Subsets of complex numbers. *To appear in Formalized Mathematics*.
[17] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(**1**):9–11, 1990.
[18] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(**2**):291–296, 1990.
[19] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(**1**):67–71, 1990.
[20] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(**1**):73–83, 1990.
[21] Yuguang Yang and Yasunari Shidama. Trigonometric functions and existence of circle ratio. *Formalized Mathematics*, 7(**2**):255–263, 1998.

Received February 13, 2004