FORMALIZED MATHEMATICS
Volume 13, Number 3, Pages 383–387
University of Białystok, 2005

Formulas and Identities of Inverse Hyperbolic Functions

Fuguo Ge
Qingdao University of Science and Technology
China

Xiquan Liang
Qingdao University of Science and Technology
China

Yuzhong Ding
Qingdao University of Science and Technology
China

Summary. This article describes definitions of inverse hyperbolic functions and their main properties, as well as some addition formulas with hyperbolic functions.

MML identifier: SIN_COS7, version: 7.5.01 4.39.921

The papers [1], [8], [4], [2], [9], [3], [6], [5], and [7] provide the terminology and notation for this paper.

1. Preliminaries

In this paper x, y, t denote real numbers.

Next we state a number of propositions:

(1) If $x > 0$, then $\frac{1}{x} = x^{-1}$.

(2) If $x > 1$, then $(\frac{\sqrt{x^2-1}}{x})^2 < 1$.

(3) $(\frac{x}{\sqrt{x^2+1}})^2 < 1$.

(4) $\sqrt{x^2+1} > 0$.

(5) $\sqrt{x^2+1} + x > 0$.

© 2005 University of Białystok
ISSN 1426–2630

(6) If $y \geq 0$ and $x \geq 1$, then $\frac{x+1}{y} \geq 0$.

(7) If $y \geq 0$ and $x \geq 1$, then $\frac{x-1}{y} \geq 0$.

(8) If $x \geq 1$, then $\sqrt{\frac{x+1}{2}} \geq 1$.

(9) If $y \geq 0$ and $x \geq 1$, then $\frac{x^2-1}{y} \geq 0$.

(10) If $x \geq 1$, then $\sqrt{\frac{x+1}{2}} + \sqrt{\frac{x-1}{2}} > 0$.

(11) If $x^2 < 1$, then $x + 1 > 0$ and $1 - x > 0$.

(12) If $x \neq 1$, then $(1 - x)^2 > 0$.

(13) If $x^2 < 1$, then $\frac{x^2+1}{1-x^2} \geq 0$.

(14) If $x^2 < 1$, then $(\frac{2 \cdot x}{1+x^2})^2 < 1$.

(15) If $0 < x$ and $x < 1$, then $\frac{1+x}{1-x} > 0$.

(16) If $0 < x$ and $x < 1$, then $x^2 < 1$.

(17) If $0 < x$ and $x < 1$, then $\frac{1}{\sqrt{1-x^2}} > 1$.

(18) If $0 < x$ and $x < 1$, then $\frac{2 \cdot x}{1-x^2} > 0$.

(19) If $0 < x$ and $x < 1$, then $0 < (1 - x^2)^2$.

(20) If $0 < x$ and $x < 1$, then $\frac{1+x^2}{1-x^2} > 1$.

(21) If $1 < x^2$, then $(\frac{1}{x})^2 < 1$.

(22) If $0 < x$ and $x \leq 1$, then $1 - x^2 \geq 0$.

(23) If $1 \leq x$, then $0 < x + \sqrt{x^2 - 1}$.

(24) If $1 \leq x$ and $1 \leq y$, then $0 \leq x \cdot \sqrt{y^2 - 1} + y \cdot \sqrt{x^2 - 1}$.

(25) If $1 \leq x$ and $1 \leq y$ and $|y| \leq |x|$, then $0 < y - \sqrt{y^2 - 1}$.

(26) If $1 \leq x$ and $1 \leq y$ and $|y| \leq |x|$, then $0 \leq y \cdot \sqrt{x^2 - 1} - x \cdot \sqrt{y^2 - 1}$.

(27) If $x^2 < 1$ and $y^2 < 1$, then $x \cdot y \neq -1$.

(28) If $x^2 < 1$ and $y^2 < 1$, then $x \cdot y \neq 1$.

(29) If $x \neq 0$, then $\exp x \neq 1$.

(30) If $0 \neq x$, then $(\exp x)^2 - 1 \neq 0$.

(31) If $0 < t$, then $\frac{t^2-1}{t^2+1} < 1$.

(32) If $-1 < t$ and $t < 1$, then $0 < \frac{t+1}{1-t}$.

2. Formulas and Identities of Inverse Hyperbolic Functions

Let x be a real number. The functor $\sinh' x$ yields a real number and is defined by:

(Def. 1) $\sinh' x = \log_e(x + \sqrt{x^2 + 1})$.

Let x be a real number. The functor $\cosh'_1 x$ yielding a real number is defined by:

(Def. 2) $\cosh'_1 x = \log_e(x + \sqrt{x^2 - 1})$.

Let x be a real number. The functor $\cosh'_2 x$ yields a real number and is defined by:

(Def. 3) $\cosh'_2 x = -\log_e(x + \sqrt{x^2 - 1})$.

Let x be a real number. The functor $\tanh' x$ yields a real number and is defined by:

(Def. 4) $\tanh' x = \frac{1}{2} \cdot \log_e(\frac{1+x}{1-x})$.

Let x be a real number. The functor $\coth' x$ yielding a real number is defined as follows:

(Def. 5) $\coth' x = \frac{1}{2} \cdot \log_e(\frac{x+1}{x-1})$.

Let x be a real number. The functor $\operatorname{sech}'_1 x$ yields a real number and is defined by:

(Def. 6) $\operatorname{sech}'_1 x = \log_e(\frac{1+\sqrt{1-x^2}}{x})$.

Let x be a real number. The functor $\operatorname{sech}'_2 x$ yielding a real number is defined as follows:

(Def. 7) $\operatorname{sech}'_2 x = -\log_e(\frac{1+\sqrt{1-x^2}}{x})$.

Let x be a real number. The functor $\operatorname{csch}' x$ yielding a real number is defined by:

(Def. 8)(i) $\operatorname{csch}' x = \log_e(\frac{1+\sqrt{1+x^2}}{x})$ if $0 < x$,

(ii) $\operatorname{csch}' x = \log_e(\frac{1-\sqrt{1+x^2}}{x})$ if $x < 0$,

(iii) $x < 0$, otherwise.

The following propositions are true:

(33) If $0 \leq x$, then $\sinh' x = \cosh'_1 \sqrt{x^2 + 1}$.

(34) If $x < 0$, then $\sinh' x = \cosh'_2 \sqrt{x^2 + 1}$.

(35) $\sinh' x = \tanh'(\frac{x}{\sqrt{x^2+1}})$.

(36) If $x \geq 1$, then $\cosh'_1 x = \sinh' \sqrt{x^2 - 1}$.

(37) If $x > 1$, then $\cosh'_1 x = \tanh'(\frac{\sqrt{x^2-1}}{x})$.

(38) If $x \geq 1$, then $\cosh'_1 x = 2 \cdot \cosh'_1 \sqrt{\frac{x+1}{2}}$.

(39) If $x \geq 1$, then $\cosh'_2 x = 2 \cdot \cosh'_2 \sqrt{\frac{x+1}{2}}$.

(40) If $x \geq 1$, then $\cosh'_1 x = 2 \cdot \sinh' \sqrt{\frac{x-1}{2}}$.

(41) If $x^2 < 1$, then $\tanh' x = \sinh'(\frac{x}{\sqrt{1-x^2}})$.

(42) If $0 < x$ and $x < 1$, then $\tanh' x = \cosh'_1(\frac{1}{\sqrt{1-x^2}})$.

(43) If $x^2 < 1$, then $\tanh' x = \frac{1}{2} \cdot \sinh'(\frac{2 \cdot x}{1-x^2})$.

(44) If $x > 0$ and $x < 1$, then $\tanh' x = \frac{1}{2} \cdot \cosh'_1(\frac{1+x^2}{1-x^2})$.

(45) If $x^2 < 1$, then $\tanh' x = \frac{1}{2} \cdot \tanh'(\frac{2 \cdot x}{1+x^2})$.

(46) If $x^{\mathbf{2}} > 1$, then $\coth' x = \tanh'(\frac{1}{x})$.

(47) If $x > 0$ and $x \leq 1$, then $\operatorname{sech}'_1 x = \cosh'_1(\frac{1}{x})$.

(48) If $x > 0$ and $x \leq 1$, then $\operatorname{sech}'_2 x = \cosh'_2(\frac{1}{x})$.

(49) If $x > 0$, then $\operatorname{csch}' x = \sinh'(\frac{1}{x})$.

(50) If $x \cdot y + \sqrt{x^{\mathbf{2}} + 1} \cdot \sqrt{y^{\mathbf{2}} + 1} \geq 0$, then $\sinh' x + \sinh' y = \sinh'(x \cdot \sqrt{1 + y^{\mathbf{2}}} + y \cdot \sqrt{1 + x^{\mathbf{2}}})$.

(51) $\sinh' x - \sinh' y = \sinh'(x \cdot \sqrt{1 + y^{\mathbf{2}}} - y \cdot \sqrt{1 + x^{\mathbf{2}}})$.

(52) If $1 \leq x$ and $1 \leq y$, then $\cosh'_1 x + \cosh'_1 y = \cosh'_1(x \cdot y + \sqrt{(x^{\mathbf{2}} - 1) \cdot (y^{\mathbf{2}} - 1)})$.

(53) If $1 \leq x$ and $1 \leq y$, then $\cosh'_2 x + \cosh'_2 y = \cosh'_2(x \cdot y + \sqrt{(x^{\mathbf{2}} - 1) \cdot (y^{\mathbf{2}} - 1)})$.

(54) If $1 \leq x$ and $1 \leq y$ and $|y| \leq |x|$, then $\cosh'_1 x - \cosh'_1 y = \cosh'_1(x \cdot y - \sqrt{(x^{\mathbf{2}} - 1) \cdot (y^{\mathbf{2}} - 1)})$.

(55) If $1 \leq x$ and $1 \leq y$ and $|y| \leq |x|$, then $\cosh'_2 x - \cosh'_2 y = \cosh'_2(x \cdot y - \sqrt{(x^{\mathbf{2}} - 1) \cdot (y^{\mathbf{2}} - 1)})$.

(56) If $x^{\mathbf{2}} < 1$ and $y^{\mathbf{2}} < 1$, then $\tanh' x + \tanh' y = \tanh'(\frac{x+y}{1+x \cdot y})$.

(57) If $x^{\mathbf{2}} < 1$ and $y^{\mathbf{2}} < 1$, then $\tanh' x - \tanh' y = \tanh'(\frac{x-y}{1-x \cdot y})$.

(58) If $0 < x$ and $(\frac{x-1}{x+1})^{\mathbf{2}} < 1$, then $\log_e x = 2 \cdot \tanh'(\frac{x-1}{x+1})$.

(59) If $0 < x$ and $(\frac{x^{\mathbf{2}}-1}{x^{\mathbf{2}}+1})^{\mathbf{2}} < 1$, then $\log_e x = \tanh'(\frac{x^{\mathbf{2}}-1}{x^{\mathbf{2}}+1})$.

(60) If $1 < x$ and $1 \leq \frac{x^{\mathbf{2}}+1}{2 \cdot x}$, then $\log_e x = \cosh'_1(\frac{x^{\mathbf{2}}+1}{2 \cdot x})$.

(61) If $0 < x$ and $x < 1$ and $1 \leq \frac{x^{\mathbf{2}}+1}{2 \cdot x}$, then $\log_e x = \cosh'_2(\frac{x^{\mathbf{2}}+1}{2 \cdot x})$.

(62) If $0 < x$, then $\log_e x = \sinh'(\frac{x^{\mathbf{2}}-1}{2 \cdot x})$.

(63) If $y = \frac{1}{2} \cdot (\exp x - \exp(-x))$, then $x = \log_e(y + \sqrt{y^{\mathbf{2}} + 1})$.

(64) If $y = \frac{1}{2} \cdot (\exp x + \exp(-x))$ and $1 \leq y$, then $x = \log_e(y + \sqrt{y^{\mathbf{2}} - 1})$ or $x = -\log_e(y + \sqrt{y^{\mathbf{2}} - 1})$.

(65) If $y = \frac{\exp x - \exp(-x)}{\exp x + \exp(-x)}$, then $x = \frac{1}{2} \cdot \log_e(\frac{1+y}{1-y})$.

(66) If $y = \frac{\exp x + \exp(-x)}{\exp x - \exp(-x)}$ and $x \neq 0$, then $x = \frac{1}{2} \cdot \log_e(\frac{y+1}{y-1})$.

(67) If $y = \frac{1}{\frac{\exp x + \exp(-x)}{2}}$, then $x = \log_e(\frac{1+\sqrt{1-y^{\mathbf{2}}}}{y})$ or $x = -\log_e(\frac{1+\sqrt{1-y^{\mathbf{2}}}}{y})$.

(68) If $y = \frac{1}{\frac{\exp x - \exp(-x)}{2}}$ and $x \neq 0$, then $x = \log_e(\frac{1+\sqrt{1+y^{\mathbf{2}}}}{y})$ or $x = \log_e(\frac{1-\sqrt{1+y^{\mathbf{2}}}}{y})$.

(69) (The function cosh)$(2 \cdot x) = 1 + 2 \cdot$ (the function sinh)$(x)^{\mathbf{2}}$.

(70) (The function cosh)$(x)^{\mathbf{2}} = 1 +$ (the function sinh)$(x)^{\mathbf{2}}$.

(71) (The function sinh)$(x)^{\mathbf{2}} =$ (the function cosh)$(x)^{\mathbf{2}} - 1$.

(72) $\sinh(5 \cdot x) = 5 \cdot \sinh x + 20 \cdot (\sinh x)^3 + 16 \cdot (\sinh x)^5$.

(73) $\cosh(5 \cdot x) = (5 \cdot \cosh x - 20 \cdot (\cosh x)^3) + 16 \cdot (\cosh x)^5$.

References

[1] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(**1**):91–96, 1990.
[2] Czesław Byliński. The complex numbers. *Formalized Mathematics*, 1(**3**):507–513, 1990.
[3] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(**2**):269–272, 1990.
[4] Rafał Kwiatek. Factorial and Newton coefficients. *Formalized Mathematics*, 1(**5**):887–890, 1990.
[5] Takashi Mitsuishi and Yuguang Yang. Properties of the trigonometric function. *Formalized Mathematics*, 8(**1**):103–106, 1999.
[6] Library Committee of the Association of Mizar Users. Binary operations on numbers. *To appear in Formalized Mathematics*.
[7] Konrad Raczkowski and Andrzej Nędzusiak. Real exponents and logarithms. *Formalized Mathematics*, 2(**2**):213–216, 1991.
[8] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers. *Formalized Mathematics*, 1(**3**):445–449, 1990.
[9] Yuguang Yang and Yasunari Shidama. Trigonometric functions and existence of circle ratio. *Formalized Mathematics*, 7(**2**):255–263, 1998.

Received May 24, 2005