FORMALIZED MATHEMATICS
Volume 13, Number 4, Pages 471–475
University of Białystok, 2005

Tietze Extension Theorem

Artur Korniłowicz
Institute of Computer Science
University of Białystok
Sosnowa 64, 15-887 Białystok
Poland

Grzegorz Bancerek
Department of Theoretical
Computer Science
Białystok Technical University
Poland

Adam Naumowicz
Institute of Computer Science
University of Białystok
Akademicka 2, 15-267 Białystok
Poland

Summary. In this paper we formalize the Tietze extension theorem using as a basis the proof presented at the PlanetMath web server[1].

MML identifier: `TIETZE`, version: `7.5.01 4.39.921`

The articles [24], [26], [1], [2], [23], [11], [4], [21], [27], [5], [28], [7], [6], [17], [16], [22], [18], [20], [19], [25], [9], [10], [13], [14], [8], [12], [3], and [15] provide the notation and terminology for this paper.

We adopt the following rules: r, s denote real numbers, X denotes a set, and f, g, h denote real-yielding functions.

The following propositions are true:

(1) For all real numbers a, b, c such that $|a - b| \leq c$ holds $b - c \leq a$ and $a \leq b + c$.

(2) If $r < s$, then $]-\infty, r]$ misses $[s, +\infty[$.

(3) If $r \leq s$, then $]-\infty, r[$ misses $]s, +\infty[$.

(4) If $f \subseteq g$, then $h - f \subseteq h - g$.

(5) If $f \subseteq g$, then $f - h \subseteq g - h$.

[1] `http://planetmath.org/encyclopedia/ProofOfTietzeExtensionTheorem2.html`

© 2005 University of Białystok
ISSN 1426–2630

Let f be a real-yielding function, let r be a real number, and let X be a set. We say that f is absolutely bounded by r in X if and only if:

(Def. 1) For every set x such that $x \in X \cap \operatorname{dom} f$ holds $|f(x)| \leq r$.

Let us mention that there exists a sequence of real numbers which is summable, constant, and convergent.

We now state the proposition

(6) For every empty topological space T_1 and for every topological space T_2 holds every map from T_1 into T_2 is continuous.

Let T_1 be a topological space and let T_2 be a non empty topological space. Observe that there exists a map from T_1 into T_2 which is continuous.

We now state several propositions:

(7) For all summable sequences f, g of real numbers such that for every natural number n holds $f(n) \leq g(n)$ holds $\sum f \leq \sum g$.

(8) For every sequence f of real numbers such that f is absolutely summable holds $|\sum f| \leq \sum |f|$.

(9) Let f be a sequence of real numbers and a, r be positive real numbers. Suppose $r < 1$ and for every natural number n holds $|f(n) - f(n+1)| \leq a \cdot r^n$. Then f is convergent and for every natural number n holds $|\lim f - f(n)| \leq \frac{a \cdot r^n}{1-r}$.

(10) Let f be a sequence of real numbers and a, r be positive real numbers. Suppose $r < 1$ and for every natural number n holds $|f(n) - f(n+1)| \leq a \cdot r^n$. Then $\lim f \geq f(0) - \frac{a}{1-r}$ and $\lim f \leq f(0) + \frac{a}{1-r}$.

(11) Let X, Z be non empty sets and F be a sequence of partial functions from X into $\mathbb{R}$. Suppose Z is common for elements of F. Let a, r be positive real numbers. Suppose $r < 1$ and for every natural number n holds $F(n) - F(n+1)$ is absolutely bounded by $a \cdot r^n$ in Z. Then F is uniform-convergent on Z and for every natural number n holds $\lim_Z F - F(n)$ is absolutely bounded by $\frac{a \cdot r^n}{1-r}$ in Z.

(12) Let X, Z be non empty sets and F be a sequence of partial functions from X into $\mathbb{R}$. Suppose Z is common for elements of F. Let a, r be positive real numbers. Suppose $r < 1$ and for every natural number n holds $F(n) - F(n+1)$ is absolutely bounded by $a \cdot r^n$ in Z. Let z be an element of Z. Then $(\lim_Z F)(z) \geq F(0)(z) - \frac{a}{1-r}$ and $(\lim_Z F)(z) \leq F(0)(z) + \frac{a}{1-r}$.

(13) Let X, Z be non empty sets and F be a sequence of partial functions from X into $\mathbb{R}$. Suppose Z is common for elements of F. Let a, r be positive real numbers and f be a function from Z into $\mathbb{R}$. Suppose $r < 1$ and for every natural number n holds $F(n) - f$ is absolutely bounded by $a \cdot r^n$ in Z. Then F is point-convergent on Z and $\lim_Z F = f$.

Let S, T be topological structures, let A be an empty subset of S, and let f be a map from S into T. Note that $f \restriction A$ is empty.

Let T be a topological space and let A be a closed subset of T. Note that $T\restriction A$ is closed.

The following propositions are true:

(14) Let X, Y be non empty topological spaces, X_1, X_2 be non empty subspaces of X, f_1 be a map from X_1 into Y, and f_2 be a map from X_2 into Y. Suppose X_1 misses X_2 or $f_1\restriction(X_1 \cap X_2) = f_2\restriction(X_1 \cap X_2)$. Let x be a point of X. Then
  (i) if $x \in$ the carrier of X_1, then $(f_1 \cup f_2)(x) = f_1(x)$, and
  (ii) if $x \in$ the carrier of X_2, then $(f_1 \cup f_2)(x) = f_2(x)$.

(15) Let X, Y be non empty topological spaces, X_1, X_2 be non empty subspaces of X, f_1 be a map from X_1 into Y, and f_2 be a map from X_2 into Y. If X_1 misses X_2 or $f_1\restriction(X_1 \cap X_2) = f_2\restriction(X_1 \cap X_2)$, then $\operatorname{rng}(f_1 \cup f_2) \subseteq \operatorname{rng} f_1 \cup \operatorname{rng} f_2$.

(16) Let X, Y be non empty topological spaces, X_1, X_2 be non empty subspaces of X, f_1 be a map from X_1 into Y, and f_2 be a map from X_2 into Y. Suppose X_1 misses X_2 or $f_1\restriction(X_1 \cap X_2) = f_2\restriction(X_1 \cap X_2)$. Then for every subset A of X_1 holds $(f_1 \cup f_2)^\circ A = {f_1}^\circ A$ and for every subset A of X_2 holds $(f_1 \cup f_2)^\circ A = {f_2}^\circ A$.

(17) If $f \subseteq g$ and g is absolutely bounded by r in X, then f is absolutely bounded by r in X.

(18) If $X \subseteq \operatorname{dom} f$ or $\operatorname{dom} g \subseteq \operatorname{dom} f$ and if $f\restriction X = g\restriction X$ and if f is absolutely bounded by r in X, then g is absolutely bounded by r in X.

In the sequel T is a non empty topological space and A is a closed subset of T.

One can prove the following propositions:

(19) Suppose $r > 0$ and T is T_4. Let f be a continuous map from $T\restriction A$ into $\mathbb{R}^{\mathbf{1}}$. Suppose f is absolutely bounded by r in A. Then there exists a continuous map g from T into $\mathbb{R}^{\mathbf{1}}$ such that g is absolutely bounded by $\frac{r}{3}$ in $\operatorname{dom} g$ and $f - g$ is absolutely bounded by $\frac{2 \cdot r}{3}$ in A.

(20) Suppose that for all non empty closed subsets A, B of T such that A misses B there exists a continuous map f from T into $\mathbb{R}^{\mathbf{1}}$ such that ${f}^\circ A = \{0\}$ and ${f}^\circ B = \{1\}$. Then T is a T_4 space.

(21) Let f be a map from T into $\mathbb{R}^{\mathbf{1}}$ and x be a point of T. Then f is continuous at x if and only if for every real number e such that $e > 0$ there exists a subset H of T such that H is open and $x \in H$ and for every point y of T such that $y \in H$ holds $|f(y) - f(x)| < e$.

(22) Let F be a sequence of partial functions from the carrier of T into $\mathbb{R}$. Suppose that
  (i) F is uniform-convergent on the carrier of T, and
  (ii) for every natural number i holds $F(i)$ is a continuous map from T into

$\mathbb{R}^{\mathbf{1}}$.

Then $\lim_{\text{the carrier of } T} F$ is a continuous map from T into $\mathbb{R}^{\mathbf{1}}$.

(23) Let T be a non empty topological space, f be a map from T into $\mathbb{R}^{\mathbf{1}}$, and r be a positive real number. Then f is absolutely bounded by r in the carrier of T if and only if f is a map from T into $[-r, r]_{\mathrm{T}}$.

(24) If $f-g$ is absolutely bounded by r in X, then $g-f$ is absolutely bounded by r in X.

(25) Suppose T is T_4. Let given A and f be a map from $T\restriction A$ into $[-1, 1]_{\mathrm{T}}$. Suppose f is continuous. Then there exists a continuous map g from T into $[-1, 1]_{\mathrm{T}}$ such that $g\restriction A = f$.

(26) Suppose that for every non empty closed subset A of T and for every continuous map f from $T\restriction A$ into $[-1, 1]_{\mathrm{T}}$ there exists a continuous map g from T into $[-1, 1]_{\mathrm{T}}$ such that $g\restriction A = f$. Then T is T_4.

References

[1] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(**1**):91–96, 1990.
[2] Grzegorz Bancerek. Sequences of ordinal numbers. *Formalized Mathematics*, 1(**2**):281–290, 1990.
[3] Leszek Borys. Paracompact and metrizable spaces. *Formalized Mathematics*, 2(**4**):481–485, 1991.
[4] Czesław Byliński. The complex numbers. *Formalized Mathematics*, 1(**3**):507–513, 1990.
[5] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(**1**):55–65, 1990.
[6] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(**1**):153–164, 1990.
[7] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(**2**):357–367, 1990.
[8] Czesław Byliński and Piotr Rudnicki. Bounding boxes for compact sets in $\mathcal{E}^2$. *Formalized Mathematics*, 6(**3**):427–440, 1997.
[9] Agata Darmochwał. Compact spaces. *Formalized Mathematics*, 1(**2**):383–386, 1990.
[10] Agata Darmochwał and Yatsuka Nakamura. Metric spaces as topological spaces – fundamental concepts. *Formalized Mathematics*, 2(**4**):605–608, 1991.
[11] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(**1**):35–40, 1990.
[12] Stanisława Kanas, Adam Lecko, and Mariusz Startek. Metric spaces. *Formalized Mathematics*, 1(**3**):607–610, 1990.
[13] Zbigniew Karno. Separated and weakly separated subspaces of topological spaces. *Formalized Mathematics*, 2(**5**):665–674, 1991.
[14] Zbigniew Karno. Continuity of mappings over the union of subspaces. *Formalized Mathematics*, 3(**1**):1–16, 1992.
[15] Artur Korniłowicz. On the real valued functions. *Formalized Mathematics*, 13(**1**):181–187, 2005.
[16] Jarosław Kotowicz. Convergent sequences and the limit of sequences. *Formalized Mathematics*, 1(**2**):273–275, 1990.
[17] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(**2**):269–272, 1990.
[18] Jarosław Kotowicz. The limit of a real function at infinity. *Formalized Mathematics*, 2(**1**):17–28, 1991.
[19] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(**1**):223–230, 1990.
[20] Beata Perkowska. Functional sequence from a domain to a domain. *Formalized Mathematics*, 3(**1**):17–21, 1992.

[21] Konrad Raczkowski and Andrzej Nędzusiak. Real exponents and logarithms. *Formalized Mathematics*, 2(**2**):213–216, 1991.
[22] Konrad Raczkowski and Andrzej Nędzusiak. Series. *Formalized Mathematics*, 2(**4**):449–452, 1991.
[23] Andrzej Trybulec. Subsets of complex numbers. *To appear in Formalized Mathematics*.
[24] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(**1**):9–11, 1990.
[25] Andrzej Trybulec. A Borsuk theorem on homotopy types. *Formalized Mathematics*, 2(**4**):535–545, 1991.
[26] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(**1**):67–71, 1990.
[27] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(**1**):73–83, 1990.
[28] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(**1**):181–186, 1990.

Received September 14, 2005