FORMALIZED MATHEMATICS
Volume 11, Number 3, 2003
University of Białystok

On Some Properties of Real Hilbert Space. Part I

Hiroshi Yamazaki
Shinshu University
Nagano

Yasumasa Suzuki
Take, Yokosuka-shi
Japan

Takao Inoué
The Iida Technical High School
Nagano

Yasunari Shidama
Shinshu University
Nagano

Summary. In this paper, we first introduce the notion of summability of an infinite set of vectors of real Hilbert space, without using index sets. Further we introduce the notion of weak summability, which is weaker than that of summability. Then, several statements for summable sets and weakly summable ones are proved. In the last part of the paper, we give a necessary and sufficient condition for summability of an infinite set of vectors of real Hilbert space as our main theorem. The last theorem is due to [8].

MML Identifier: BHSP_6.

The terminology and notation used here are introduced in the following articles: [18], [21], [6], [1], [16], [9], [22], [4], [5], [7], [12], [20], [13], [14], [15], [3], [10], [17], [11], [2], [19], and [23].

1. Preliminaries

In this paper X is a real unitary space, x is a point of X, and i is a natural number.

Let us consider X. Let us assume that the addition of X is commutative and associative and the addition of X has a unity. Let Y be a finite subset of the carrier of X. The functor Setsum(Y) yielding an element of the carrier of X is defined by the condition (Def. 1).

© 2003 University of Białystok
ISSN 1426–2630

(Def. 1) There exists a finite sequence p of elements of the carrier of X such that p is one-to-one and $\operatorname{rng} p = Y$ and $\operatorname{Setsum}(Y) =$ the addition of $X \odot p$.

We now state two propositions:

(1) Let given X. Suppose the addition of X is commutative and associative and the addition of X has a unity. Let Y be a finite subset of the carrier of X and I be a function from the carrier of X into the carrier of X. Suppose $Y \subseteq \operatorname{dom} I$ and for every set x such that $x \in \operatorname{dom} I$ holds $I(x) = x$. Then $\operatorname{Setsum}(Y) = \operatorname{setopfunc}(Y,$ the carrier of X, the carrier of X, $I,$ the addition of $X)$.

(2) Let given X. Suppose the addition of X is commutative and associative and the addition of X has a unity. Let Y_1, Y_2 be finite subsets of the carrier of X. Suppose Y_1 misses Y_2. Let Z be a finite subset of the carrier of X. If $Z = Y_1 \cup Y_2$, then $\operatorname{Setsum}(Z) = \operatorname{Setsum}(Y_1) + \operatorname{Setsum}(Y_2)$.

2. Summability

Let us consider X and let Y be a subset of the carrier of X. We say that Y is summable_set if and only if the condition (Def. 2) is satisfied.

(Def. 2) There exists x such that for every real number e if $e > 0$, then there exists a finite subset Y_0 of the carrier of X such that Y_0 is non empty and $Y_0 \subseteq Y$ and for every finite subset Y_1 of the carrier of X such that $Y_0 \subseteq Y_1$ and $Y_1 \subseteq Y$ holds $\|x - \operatorname{Setsum}(Y_1)\| < e$.

Let us consider X and let Y be a subset of the carrier of X. Let us assume that Y is summable_set. The functor $\operatorname{sum} Y$ yielding a point of X is defined by the condition (Def. 3).

(Def. 3) Let e be a real number. Suppose $e > 0$. Then there exists a finite subset Y_0 of the carrier of X such that Y_0 is non empty and $Y_0 \subseteq Y$ and for every finite subset Y_1 of the carrier of X such that $Y_0 \subseteq Y_1$ and $Y_1 \subseteq Y$ holds $\|\operatorname{sum} Y - \operatorname{Setsum}(Y_1)\| < e$.

Let us consider X and let L be a linear functional in X. We say that L is Bounded if and only if:

(Def. 4) There exists a real number K such that $K > 0$ and for every x holds $|L(x)| \leqslant K \cdot \|x\|$.

Let us consider X and let Y be a subset of the carrier of X. We say that Y is weakly summable_set if and only if the condition (Def. 5) is satisfied.

(Def. 5) There exists x such that for every linear functional L in X if L is Bounded, then for every real number e such that $e > 0$ there exists a finite subset Y_0 of the carrier of X such that Y_0 is non empty and $Y_0 \subseteq Y$ and for every finite subset Y_1 of the carrier of X such that $Y_0 \subseteq Y_1$ and $Y_1 \subseteq Y$ holds $|L(x - \operatorname{Setsum}(Y_1))| < e$.

Let us consider X, let Y be a subset of the carrier of X, and let L be a functional in X. We say that Y is summable set by L if and only if the condition (Def. 6) is satisfied.

(Def. 6) There exists a real number r such that for every real number e if $e > 0$, then there exists a finite subset Y_0 of the carrier of X such that Y_0 is non empty and $Y_0 \subseteq Y$ and for every finite subset Y_1 of the carrier of X such that $Y_0 \subseteq Y_1$ and $Y_1 \subseteq Y$ holds $|r - \mathrm{setopfunc}(Y_1, \text{the carrier of } X, \mathbb{R}, L, +_{\mathbb{R}})| < e$.

Let us consider X, let Y be a subset of the carrier of X, and let L be a functional in X. Let us assume that Y is summable set by L. The functor $\mathrm{SumByfunc}(Y, L)$ yielding a real number is defined by the condition (Def. 7).

(Def. 7) Let e be a real number. Suppose $e > 0$. Then there exists a finite subset Y_0 of the carrier of X such that

(i) Y_0 is non empty,

(ii) $Y_0 \subseteq Y$, and

(iii) for every finite subset Y_1 of the carrier of X such that $Y_0 \subseteq Y_1$ and $Y_1 \subseteq Y$ holds $|\mathrm{SumByfunc}(Y, L) - \mathrm{setopfunc}(Y_1, \text{the carrier of } X, \mathbb{R}, L, +_{\mathbb{R}})| < e$.

The following propositions are true:

(3) For every subset Y of the carrier of X such that Y is summable_set holds Y is weakly summable_set.

(4) Let L be a linear functional in X and p be a finite sequence of elements of the carrier of X. Suppose $\mathrm{len}\, p \geqslant 1$. Let q be a finite sequence of elements of $\mathbb{R}$. Suppose $\mathrm{dom}\, p = \mathrm{dom}\, q$ and for every i such that $i \in \mathrm{dom}\, q$ holds $q(i) = L(p(i))$. Then $L(\text{the addition of } X \odot p) = +_{\mathbb{R}} \odot q$.

(5) Let given X. Suppose the addition of X is commutative and associative and the addition of X has a unity. Let S be a finite subset of the carrier of X. Suppose S is non empty. Let L be a linear functional in X. Then $L(\mathrm{Setsum}(S)) = \mathrm{setopfunc}(S, \text{the carrier of } X, \mathbb{R}, L, +_{\mathbb{R}})$.

(6) Let given X. Suppose the addition of X is commutative and associative and the addition of X has a unity. Let Y be a subset of the carrier of X. Suppose Y is weakly summable_set. Then there exists x such that for every linear functional L in X if L is Bounded, then for every real number e such that $e > 0$ there exists a finite subset Y_0 of the carrier of X such that Y_0 is non empty and $Y_0 \subseteq Y$ and for every finite subset Y_1 of the carrier of X such that $Y_0 \subseteq Y_1$ and $Y_1 \subseteq Y$ holds $|L(x) - \mathrm{setopfunc}(Y_1, \text{the carrier of } X, \mathbb{R}, L, +_{\mathbb{R}})| < e$.

(7) Let given X. Suppose the addition of X is commutative and associative and the addition of X has a unity. Let Y be a subset of the carrier of X. Suppose Y is weakly summable_set. Let L be a linear functional in X. If L is Bounded, then Y is summable set by L.

(8) Let given X. Suppose the addition of X is commutative and associative and the addition of X has a unity. Let Y be a subset of the carrier of X. Suppose Y is summable_set. Let L be a linear functional in X. If L is Bounded, then Y is summable set by L.

(9) For every finite subset Y of the carrier of X such that Y is non empty holds Y is summable_set.

3. Necessary and Sufficient Condition for Summability

One can prove the following proposition

(10) Let given X. Suppose the addition of X is commutative and associative and the addition of X has a unity and X is a Hilbert space. Let Y be a subset of the carrier of X. Then Y is summable_set if and only if for every real number e such that $e > 0$ there exists a finite subset Y_0 of the carrier of X such that Y_0 is non empty and $Y_0 \subseteq Y$ and for every finite subset Y_1 of the carrier of X such that Y_1 is non empty and $Y_1 \subseteq Y$ and Y_0 misses Y_1 holds $\|\mathrm{Setsum}(Y_1)\| < e$.

References

[1] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(**2**):377–382, 1990.
[2] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(**1**):107–114, 1990.
[3] Czesław Byliński. Binary operations. *Formalized Mathematics*, 1(**1**):175–180, 1990.
[4] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(**1**):55–65, 1990.
[5] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(**1**):153–164, 1990.
[6] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(**1**):47–53, 1990.
[7] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(**1**):165–167, 1990.
[8] P. R. Halmos. *Introduction to Hilbert Space*. American Mathematical Society, 1987.
[9] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(**1**):35–40, 1990.
[10] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Formalized Mathematics*, 1(**2**):335–342, 1990.
[11] Bogdan Nowak and Andrzej Trybulec. Hahn-Banach theorem. *Formalized Mathematics*, 4(**1**):29–34, 1993.
[12] Jan Popiołek. Some properties of functions modul and signum. *Formalized Mathematics*, 1(**2**):263–264, 1990.
[13] Jan Popiołek. Introduction to Banach and Hilbert spaces - part I. *Formalized Mathematics*, 2(**4**):511–516, 1991.
[14] Jan Popiołek. Introduction to Banach and Hilbert spaces - part III. *Formalized Mathematics*, 2(**4**):523–526, 1991.
[15] Jan Popiołek. Real normed space. *Formalized Mathematics*, 2(**1**):111–115, 1991.
[16] Andrzej Trybulec. Introduction to arithmetics. *To appear in Formalized Mathematics*.
[17] Andrzej Trybulec. Semilattice operations on finite subsets. *Formalized Mathematics*, 1(**2**):369–376, 1990.
[18] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(**1**):9–11, 1990.

[19] Wojciech A. Trybulec. Binary operations on finite sequences. *Formalized Mathematics*, 1(**5**):979–981, 1990.
[20] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(**2**):291–296, 1990.
[21] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(**1**):67–71, 1990.
[22] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(**1**):73–83, 1990.
[23] Hiroshi Yamazaki, Yasunari Shidama, and Yatsuka Nakamura. Bessel's inequality. *Formalized Mathematics*, 11(**2**):169–173, 2003.

Received February 25, 2003