FORMALIZED MATHEMATICS
Volume 11, Number 3, 2003
University of Białystok

Some Properties for Convex Combinations

Noboru Endou
Gifu National College of Technology

Yasumasa Suzuki
Miyagi University

Yasunari Shidama
Shinshu University
Nagano

Summary. This is a continuation of [6]. In this article, we proved that convex combination on convex family is convex.

MML Identifier: CONVEX2.

The notation and terminology used in this paper are introduced in the following articles: [13], [18], [12], [8], [2], [19], [3], [5], [1], [10], [4], [17], [16], [15], [14], [11], [7], [6], and [9].

1. Convex Combinations on Convex Family

The following propositions are true:

(1) For every non empty RLS structure V and for all convex subsets M, N of V holds $M \cap N$ is convex.

(2) Let V be a real unitary space-like non empty unitary space structure, M be a subset of V, F be a finite sequence of elements of the carrier of V, and B be a finite sequence of elements of $\mathbb{R}$. Suppose $M = \{u;$ u ranges over vectors of V: $\bigwedge_{i\,:\,\text{natural number}}$ $(i \in \operatorname{dom} F \cap \operatorname{dom} B \Rightarrow \bigvee_{v\,:\,\text{vector of } V}$ $(v = F(i) \wedge (u|v) \leqslant B(i)))\}$. Then M is convex.

(3) Let V be a real unitary space-like non empty unitary space structure, M be a subset of V, F be a finite sequence of elements of the carrier of V, and B be a finite sequence of elements of $\mathbb{R}$. Suppose $M = \{u;$ u ranges over vectors of V: $\bigwedge_{i\,:\,\text{natural number}}$ $(i \in \operatorname{dom} F \cap \operatorname{dom} B \Rightarrow \bigvee_{v\,:\,\text{vector of } V}$ $(v = F(i) \wedge (u|v) < B(i)))\}$. Then M is convex.

© 2003 University of Białystok
ISSN 1426–2630

(4) Let V be a real unitary space-like non empty unitary space structure, M be a subset of V, F be a finite sequence of elements of the carrier of V, and B be a finite sequence of elements of $\mathbb{R}$. Suppose $M = \{u; u \text{ ranges over vectors of } V\colon \bigwedge_{i\colon \text{natural number}} (i \in \operatorname{dom} F \cap \operatorname{dom} B \Rightarrow \bigvee_{v\colon \text{vector of } V} (v = F(i) \wedge (u|v) \geqslant B(i)))\}$. Then M is convex.

(5) Let V be a real unitary space-like non empty unitary space structure, M be a subset of V, F be a finite sequence of elements of the carrier of V, and B be a finite sequence of elements of $\mathbb{R}$. Suppose $M = \{u; u \text{ ranges over vectors of } V\colon \bigwedge_{i\colon \text{natural number}} (i \in \operatorname{dom} F \cap \operatorname{dom} B \Rightarrow \bigvee_{v\colon \text{vector of } V} (v = F(i) \wedge (u|v) > B(i)))\}$. Then M is convex.

(6) Let V be a real linear space and M be a subset of V. Then for every subset N of V and for every linear combination L of N such that L is convex and $N \subseteq M$ holds $\sum L \in M$ if and only if M is convex.

Let V be a real linear space and let M be a subset of V. The functor LC_M yielding a set is defined as follows:

(Def. 1) For every set L holds $L \in \mathrm{LC}_M$ iff L is a linear combination of M.

Let V be a real linear space. Observe that there exists a linear combination of V which is convex.

Let V be a real linear space. A convex combination of V is a convex linear combination of V.

Let V be a real linear space and let M be a non empty subset of V. One can verify that there exists a linear combination of M which is convex.

Let V be a real linear space and let M be a non empty subset of V. A convex combination of M is a convex linear combination of M.

The following propositions are true:

(7) For every real linear space V and for every subset M of V holds Convex-Family $M \neq \emptyset$.

(8) For every real linear space V and for every subset M of V holds $M \subseteq \operatorname{conv} M$.

(9) Let V be a real linear space, L_1, L_2 be convex combinations of V, and r be a real number. If $0 < r$ and $r < 1$, then $r \cdot L_1 + (1 - r) \cdot L_2$ is a convex combination of V.

(10) Let V be a real linear space, M be a non empty subset of V, L_1, L_2 be convex combinations of M, and r be a real number. If $0 < r$ and $r < 1$, then $r \cdot L_1 + (1 - r) \cdot L_2$ is a convex combination of M.

(11) For every real linear space V holds there exists a linear combination of V which is convex.

(12) For every real linear space V and for every non empty subset M of V holds there exists a linear combination of M which is convex.

2. Miscellaneous

We now state several propositions:

(13) For every real linear space V and for all subspaces W_1, W_2 of V holds $\mathrm{Up}(W_1 + W_2) = \mathrm{Up}(W_1) + \mathrm{Up}(W_2)$.

(14) For every real linear space V and for all subspaces W_1, W_2 of V holds $\mathrm{Up}(W_1 \cap W_2) = \mathrm{Up}(W_1) \cap \mathrm{Up}(W_2)$.

(15) Let V be a real linear space, L_1, L_2 be convex combinations of V, and a, b be real numbers. Suppose $a \cdot b > 0$. Then the support of $a \cdot L_1 + b \cdot L_2 =$ (the support of $a \cdot L_1$) $\cup$ (the support of $b \cdot L_2$).

(16) Let F, G be functions. Suppose F and G are fiberwise equipotent. Let x_1, x_2 be sets. Suppose $x_1 \in \mathrm{dom}\, F$ and $x_2 \in \mathrm{dom}\, F$ and $x_1 \neq x_2$. Then there exist sets z_1, z_2 such that $z_1 \in \mathrm{dom}\, G$ and $z_2 \in \mathrm{dom}\, G$ and $z_1 \neq z_2$ and $F(x_1) = G(z_1)$ and $F(x_2) = G(z_2)$.

(17) Let V be a real linear space, L be a linear combination of V, and A be a subset of V. Suppose $A \subseteq$ the support of L. Then there exists a linear combination L_1 of V such that the support of $L_1 = A$.

References

[1] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(**2**):377–382, 1990.
[2] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(**1**):107–114, 1990.
[3] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(**1**):55–65, 1990.
[4] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(**1**):153–164, 1990.
[5] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(**1**):165–167, 1990.
[6] Noboru Endou, Takashi Mitsuishi, and Yasunari Shidama. Convex sets and convex combinations. *Formalized Mathematics*, 11(**1**):53–58, 2003.
[7] Noboru Endou, Takashi Mitsuishi, and Yasunari Shidama. Dimension of real unitary space. *Formalized Mathematics*, 11(**1**):23–28, 2003.
[8] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(**1**):35–40, 1990.
[9] Jarosław Kotowicz. Functions and finite sequences of real numbers. *Formalized Mathematics*, 3(**2**):275–278, 1992.
[10] Beata Padlewska. Families of sets. *Formalized Mathematics*, 1(**1**):147–152, 1990.
[11] Jan Popiołek. Introduction to Banach and Hilbert spaces - part I. *Formalized Mathematics*, 2(**4**):511–516, 1991.
[12] Andrzej Trybulec. Introduction to arithmetics. *To appear in Formalized Mathematics.*
[13] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(**1**):9–11, 1990.
[14] Wojciech A. Trybulec. Linear combinations in real linear space. *Formalized Mathematics*, 1(**3**):581–588, 1990.
[15] Wojciech A. Trybulec. Operations on subspaces in real linear space. *Formalized Mathematics*, 1(**2**):395–399, 1990.
[16] Wojciech A. Trybulec. Subspaces and cosets of subspaces in real linear space. *Formalized Mathematics*, 1(**2**):297–301, 1990.
[17] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(**2**):291–296, 1990.
[18] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(**1**):67–71, 1990.

[19] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(**1**):73–83, 1990.

Received April 3, 2003