FORMALIZED MATHEMATICS
Volume 11, Number 3, 2003
University of Białystok

The Class of Series-Parallel Graphs. Part II[1]

Krzysztof Retel
University of Białystok

Summary. In this paper we introduce two new operations on graphs: sum and union corresponding to parallel and series operation respectively. We determine N-free graph as the graph that does not embed Necklace 4. We define "fin_RelStr" as the set of all graphs with finite carriers. We also define the smallest class of graphs which contains the one-element graph and which is closed under parallel and series operations. The goal of the article is to prove the theorem that the class of finite series-parallel graphs is the class of finite N-free graphs. This paper formalizes the next part of [12].

MML Identifier: NECKLA_2.

The terminology and notation used in this paper are introduced in the following papers: [15], [14], [18], [7], [20], [8], [1], [2], [3], [13], [16], [4], [17], [19], [11], [5], [6], [9], and [10].

In this paper U denotes a universal class.

Next we state two propositions:

(1) For all sets X, Y such that $X \in U$ and $Y \in U$ and for every relation R between X and Y holds $R \in U$.

(2) The internal relation of Necklace 4 $= \{\langle 0, 1\rangle, \langle 1, 0\rangle, \langle 1, 2\rangle, \langle 2, 1\rangle, \langle 2, 3\rangle, \langle 3, 2\rangle\}$.

Let n be a natural number. One can check that every element of $\mathbf{R}_n$ is finite.

Next we state the proposition

(3) For every set x such that $x \in \mathbf{U}_0$ holds x is finite.

Let us mention that every element of $\mathbf{U}_0$ is finite.

Let us note that every number which is finite and ordinal is also natural.

[1] This work has been partially supported by CALCULEMUS grant HPRN-CT-2000-00102.

© 2003 University of Białystok
ISSN 1426–2630

Let G be a non empty relational structure. We say that G is N-free if and only if:

(Def. 1) G does not embed Necklace 4.

Let us mention that there exists a non empty relational structure which is N-free.

Let R, S be relational structures. The functor UnionOf(R, S) yielding a strict relational structure is defined by the conditions (Def. 2).

(Def. 2)(i) The carrier of UnionOf(R, S) = (the carrier of R) $\cup$ (the carrier of S), and

(ii) the internal relation of UnionOf(R, S) = (the internal relation of R) $\cup$ (the internal relation of S).

Let R, S be relational structures. The functor SumOf(R, S) yielding a strict relational structure is defined by the conditions (Def. 3).

(Def. 3)(i) The carrier of SumOf(R, S) = (the carrier of R) $\cup$ (the carrier of S), and

(ii) the internal relation of SumOf(R, S) = (the internal relation of R) $\cup$ (the internal relation of S) $\cup$ $[:$ the carrier of R, the carrier of $S :] \cup [:$ the carrier of S, the carrier of $R :]$.

The functor FinRelStr is defined by the condition (Def. 4).

(Def. 4) Let X be a set. Then $X \in$ FinRelStr if and only if there exists a strict relational structure R such that $X = R$ and the carrier of $R \in \mathbf{U}_0$.

Let us mention that FinRelStr is non empty.

The subset FinRelStrSp of FinRelStr is defined by the conditions (Def. 5).

(Def. 5)(i) For every strict relational structure R such that the carrier of R is non empty and trivial and the carrier of $R \in \mathbf{U}_0$ holds $R \in$ FinRelStrSp,

(ii) for all strict relational structures H_1, H_2 such that the carrier of H_1 misses the carrier of H_2 and $H_1 \in$ FinRelStrSp and $H_2 \in$ FinRelStrSp holds UnionOf$(H_1, H_2) \in$ FinRelStrSp and SumOf$(H_1, H_2) \in$ FinRelStrSp, and

(iii) for every subset M of FinRelStr such that for every strict relational structure R such that the carrier of R is non empty and trivial and the carrier of $R \in \mathbf{U}_0$ holds $R \in M$ and for all strict relational structures H_1, H_2 such that the carrier of H_1 misses the carrier of H_2 and $H_1 \in M$ and $H_2 \in M$ holds UnionOf$(H_1, H_2) \in M$ and SumOf$(H_1, H_2) \in M$ holds FinRelStrSp $\subseteq M$.

One can verify that FinRelStrSp is non empty.

We now state four propositions:

(4) For every set X such that $X \in$ FinRelStrSp holds X is a finite strict non empty relational structure.

(5) For every relational structure R such that $R \in$ FinRelStrSp holds the carrier of $R \in \mathbf{U}_0$.

(6) Let X be a set. Suppose $X \in$ FinRelStrSp . Then

(i) X is a strict non empty trivial relational structure, or

(ii) there exist strict relational structures H_1, H_2 such that the carrier of H_1 misses the carrier of H_2 and $H_1 \in$ FinRelStrSp and $H_2 \in$ FinRelStrSp and $X = \mathrm{UnionOf}(H_1, H_2)$ or $X = \mathrm{SumOf}(H_1, H_2)$.

(7) For every strict non empty relational structure R such that $R \in$ FinRelStrSp holds R is N-free.

References

[1] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(**2**):377–382, 1990.
[2] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(**1**):91–96, 1990.
[3] Grzegorz Bancerek. Sequences of ordinal numbers. *Formalized Mathematics*, 1(**2**):281–290, 1990.
[4] Józef Białas. Group and field definitions. *Formalized Mathematics*, 1(**3**):433–439, 1990.
[5] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(**1**):55–65, 1990.
[6] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(**1**):153–164, 1990.
[7] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(**1**):47–53, 1990.
[8] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(**1**):165–167, 1990.
[9] Bogdan Nowak and Grzegorz Bancerek. Universal classes. *Formalized Mathematics*, 1(**3**):595–600, 1990.
[10] Beata Padlewska. Families of sets. *Formalized Mathematics*, 1(**1**):147–152, 1990.
[11] Krzysztof Retel. The class of series – parallel graphs. Part I. *Formalized Mathematics*, 11(**1**):99–103, 2003.
[12] Stephan Thomasse. On better-quasi-ordering countable series-parallel orders. *Transactions of the American Mathematical Society*, 352(6):2491–2505, 2000.
[13] Andrzej Trybulec. Introduction to arithmetics. *To appear in Formalized Mathematics*.
[14] Andrzej Trybulec. Enumerated sets. *Formalized Mathematics*, 1(**1**):25–34, 1990.
[15] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(**1**):9–11, 1990.
[16] Wojciech A. Trybulec. Groups. *Formalized Mathematics*, 1(**5**):821–827, 1990.
[17] Wojciech A. Trybulec. Partially ordered sets. *Formalized Mathematics*, 1(**2**):313–319, 1990.
[18] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(**1**):67–71, 1990.
[19] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(**1**):73–83, 1990.
[20] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(**1**):181–186, 1990.

Received May 29, 2003