FORMALIZED MATHEMATICS
Volume 11, Number 3, 2003
University of Białystok

# On the Subcontinua of a Real Line[1]

Adam Grabowski
University of Białystok

**Summary.** In [10] we showed that the only proper subcontinua of the simple closed curve are arcs and single points. In this article we prove that the only proper subcontinua of the real line are closed intervals. We introduce some auxiliary notions such as $]a, b[_{\mathbb{Q}}$, $]a, b[_{\mathbb{IQ}}$ – intervals consisting of rational and irrational numbers respectively. We show also some basic topological properties of intervals.

MML Identifier: BORSUK_5.

The notation and terminology used in this paper are introduced in the following papers: [24], [27], [22], [23], [18], [25], [28], [3], [4], [26], [19], [6], [21], [13], [16], [17], [1], [8], [5], [9], [14], [7], [20], [15], [12], [11], and [2].

## 1. Preliminaries

The following three propositions are true:

(1) For all sets $A$, $B$, $C$, $D$ holds $(A \cup B \cup C) \cup D = A \cup (B \cup C \cup D)$.

(2) For all sets $A$, $B$, $a$ such that $A \subseteq B$ and $B \subseteq A \cup \{a\}$ holds $A \cup \{a\} = B$ or $A = B$.

(3) For all sets $x_1$, $x_2$, $x_3$, $x_4$, $x_5$, $x_6$ holds $\{x_1, x_2, x_3, x_4, x_5, x_6\} = \{x_1, x_3, x_6\} \cup \{x_2, x_4, x_5\}$.

In the sequel $x_1$, $x_2$, $x_3$, $x_4$, $x_5$, $x_6$, $x_7$ are sets.

Let $x_1$, $x_2$, $x_3$, $x_4$, $x_5$, $x_6$ be sets. We say that $x_1$, $x_2$, $x_3$, $x_4$, $x_5$, $x_6$ are mutually different if and only if the conditions (Def. 1) are satisfied.

(Def. 1) $x_1 \neq x_2$ and $x_1 \neq x_3$ and $x_1 \neq x_4$ and $x_1 \neq x_5$ and $x_1 \neq x_6$ and $x_2 \neq x_3$ and $x_2 \neq x_4$ and $x_2 \neq x_5$ and $x_2 \neq x_6$ and $x_3 \neq x_4$ and $x_3 \neq x_5$ and $x_3 \neq x_6$ and $x_4 \neq x_5$ and $x_4 \neq x_6$ and $x_5 \neq x_6$.

[1] This work has been partially supported by CALCULEMUS grant HPRN-CT-2000-00102.



© 2003 University of Białystok
ISSN 1426–2630

Let $x_1$, $x_2$, $x_3$, $x_4$, $x_5$, $x_6$, $x_7$ be sets. We say that $x_1$, $x_2$, $x_3$, $x_4$, $x_5$, $x_6$, $x_7$ are mutually different if and only if the conditions (Def. 2) are satisfied.

(Def. 2) $x_1 \neq x_2$ and $x_1 \neq x_3$ and $x_1 \neq x_4$ and $x_1 \neq x_5$ and $x_1 \neq x_6$ and $x_1 \neq x_7$ and $x_2 \neq x_3$ and $x_2 \neq x_4$ and $x_2 \neq x_5$ and $x_2 \neq x_6$ and $x_2 \neq x_7$ and $x_3 \neq x_4$ and $x_3 \neq x_5$ and $x_3 \neq x_6$ and $x_3 \neq x_7$ and $x_4 \neq x_5$ and $x_4 \neq x_6$ and $x_4 \neq x_7$ and $x_5 \neq x_6$ and $x_5 \neq x_7$ and $x_6 \neq x_7$.

One can prove the following propositions:

(4) For all sets $x_1$, $x_2$, $x_3$, $x_4$, $x_5$, $x_6$ such that $x_1$, $x_2$, $x_3$, $x_4$, $x_5$, $x_6$ are mutually different holds $\mathrm{card}\{x_1, x_2, x_3, x_4, x_5, x_6\} = 6$.

(5) For all sets $x_1$, $x_2$, $x_3$, $x_4$, $x_5$, $x_6$, $x_7$ such that $x_1$, $x_2$, $x_3$, $x_4$, $x_5$, $x_6$, $x_7$ are mutually different holds $\mathrm{card}\{x_1, x_2, x_3, x_4, x_5, x_6, x_7\} = 7$.

(6) If $\{x_1, x_2, x_3\}$ misses $\{x_4, x_5, x_6\}$, then $x_1 \neq x_4$ and $x_1 \neq x_5$ and $x_1 \neq x_6$ and $x_2 \neq x_4$ and $x_2 \neq x_5$ and $x_2 \neq x_6$ and $x_3 \neq x_4$ and $x_3 \neq x_5$ and $x_3 \neq x_6$.

(7) Suppose $x_1$, $x_2$, $x_3$ are mutually different and $x_4$, $x_5$, $x_6$ are mutually different and $\{x_1, x_2, x_3\}$ misses $\{x_4, x_5, x_6\}$. Then $x_1$, $x_2$, $x_3$, $x_4$, $x_5$, $x_6$ are mutually different.

(8) Suppose $x_1$, $x_2$, $x_3$, $x_4$, $x_5$, $x_6$ are mutually different and $\{x_1, x_2, x_3, x_4, x_5, x_6\}$ misses $\{x_7\}$. Then $x_1$, $x_2$, $x_3$, $x_4$, $x_5$, $x_6$, $x_7$ are mutually different.

(9) If $x_1$, $x_2$, $x_3$, $x_4$, $x_5$, $x_6$, $x_7$ are mutually different, then $x_7$, $x_1$, $x_2$, $x_3$, $x_4$, $x_5$, $x_6$ are mutually different.

(10) If $x_1$, $x_2$, $x_3$, $x_4$, $x_5$, $x_6$, $x_7$ are mutually different, then $x_1$, $x_2$, $x_5$, $x_3$, $x_6$, $x_7$, $x_4$ are mutually different.

(11) Let $T$ be a non empty topological space and $a$, $b$ be points of $T$. Given a map $f$ from $\mathbb{I}$ into $T$ such that $f$ is continuous and $f(0) = a$ and $f(1) = b$. Then there exists a map $g$ from $\mathbb{I}$ into $T$ such that $g$ is continuous and $g(0) = b$ and $g(1) = a$.

Let us observe that $\mathbb{R}^{\mathbf{1}}$ is arcwise connected.

Let us note that there exists a topological space which is connected and non empty.

## 2. Intervals

The following two propositions are true:

(12) Every subset of $\mathbb{R}$ is a subset of $\mathbb{R}^{\mathbf{1}}$.

(13) $\Omega_{\mathbb{R}^{\mathbf{1}}} = \mathbb{R}$.

Let $a$ be a real number. We introduce $]-\infty, a]$ as a synonym of $]-\infty, a]$. We introduce $]-\infty, a[$ as a synonym of $]-\infty, a[$. We introduce $[a, +\infty[$ as a synonym of $[a, +\infty[$. We introduce $]a, +\infty[$ as a synonym of $]a, +\infty[$.

Next we state a number of propositions:

(14) For all real numbers $a$, $b$ holds $a \in ]b, +\infty[$ iff $a > b$.

(15) For all real numbers $a$, $b$ holds $a \in [b, +\infty[$ iff $a \geqslant b$.

(16) For all real numbers $a$, $b$ holds $a \in ]-\infty, b]$ iff $a \leqslant b$.

(17) For all real numbers $a$, $b$ holds $a \in ]-\infty, b[$ iff $a < b$.

(18) For every real number $a$ holds $\mathbb{R} \setminus \{a\} = ]-\infty, a[ \cup ]a, +\infty[$.

(19) For all real numbers $a$, $b$, $c$, $d$ such that $a < b$ and $b \leqslant c$ holds $[a, b]$ misses $]c, d]$.

(20) For all real numbers $a$, $b$, $c$, $d$ such that $a < b$ and $b \leqslant c$ holds $[a, b[$ misses $[c, d]$.

(21) Let $A$, $B$ be subsets of the carrier of $\mathbb{R}^{\mathbf{1}}$ and $a$, $b$, $c$, $d$ be real numbers. Suppose $a < b$ and $b \leqslant c$ and $c < d$ and $A = [a, b[$ and $B = ]c, d]$. Then $A$ and $B$ are separated.

(22) For every real number $a$ holds $\mathbb{R} \setminus ]-\infty, a[ = [a, +\infty[$.

(23) For every real number $a$ holds $\mathbb{R} \setminus ]-\infty, a] = ]a, +\infty[$.

(24) For every real number $a$ holds $\mathbb{R} \setminus ]a, +\infty[ = ]-\infty, a]$.

(25) For every real number $a$ holds $\mathbb{R} \setminus [a, +\infty[ = ]-\infty, a[$.

(26) For every real number $a$ holds $]-\infty, a]$ misses $]a, +\infty[$.

(27) For every real number $a$ holds $]-\infty, a[$ misses $[a, +\infty[$.

(28) For all real numbers $a$, $b$, $c$ such that $a \leqslant c$ and $c \leqslant b$ holds $[a, b] \cup [c, +\infty[ = [a, +\infty[$.

(29) For all real numbers $a$, $b$, $c$ such that $a \leqslant c$ and $c \leqslant b$ holds $]-\infty, c] \cup [a, b] = ]-\infty, b]$.

(30) For every 1-sorted structure $T$ and for every subset $A$ of $T$ holds $\{A\}$ is a family of subsets of $T$.

(31) For every 1-sorted structure $T$ and for all subsets $A$, $B$ of $T$ holds $\{A, B\}$ is a family of subsets of $T$.

(32) For every 1-sorted structure $T$ and for all subsets $A$, $B$, $C$ of $T$ holds $\{A, B, C\}$ is a family of subsets of $T$.

Let us observe that every element of $\mathbb{Q}$ is real.

Let us observe that every element of the carrier of the metric space of real numbers is real.

Next we state four propositions:

(33) Let $A$ be a subset of the carrier of $\mathbb{R}^{\mathbf{1}}$ and $p$ be a point of the metric space of real numbers. Then $p \in \overline{A}$ if and only if for every real number $r$ such that $r > 0$ holds $\mathrm{Ball}(p, r)$ meets $A$.

(34) For all elements $p$, $q$ of the carrier of the metric space of real numbers such that $q \geqslant p$ holds $\rho(p, q) = q - p$.

(35) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ such that $A = \mathbb{Q}$ holds $\overline{A}$ = the carrier of $\mathbb{R}^{\mathbf{1}}$.

(36) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ and for all real numbers $a$, $b$ such that $A = ]a, b[$ and $a \neq b$ holds $\overline{A} = [a, b]$.

## 3. Rational and Irrational Numbers

Let us mention that $e$ is irrational.

The subset $\mathbb{IQ}$ of $\mathbb{R}$ is defined by:

(Def. 3) $\mathbb{IQ} = \mathbb{R} \setminus \mathbb{Q}$.

Let $a$, $b$ be real numbers. The functor $]a, b[_{\mathbb{Q}}$ yielding a subset of $\mathbb{R}$ is defined by:

(Def. 4) $]a, b[_{\mathbb{Q}} = \mathbb{Q} \cap ]a, b[$.

The functor $]a, b[_{\mathbb{IQ}}$ yielding a subset of $\mathbb{R}$ is defined as follows:

(Def. 5) $]a, b[_{\mathbb{IQ}} = \mathbb{IQ} \cap ]a, b[$.

One can prove the following proposition

(37) For every real number $x$ holds $x$ is irrational iff $x \in \mathbb{IQ}$.

Let us observe that there exists a real number which is irrational.

Let us note that $\mathbb{IQ}$ is non empty.

Next we state several propositions:

(38) For every rational number $a$ and for every irrational real number $b$ holds $a + b$ is irrational.

(39) For every irrational real number $a$ holds $-a$ is irrational.

(40) For every rational number $a$ and for every irrational real number $b$ holds $a - b$ is irrational.

(41) For every rational number $a$ and for every irrational real number $b$ holds $b - a$ is irrational.

(42) For every rational number $a$ and for every irrational real number $b$ such that $a \neq 0$ holds $a \cdot b$ is irrational.

(43) For every rational number $a$ and for every irrational real number $b$ such that $a \neq 0$ holds $\frac{b}{a}$ is irrational.

One can check that every real number which is irrational is also non zero.

The following propositions are true:

(44) For every rational number $a$ and for every irrational real number $b$ such that $a \neq 0$ holds $\frac{a}{b}$ is irrational.

(45) For every irrational real number $r$ holds $\operatorname{frac} r$ is irrational.

Let $r$ be an irrational real number. Note that $\operatorname{frac} r$ is irrational.

Let $a$ be an irrational real number. Note that $-a$ is irrational.

Let $a$ be a rational number and let $b$ be an irrational real number. One can verify the following observations:

* $a + b$ is irrational,
* $b + a$ is irrational,
* $a - b$ is irrational, and
* $b - a$ is irrational.

Let us observe that there exists a rational number which is non zero.

Let $a$ be a non zero rational number and let $b$ be an irrational real number. One can check the following observations:

* $a \cdot b$ is irrational,
* $b \cdot a$ is irrational,
* $\frac{a}{b}$ is irrational, and
* $\frac{b}{a}$ is irrational.

The following propositions are true:

(46) For every irrational real number $r$ holds $0 < \operatorname{frac} r$.

(47) For all real numbers $a$, $b$ such that $a < b$ there exist rational numbers $p_1$, $p_2$ such that $a < p_1$ and $p_1 < p_2$ and $p_2 < b$.

(48) For all real numbers $s_1$, $s_3$, $s_4$, $l$ such that $s_1 \leqslant s_3$ and $s_1 < s_4$ and $0 < l$ and $l < 1$ holds $s_1 < (1 - l) \cdot s_3 + l \cdot s_4$.

(49) For all real numbers $s_1$, $s_3$, $s_4$, $l$ such that $s_3 < s_1$ and $s_4 \leqslant s_1$ and $0 < l$ and $l < 1$ holds $(1 - l) \cdot s_3 + l \cdot s_4 < s_1$.

(50) For all real numbers $a$, $b$ such that $a < b$ there exists an irrational real number $p$ such that $a < p$ and $p < b$.

(51) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ such that $A = \mathbb{IQ}$ holds $\overline{A}$ = the carrier of $\mathbb{R}^{\mathbf{1}}$.

(52) For all real numbers $a$, $b$, $c$ such that $a < b$ holds $c \in ]a, b[_{\mathbb{Q}}$ iff $c$ is rational and $a < c$ and $c < b$.

(53) For all real numbers $a$, $b$, $c$ such that $a < b$ holds $c \in ]a, b[_{\mathbb{IQ}}$ iff $c$ is irrational and $a < c$ and $c < b$.

(54) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ and for all real numbers $a$, $b$ such that $a < b$ and $A = ]a, b[_{\mathbb{Q}}$ holds $\overline{A} = [a, b]$.

(55) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ and for all real numbers $a$, $b$ such that $a < b$ and $A = ]a, b[_{\mathbb{IQ}}$ holds $\overline{A} = [a, b]$.

(56) For every connected topological space $T$ and for every closed open subset $A$ of $T$ holds $A = \emptyset$ or $A = \Omega_T$.

(57) For every subset $A$ of $\mathbb{R}^{\mathbf{1}}$ such that $A$ is closed and open holds $A = \emptyset$ or $A = \mathbb{R}$.

## 4. Topological Properties of Intervals

We now state a number of propositions:

(58) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ and for all real numbers $a$, $b$ such that $A = [a, b[$ and $a \neq b$ holds $\overline{A} = [a, b]$.

(59) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ and for all real numbers $a$, $b$ such that $A = ]a, b]$ and $a \neq b$ holds $\overline{A} = [a, b]$.

(60) Let $A$ be a subset of the carrier of $\mathbb{R}^{\mathbf{1}}$ and $a$, $b$, $c$ be real numbers. If $A = [a, b[\cup]b, c]$ and $a < b$ and $b < c$, then $\overline{A} = [a, c]$.

(61) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ and for every real number $a$ such that $A = \{a\}$ holds $\overline{A} = \{a\}$.

(62) For every subset $A$ of $\mathbb{R}$ and for every subset $B$ of $\mathbb{R}^{\mathbf{1}}$ such that $A = B$ holds $A$ is open iff $B$ is open.

(63) For every subset $A$ of $\mathbb{R}^{\mathbf{1}}$ and for every real number $a$ such that $A = ]a, +\infty[$ holds $A$ is open.

(64) For every subset $A$ of $\mathbb{R}^{\mathbf{1}}$ and for every real number $a$ such that $A = ]-\infty, a[$ holds $A$ is open.

(65) For every subset $A$ of $\mathbb{R}^{\mathbf{1}}$ and for every real number $a$ such that $A = ]-\infty, a]$ holds $A$ is closed.

(66) For every subset $A$ of $\mathbb{R}^{\mathbf{1}}$ and for every real number $a$ such that $A = [a, +\infty[$ holds $A$ is closed.

(67) For every real number $a$ holds $[a, +\infty[ = \{a\}\cup]a, +\infty[$.

(68) For every real number $a$ holds $]-\infty, a] = \{a\}\cup]-\infty, a[$.

(69) For every real number $a$ holds $]a, +\infty[ \subseteq [a, +\infty[$.

(70) For every real number $a$ holds $]-\infty, a[ \subseteq ]-\infty, a]$.

Let $a$ be a real number. One can check the following observations:

* $]a, +\infty[$ is non empty,
* $]-\infty, a]$ is non empty,
* $]-\infty, a[$ is non empty, and
* $[a, +\infty[$ is non empty.

The following propositions are true:

(71) For every real number $a$ holds $]a, +\infty[ \neq \mathbb{R}$.

(72) For every real number $a$ holds $[a, +\infty[ \neq \mathbb{R}$.

(73) For every real number $a$ holds $]-\infty, a] \neq \mathbb{R}$.

(74) For every real number $a$ holds $]-\infty, a[ \neq \mathbb{R}$.

(75) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ and for every real number $a$ such that $A = ]a, +\infty[$ holds $\overline{A} = [a, +\infty[$.

(76) For every real number $a$ holds $\overline{]a, +\infty[} = [a, +\infty[$.

(77) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ and for every real number $a$ such that $A = ]-\infty, a[$ holds $\overline{A} = ]-\infty, a]$.

(78) For every real number $a$ holds $\overline{]-\infty, a[} = ]-\infty, a]$.

(79) Let $A$, $B$ be subsets of the carrier of $\mathbb{R}^{\mathbf{1}}$ and $b$ be a real number. If $A = ]-\infty, b[$ and $B = ]b, +\infty[$, then $A$ and $B$ are separated.

(80) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ and for all real numbers $a$, $b$ such that $a < b$ and $A = [a, b[ \cup ]b, +\infty[$ holds $\overline{A} = [a, +\infty[$.

(81) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ and for all real numbers $a$, $b$ such that $a < b$ and $A = ]a, b[ \cup ]b, +\infty[$ holds $\overline{A} = [a, +\infty[$.

(82) Let $A$ be a subset of the carrier of $\mathbb{R}^{\mathbf{1}}$ and $a$, $b$, $c$ be real numbers. If $a < b$ and $b < c$ and $A = ]a, b[_{\mathbb{Q}} \cup ]b, c[ \cup ]c, +\infty[$, then $\overline{A} = [a, +\infty[$.

(83) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ holds $-A = \mathbb{R} \setminus A$.

(84) For all real numbers $a$, $b$ such that $a < b$ holds $]a, b[_{\mathbb{IQ}}$ misses $]a, b[_{\mathbb{Q}}$.

(85) For all real numbers $a$, $b$ such that $a < b$ holds $\mathbb{R} \setminus ]a, b[_{\mathbb{Q}} = ]-\infty, a] \cup ]a, b[_{\mathbb{IQ}} \cup [b, +\infty[$.

(86) For all real numbers $a$, $b$, $c$ such that $a \leqslant b$ and $b < c$ holds $a \notin ]b, c[ \cup ]c, +\infty[$.

(87) For all real numbers $a$, $b$ such that $a < b$ holds $b \notin ]a, b[ \cup ]b, +\infty[$.

(88) For all real numbers $a$, $b$ such that $a < b$ holds $[a, +\infty[ \setminus (]a, b[ \cup ]b, +\infty[) = \{a\} \cup \{b\}$.

(89) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ such that $A = ]2, 3[_{\mathbb{Q}} \cup ]3, 4[ \cup ]4, +\infty[$ holds $-A = ]-\infty, 2] \cup ]2, 3[_{\mathbb{IQ}} \cup \{3\} \cup \{4\}$.

(90) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ and for every real number $a$ such that $A = \{a\}$ holds $-A = ]-\infty, a[ \cup ]a, +\infty[$.

(91) For all real numbers $a$, $b$ such that $a < b$ holds $]a, +\infty[ \cap ]-\infty, b] = ]a, b]$.

(92) $(]-\infty, 1[ \cup ]1, +\infty[) \cap (]-\infty, 2] \cup ]2, 3[_{\mathbb{IQ}} \cup \{3\} \cup \{4\}) = ]-\infty, 1[ \cup ]1, 2] \cup ]2, 3[_{\mathbb{IQ}} \cup \{3\} \cup \{4\}$.

(93) For all real numbers $a$, $b$ such that $a \leqslant b$ holds $]-\infty, b[ \setminus \{a\} = ]-\infty, a[ \cup ]a, b[$.

(94) For all real numbers $a$, $b$ such that $a \leqslant b$ holds $]a, +\infty[ \setminus \{b\} = ]a, b[ \cup ]b, +\infty[$.

(95) Let $A$ be a subset of the carrier of $\mathbb{R}^{\mathbf{1}}$ and $a$, $b$ be real numbers. If $a \leqslant b$ and $A = \{a\} \cup [b, +\infty[$, then $-A = ]-\infty, a[ \cup ]a, b[$.

(96) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ and for all real numbers $a$, $b$ such that $a < b$ and $A = ]-\infty, a[ \cup ]a, b[$ holds $\overline{A} = ]-\infty, b]$.

(97) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ and for all real numbers $a$, $b$ such that $a < b$ and $A = ]-\infty, a[ \cup ]a, b]$ holds $\overline{A} = ]-\infty, b]$.

(98) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ and for every real number $a$ such that $A = ]-\infty, a]$ holds $-A = ]a, +\infty[$.

(99) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ and for every real number $a$ such that $A = [a, +\infty[$ holds $-A = ]-\infty, a[$.

(100) Let $A$ be a subset of the carrier of $\mathbb{R}^{\mathbf{1}}$ and $a$, $b$, $c$ be real numbers. If $a < b$ and $b < c$ and $A = ]-\infty, a[ \cup ]a, b] \cup ]b, c[_{\mathbb{IQ}} \cup \{c\}$, then $\overline{A} = ]-\infty, c]$.

(101) Let $A$ be a subset of the carrier of $\mathbb{R}^{\mathbf{1}}$ and $a$, $b$, $c$, $d$ be real numbers. If $a < b$ and $b < c$ and $A = ]-\infty, a[ \cup ]a, b] \cup ]b, c[_{\mathbb{IQ}} \cup \{c\} \cup \{d\}$, then $\overline{A} = ]-\infty, c] \cup \{d\}$.

(102) Let $A$ be a subset of the carrier of $\mathbb{R}^{\mathbf{1}}$ and $a$, $b$ be real numbers. If $a \leqslant b$ and $A = ]-\infty, a] \cup \{b\}$, then $-A = ]a, b[ \cup ]b, +\infty[$.

(103) For all real numbers $a$, $b$ holds $[a, +\infty[ \cup \{b\} \neq \mathbb{R}$.

(104) For all real numbers $a$, $b$ holds $]-\infty, a] \cup \{b\} \neq \mathbb{R}$.

(105) For every topological structure $T_1$ and for all subsets $A$, $B$ of the carrier of $T_1$ such that $A \neq B$ holds $-A \neq -B$.

(106) For every subset $A$ of the carrier of $\mathbb{R}^{\mathbf{1}}$ such that $\mathbb{R} = -A$ holds $A = \emptyset$.

## 5. Subcontinua of a Real Line

Let us mention that $\mathbb{I}$ is arcwise connected.

We now state several propositions:

(107) Let $X$ be a compact subset of $\mathbb{R}^{\mathbf{1}}$ and $X'$ be a subset of $\mathbb{R}$. If $X' = X$, then $X'$ is upper bounded and lower bounded.

(108) Let $X$ be a compact subset of $\mathbb{R}^{\mathbf{1}}$, $X'$ be a subset of $\mathbb{R}$, and $x$ be a real number. If $x \in X'$ and $X' = X$, then $\inf X' \leqslant x$ and $x \leqslant \sup X'$.

(109) Let $C$ be a non empty compact connected subset of $\mathbb{R}^{\mathbf{1}}$ and $C'$ be a subset of $\mathbb{R}$. If $C = C'$ and $[\inf C', \sup C'] \subseteq C'$, then $[\inf C', \sup C'] = C'$.

(110) Let $A$ be a connected subset of $\mathbb{R}^{\mathbf{1}}$ and $a$, $b$, $c$ be real numbers. If $a \leqslant b$ and $b \leqslant c$ and $a \in A$ and $c \in A$, then $b \in A$.

(111) For every connected subset $A$ of $\mathbb{R}^{\mathbf{1}}$ and for all real numbers $a$, $b$ such that $a \in A$ and $b \in A$ holds $[a, b] \subseteq A$.

(112) Every non empty compact connected subset of $\mathbb{R}^{\mathbf{1}}$ is a non empty closed-interval subset of $\mathbb{R}$.

(113) For every non empty compact connected subset $A$ of $\mathbb{R}^{\mathbf{1}}$ there exist real numbers $a$, $b$ such that $a \leqslant b$ and $A = [a, b]$.

## 6. Sets with Proper Subsets Only

Let $T_1$ be a topological structure and let $F$ be a family of subsets of $T_1$. We say that $F$ has proper subsets if and only if:

(Def. 6) The carrier of $T_1 \notin F$.

One can prove the following proposition

(114) Let $T_1$ be a topological structure and $F$, $G$ be families of subsets of $T_1$ such that $F$ has proper subsets and $G \subseteq F$. Then $G$ has proper subsets.

Let $T_1$ be a non empty topological structure. Observe that there exists a family of subsets of $T_1$ which has proper subsets.

We now state the proposition

(115) Let $T_1$ be a non empty topological structure and $A$, $B$ be families of subsets of $T_1$ with proper subsets. Then $A \cup B$ has proper subsets.

Let $T$ be a topological structure and let $F$ be a family of subsets of $T$. We say that $F$ is open if and only if:

(Def. 7) For every subset $P$ of $T$ such that $P \in F$ holds $P$ is open.

We say that $F$ is closed if and only if:

(Def. 8) For every subset $P$ of $T$ such that $P \in F$ holds $P$ is closed.

Let $T$ be a topological space. Note that there exists a family of subsets of $T$ which is open, closed, and non empty.

## References


[1] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(**2**):377–382, 1990.
[2] Leszek Borys. Paracompact and metrizable spaces. *Formalized Mathematics*, 2(**4**):481–485, 1991.
[3] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(**1**):55–65, 1990.
[4] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(**1**):153–164, 1990.
[5] Czesław Byliński and Piotr Rudnicki. Bounding boxes for compact sets in $\mathcal{E}^2$. *Formalized Mathematics*, 6(**3**):427–440, 1997.
[6] Agata Darmochwał. Compact spaces. *Formalized Mathematics*, 1(**2**):383–386, 1990.
[7] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(**1**):165–167, 1990.
[8] Agata Darmochwał and Yatsuka Nakamura. Metric spaces as topological spaces - fundamental concepts. *Formalized Mathematics*, 2(**4**):605–608, 1991.
[9] Noboru Endou and Artur Korniłowicz. The definition of the Riemann definite integral and some related lemmas. *Formalized Mathematics*, 8(**1**):93–102, 1999.
[10] Adam Grabowski. On the decompositions of intervals and simple closed curves. *Formalized Mathematics*, 10(**3**):145–151, 2002.
[11] Stanisława Kanas, Adam Lecko, and Mariusz Startek. Metric spaces. *Formalized Mathematics*, 1(**3**):607–610, 1990.
[12] Andrzej Kondracki. Basic properties of rational numbers. *Formalized Mathematics*, 1(**5**):841–845, 1990.
[13] Jarosław Kotowicz. Convergent real sequences. Upper and lower bound of sets of real numbers. *Formalized Mathematics*, 1(**3**):477–481, 1990.
[14] Jarosław Kotowicz. The limit of a real function at infinity. *Formalized Mathematics*, 2(**1**):17–28, 1991.
[15] Wojciech Leończuk and Krzysztof Prażmowski. Incidence projective spaces. *Formalized Mathematics*, 2(**2**):225–232, 1991.
[16] Yatsuka Nakamura. Half open intervals in real numbers. *Formalized Mathematics*, 10(**1**):21–22, 2002.
[17] Beata Padlewska. Connected spaces. *Formalized Mathematics*, 1(**1**):239–244, 1990.
[18] Beata Padlewska. Families of sets. *Formalized Mathematics*, 1(**1**):147–152, 1990.
[19] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(**1**):223–230, 1990.

[20] Jan Popiołek. Some properties of functions modul and signum. *Formalized Mathematics*, 1(**2**):263–264, 1990.
[21] Konrad Raczkowski and Paweł Sadowski. Topological properties of subsets in real numbers. *Formalized Mathematics*, 1(**4**):777–780, 1990.
[22] Andrzej Trybulec. Introduction to arithmetics. *To appear in Formalized Mathematics*.
[23] Andrzej Trybulec. Enumerated sets. *Formalized Mathematics*, 1(**1**):25–34, 1990.
[24] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(**1**):9–11, 1990.
[25] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers. *Formalized Mathematics*, 1(**3**):445–449, 1990.
[26] Michał J. Trybulec. Integers. *Formalized Mathematics*, 1(**3**):501–505, 1990.
[27] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(**1**):67–71, 1990.
[28] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(**1**):73–83, 1990.

*Received June 12, 2003*