

The Friendship Theorem¹

Karol Pąk
Institute of Informatics
University of Białystok
Poland

Summary. In this article we prove the friendship theorem according to the article [1], which states that if a group of people has the property that any pair of persons have exactly one common friend, then there is a universal friend, i.e. a person who is a friend of every other person in the group.

MML identifier: FRIENDS1, version: 7.15.01 4.184.1155

The papers [3], [2], [6], [7], [11], [8], [9], [15], [14], [4], [13], [5], [17], [18], [12], [16], and [10] provide the terminology and notation for this paper.

1. PRELIMINARIES

For simplicity, we adopt the following rules: x, y, z are sets, i, k, n are natural numbers, R is a binary relation, P is a finite binary relation, and p, q are finite sequences.

Let us consider P, x . Observe that $P^\circ x$ is finite.

We now state several propositions:

- (1) $\overline{\overline{R}} = \overline{\overline{R^\smile}}$.
- (2) If R is symmetric, then $R^\circ x = R^{-1}(x)$.
- (3) If $(p|_k) \frown (p|k) = (q|_n) \frown (q|n)$ and $k \leq n \leq \text{len } p$, then $p = (q|_{n-k}) \frown (q|(n-k))$.
- (4) If $n \in \text{dom } q$ and $p = (q|_n) \frown (q|n)$, then $q = (p|_{\text{len } p - n}) \frown (p|(\text{len } p - n))$.

¹This work has been supported by the Polish Ministry of Science and Higher Education project “Managing a Large Repository of Computer-verified Mathematical Knowledge” (N N519 385136).

- (5) If $(p \upharpoonright k) \cap (p \upharpoonright k) = (q \upharpoonright n) \cap (q \upharpoonright n)$, then there exists i such that $p = (q \upharpoonright i) \cap (q \upharpoonright i)$.

The scheme *Sch* deals with a non empty set \mathcal{A} , a non zero natural number \mathcal{B} , and a unary predicate \mathcal{P} , and states that:

There exists a cardinal number C such that $\mathcal{B} \cdot C = \{F \in \mathcal{A}^{\mathcal{B}}: \mathcal{P}[F]\}$

provided the following requirements are met:

- For all finite sequences p, q of elements of \mathcal{A} such that $p \cap q$ is \mathcal{B} -element and $\mathcal{P}[p \cap q]$ holds $\mathcal{P}[q \cap p]$, and
- For every element p of $\mathcal{A}^{\mathcal{B}}$ such that $\mathcal{P}[p]$ and for every natural number i such that $i < \mathcal{B}$ and $p = (p \upharpoonright i) \cap (p \upharpoonright i)$ holds $i = 0$.

One can prove the following propositions:

- (6) Let X be a non empty set, A be a non empty finite subset of X , and P be a function from X into 2^X . Suppose that for every x such that $x \in X$ holds $\overline{P(x)} = n$. Then $\overline{\{F \in X^{k+1}: F(1) \in A \wedge \bigwedge_i (i \in \text{Seg } k \Rightarrow F(i+1) \in P(F(i)))\}} = \overline{A} \cdot n^k$.
- (7) If $\text{len } p$ is prime and there exists i such that $0 < i < \text{len } p$ and $p = (p \upharpoonright i) \cap (p \upharpoonright i)$, then $\text{rng } p \subseteq \{p(1)\}$.

2. THE FRIENDSHIP GRAPH

Let us consider R and let x be an element of field R . We say that x is universal friend if and only if:

- (Def. 1) For every y such that $y \in \text{field } R \setminus \{x\}$ holds $\langle x, y \rangle \in R$.

Let R be a binary relation. We say that R has universal friend if and only if:

- (Def. 2) There exists an element of field R which is universal friend.

Let R be a binary relation. We introduce R is without universal friend as an antonym of R has universal friend.

Let R be a binary relation. We say that R is friendship graph like if and only if:

- (Def. 3) For all x, y such that $x, y \in \text{field } R$ and $x \neq y$ there exists z such that $R^\circ x \cap \text{Coim}(R, y) = \{z\}$.

Let us observe that there exists a binary relation which is finite, symmetric, irreflexive, and friendship graph like.

A friendship graph is a finite symmetric irreflexive friendship graph like binary relation.

In the sequel F_1 is a friendship graph.

The following propositions are true:

- (8) $2 \mid \overline{F_1^\circ x}$.
- (9) If $x, y \in \text{field } F_1$ and $\langle x, y \rangle \notin F_1$, then $\overline{F_1^\circ x} = \overline{F_1^\circ y}$.
- (10) If F_1 is without universal friend and $x \in \text{field } F_1$, then $\overline{F_1^\circ x} > 2$.
- (11) If F_1 is without universal friend and $x, y \in \text{field } F_1$, then $\overline{F_1^\circ x} = \overline{F_1^\circ y}$.
- (12) If F_1 is without universal friend and $x \in \text{field } F_1$, then $\overline{\text{field } F_1} = 1 + \overline{F_1^\circ x} \cdot (\overline{F_1^\circ x} - 1)$.
- (13) For all elements x, y of field F_1 such that x is universal friend and $x \neq y$ there exists z such that $F_1^\circ y = \{x, z\}$ and $F_1^\circ z = \{x, y\}$.

3. THE FRIENDSHIP THEOREM

Next we state the proposition

- (14) If F_1 is non empty, then F_1 has universal friend.

REFERENCES

- [1] Michael Albert. Notes on the friendship theorem, <http://www.math.auckland.ac.nz/~olympiad/training/2006/friendship.pdf>.
- [2] Grzegorz Bancerek. Cardinal arithmetics. *Formalized Mathematics*, 1(3):543–547, 1990.
- [3] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(2):377–382, 1990.
- [4] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [5] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [6] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [7] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(3):529–536, 1990.
- [8] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [9] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [10] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [11] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(1):165–167, 1990.
- [12] Jarosław Kotowicz. Functions and finite sequences of real numbers. *Formalized Mathematics*, 3(2):275–278, 1992.
- [13] Rafał Kwiatek. Factorial and Newton coefficients. *Formalized Mathematics*, 1(5):887–890, 1990.
- [14] Rafał Kwiatek and Grzegorz Zwara. The divisibility of integers and integer relative primes. *Formalized Mathematics*, 1(5):829–832, 1990.
- [15] Michał J. Trybulec. Integers. *Formalized Mathematics*, 1(3):501–505, 1990.
- [16] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [17] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [18] Edmund Woronowicz and Anna Zalewska. Properties of binary relations. *Formalized Mathematics*, 1(1):85–89, 1990.

Received May 15, 2012