

Brouwer Invariance of Domain Theorem¹

Karol Pałk
Institute of Informatics
University of Białystok
Sosnowa 64, 15-887 Białystok
Poland

Summary. In this article we focus on a special case of the Brouwer invariance of domain theorem. Let us A, B be a subsets of \mathcal{E}^n , and $f : A \rightarrow B$ be a homeomorphic. We prove that, if A is closed then f transform the boundary of A to the boundary of B ; and if B is closed then f transform the interior of A to the interior of B . These two cases are sufficient to prove the topological invariance of dimension, which is used to prove basic properties of the n -dimensional manifolds, and also to prove basic properties of the boundary and the interior of manifolds, e.g. the boundary of an n -dimension manifold with boundary is an $(n - 1)$ -dimension manifold. This article is based on [18]; [21] and [20] can also serve as reference books.

MSC: 54A05 03B35

Keywords: continuous transformations; topological dimension

MML identifier: BROUWER3, version: 8.1.02 5.22.1199

The notation and terminology used in this paper have been introduced in the following articles: [27], [1], [14], [4], [6], [15], [37], [7], [8], [40], [31], [34], [38], [2], [3], [9], [5], [33], [13], [44], [45], [10], [42], [43], [35], [17], [28], [29], [25], [46], [16], [47], [26], [30], [32], and [12].

1. PRELIMINARIES

From now on x, X denote sets, n, m, i denote natural numbers, p, q denote points of \mathcal{E}_T^n , A, B denote subsets of \mathcal{E}_T^n , and r, s denote real numbers.

Let us consider X and n . One can verify that every function from X into \mathcal{E}_T^n is finite sequence-yielding.

¹The paper has been financed by the resources of the Polish National Science Centre granted by decision no DEC-2012/07/N/ST6/02147.

Let us consider m . Let f be a function from X into \mathcal{E}_T^n and g be a function from X into \mathcal{E}_T^m . Let us observe that the functor $f \frown g$ yields a function from X into \mathcal{E}_T^{n+m} . Let T be a topological space. Let f be a continuous function from T into \mathcal{E}_T^n and g be a continuous function from T into \mathcal{E}_T^m . Note that $f \frown g$ is continuous as a function from T into \mathcal{E}_T^{n+m} .

Let f be a real-valued function. The functor $[[f]]$ yielding a function is defined by

(Def. 1) (i) $\text{dom } it = \text{dom } f$, and

(ii) for every object x such that $x \in \text{dom } it$ holds $it(x) = [[f(x)]]$.

One can verify that $[[f]]$ is (the carrier of \mathcal{E}_T^1)-valued.

Let us consider X . Let Y be a non empty real-membered set and f be a function from X into Y . One can verify that the functor $[[f]]$ yields a function from X into \mathcal{E}_T^1 . Let T be a non empty topological space and f be a continuous function from T into \mathbb{R}^1 . Note that $[[f]]$ is continuous as a function from T into \mathcal{E}_T^1 .

Let f be a continuous real map of T . Observe that $[[f]]$ is continuous as a function from T into \mathcal{E}_T^1 .

2. A DISTRIBUTION OF SPHERE

In the sequel N denotes a non zero natural number and u, t denote points of \mathcal{E}_T^{N+1} .

Now we state the propositions:

(1) Let us consider an element F of $((\text{the carrier of } \mathbb{R}^1)^\alpha)^N$. Suppose If $i \in \text{dom } F$, then $F(i) = \text{PROJ}(N+1, i)$. Then

(i) for every t , $(\prod^* F)(t) = t \upharpoonright N$, and

(ii) for every subsets S_3, S_2 of \mathcal{E}_T^{N+1} such that $S_3 = \{u : u(N+1) \geq 0 \text{ and } |u| = 1\}$ and $S_2 = \{t : t(N+1) \leq 0 \text{ and } |t| = 1\}$ holds $(\prod^* F)^\circ S_3 = \overline{\text{Ball}}(0_{\mathcal{E}_T^N}, 1)$ and $(\prod^* F)^\circ S_2 = \overline{\text{Ball}}(0_{\mathcal{E}_T^N}, 1)$ and

$(\prod^* F)^\circ (S_3 \cap S_2) = \text{Sphere}(0_{\mathcal{E}_T^N}, 1)$ and for every function H from $\mathcal{E}_T^{N+1} \upharpoonright S_3$ into $\text{Tdisk}(0_{\mathcal{E}_T^N}, 1)$ such that $H = \prod^* F \upharpoonright S_3$ holds H is a homeomorphism and for every function H from $\mathcal{E}_T^{N+1} \upharpoonright S_2$ into

$\text{Tdisk}(0_{\mathcal{E}_T^N}, 1)$ such that $H = \prod^* F \upharpoonright S_2$ holds H is a homeomorphism,

where α is the carrier of \mathcal{E}_T^{N+1} . PROOF: Set $N_2 = N+1$. Set $T_{10} = \mathcal{E}_T^{N_2}$. Set $T_4 = \mathcal{E}_T^N$. Set $N_3 = N \text{ NormF}$. Set $N_4 = N_3 \cdot N_3$. Reconsider $O = 1$ as an element of \mathbb{N} . Set $T_3 = \text{Tdisk}(0_{\mathcal{E}_T^N}, 1)$. Reconsider $m_2 = -N_4$ as a function from T_4 into \mathbb{R}^1 . Reconsider $m_1 = 1 + m_2$ as a function from T_4 into \mathbb{R}^1 . Set $F_1 = \prod^* F$. For every t , $(\prod^* F)(t) = t \upharpoonright N$ by [2, (13)], [41, (25)], [4, (1)]. $\overline{\text{Ball}}(0_{T_4}, 1) \subseteq F_1^\circ S_3$ by [14, (22)], [28, (11)], [6, (16)],

[11, (145)]. $\overline{\text{Ball}}(0_{T_4}, 1) \subseteq F_1^\circ S_2$ by [14, (22)], [28, (11)], [6, (16)], [11, (145)]. $\text{Sphere}(0_{T_4}, 1) \subseteq F_1^\circ(S_2 \cap S_3)$ by [14, (22)], [28, (12)], [6, (16), (92)]. $F_1^\circ S_3 \subseteq \overline{\text{Ball}}(0_{T_4}, 1)$ by [14, (22)], [4, (59)], [24, (17)], [19, (10)]. $F_1^\circ S_2 \subseteq \overline{\text{Ball}}(0_{T_4}, 1)$ by [14, (22)], [4, (59)], [24, (17)], [19, (10)]. $F_1^\circ(S_2 \cap S_3) \subseteq \text{Sphere}(0_{T_4}, 1)$ by [14, (22)], [4, (59)], [24, (17)], [19, (10)]. For every function H from $\mathcal{E}_T^{N+1} \upharpoonright S_3$ into $\text{Tdisk}(0_{\mathcal{E}_T^N}, 1)$ such that $H = \prod^* F \upharpoonright S_3$ holds H is a homeomorphism by [24, (17)], [17, (17)], [2, (11)], [25, (13)]. For every objects x_1, x_2 such that $x_1, x_2 \in \text{dom } H$ and $H(x_1) = H(x_2)$ holds $x_1 = x_2$ by [14, (22)], [19, (10)], [7, (47)], [39, (40)]. Set $T_3 = \text{Tdisk}(0_{T_4}, 1)$. Set $M = m_1 \upharpoonright T_3$. Reconsider $M_1 = M$ as a continuous function from T_3 into \mathbb{R} . Reconsider $M_2 = -\sqrt{M_1}$ as a function from T_3 into \mathbb{R} . For every point p of T_4 such that $p \in$ the carrier of T_3 holds $M_1(p) = 1 - |p| \cdot |p|$ by [7, (49)]. Reconsider $S_1 = |[M_2]|$ as a continuous function from T_3 into \mathcal{E}_T^1 . Reconsider $I_3 = \text{id}_{T_3}$ as a continuous function from T_3 into T_4 . Reconsider $I_4 = I_3 \frown S_1$ as a continuous function from T_3 into \mathcal{E}_T^{N+O} . For every objects $y, x, z \in \text{rng } H$ and $x = I_4(y)$ iff $x \in \text{dom } H$ and $y = H(x)$ by [7, (17)], [11, (145), (144), (55)]. For every subset P of $T_{10} \upharpoonright S_2$, P is open iff $H^\circ P$ is open by [4, (1)], [2, (13)], [25, (57)]. \square

- (2) Let us consider subsets S_3, S_2 of \mathcal{E}_T^n . Suppose
- (i) $S_3 = \{s, \text{ where } s \text{ is a point of } \mathcal{E}_T^n : s(n) \geq 0 \text{ and } |s| = 1\}$, and
 - (ii) $S_2 = \{t, \text{ where } t \text{ is a point of } \mathcal{E}_T^n : t(n) \leq 0 \text{ and } |t| = 1\}$.

Then

- (iii) S_3 is closed, and
 - (iv) S_2 is closed.
- (3) Let us consider a metrizable topological space T_2 . Suppose T_2 is finite-ind and second-countable. Let us consider a closed subset F of T_2 . Suppose $\text{ind } F^c \leq n$. Let us consider a continuous function f from $T_2 \upharpoonright F$ into $\text{TopUnitCircle}(n+1)$. Then there exists a continuous function g from T_2 into $\text{TopUnitCircle}(n+1)$ such that $g \upharpoonright F = f$. PROOF: Define $\mathcal{P}[\text{natural number}] \equiv$ for every metrizable topological space T_2 such that T_2 is finite-ind and second-countable for every closed subset F of T_2 such that $\text{ind } F^c \leq \$_1$ for every continuous function f from $T_2 \upharpoonright F$ into $\text{TopUnitCircle}(\$_1+1)$, there exists a function g from T_2 into $\text{TopUnitCircle}(\$_1+1)$ such that g is continuous and $g \upharpoonright F = f$. For every n such that $\mathcal{P}[n]$ holds $\mathcal{P}[n+1]$ by (2), [29, (9)], [42, (13)], [44, (121)]. $\mathcal{P}[(0 \text{ qua natural number})]$ by [44, (143), (135)], [29, (9)], [14, (70)]. For every n , $\mathcal{P}[n]$ from [2, Sch. 2]. \square
- (4) Suppose $p \notin A$ and $r > 0$. Then there exists a function h from $\mathcal{E}_T^n \upharpoonright A$ into $\mathcal{E}_T^n \upharpoonright \text{Sphere}(p, r)$ such that
- (i) h is continuous, and

(ii) $h \upharpoonright \text{Sphere}(p, r) = \text{id}_{A \cap \text{Sphere}(p, r)}$.

(5) If $r + |p - q| \leq s$, then $\text{Ball}(p, r) \subseteq \text{Ball}(q, s)$.

(6) If A is not boundary, then $\text{ind } A = n$.

Now we state the proposition:

(7) THE SMALL INDUCTIVE DIMENSION OF THE SPHERE:

If $r > 0$, then $\text{ind } \text{Sphere}(p, r) = n - 1$. PROOF: If $\text{ind } A \leq i$ and $\text{ind } B \leq i$ and A is closed, then $\text{ind}(A \cup B) \leq i$ by [33, (31)], [23, (93)], [35, (22)], [36, (5)]. \square

3. A CHARACTERIZATION OF OPEN SETS IN EUCLIDEAN SPACE IN TERMS OF CONTINUOUS TRANSFORMATIONS

Now we state the propositions:

(8) Suppose $n > 0$ and $p \in A$ and for every r such that $r > 0$ there exists an open subset U of $\mathcal{E}_T^n \upharpoonright A$ such that $p \in U$ and $U \subseteq \text{Ball}(p, r)$ and for every function f from $\mathcal{E}_T^n \upharpoonright (A \setminus U)$ into $\text{TopUnitCircle } n$ such that f is continuous there exists a function h from $\mathcal{E}_T^n \upharpoonright A$ into $\text{TopUnitCircle } n$ such that h is continuous and $h \upharpoonright (A \setminus U) = f$. Then $p \in \text{Fr } A$. PROOF: Set $T_7 = \mathcal{E}_T^n$. Set $c_1 =$ the carrier of T_7 . Set $S = \text{Sphere}(0_{T_7}, 1)$. Set $T_9 = \text{TopUnitCircle } n$. Reconsider $c = c_1 \setminus \{0_{T_7}\}$ as a non empty open subset of T_7 . Set $n_3 = n \text{ NormF}$. Set $T_8 = T_7 \upharpoonright c$. Set $G = \text{transl}(p, T_7)$. Reconsider $I = \xrightarrow{T_8}$ as a continuous function from T_8 into T_7 . $0 \notin \text{rng}(n_3 \upharpoonright T_8)$ by [44, (57)], [14, (22)], [7, (47)], [14, (8), (70)]. Reconsider $n_2 = n_3 \upharpoonright T_8$ as a non-empty continuous function from T_8 into \mathbb{R}^1 . Reconsider $b = I/n_2$ as a function from T_8 into T_7 . Set $E_1 = \mathcal{E}^n$. Set $T_2 = E_{1\text{top}}$. Reconsider $e = p$ as a point of E_1 . Reconsider $I_1 = \text{Int } A$ as a subset of T_2 . Consider r being a real number such that $r > 0$ and $\text{Ball}(e, r) \subseteq I_1$. Set $r_2 = \frac{r}{2}$. Consider U being an open subset of $T_7 \upharpoonright A$ such that $p \in U$ and $U \subseteq \text{Ball}(p, r_2)$ and for every function f from $T_7 \upharpoonright (A \setminus U)$ into T_9 such that f is continuous there exists a function h from $T_7 \upharpoonright A$ into T_9 such that h is continuous and $h \upharpoonright (A \setminus U) = f$. Reconsider $S_4 = \text{Sphere}(p, r_2)$ as a non empty subset of T_7 . Consider a being an object such that $a \in S_4$. Reconsider $C_2 = \overline{\text{Ball}}(p, r_2)$ as a non empty subset of T_7 . Reconsider $s_2 = S_4$ as a non empty subset of $T_7 \upharpoonright C_2$. Reconsider $A_1 = A \setminus U$ as a non empty subset of T_7 . Set $T_1 = T_7 \upharpoonright A_1$. Set $t = \text{transl}(-p, T_7)$. Set $T = t \upharpoonright T_1$. $\text{rng } T \subseteq c$ by [7, (47)], [42, (21)]. Reconsider $T_1 1 = T$ as a continuous function from T_1 into T_8 . For every point p of T_7 such that $p \in c$ holds $b(p) = \frac{1}{|p|} \cdot p$ and $|\frac{1}{|p|} \cdot p| = 1$ by [22, (84)], [7, (49)], [26, (72)], [12, (56)]. $\text{rng } b \subseteq S$ by [42, (13)]. Reconsider $B = b$ as a function from T_8 into T_9 . Set $m = r_2 \bullet T_7$. Set $M = m \upharpoonright T_9$. Reconsider $M = m \upharpoonright T_9$ as a continuous function from T_9 into T_7 . Reconsider $c_2 = C_2$ as a subset of $T_7 \upharpoonright A$. Consider h being a function from $T_7 \upharpoonright A$ into T_9 such

that h is continuous and $h \upharpoonright (A \setminus U) = B \cdot T_1 1$. Reconsider $G_2 = G \cdot (M \cdot h)$ as a continuous function from $T_7 \upharpoonright A$ into T_7 . $\text{rng } G_2 \subseteq S_4$ by [7, (12), (11), (47)], [42, (28), (15)]. Reconsider $g_2 = G_2$ as a function from $T_7 \upharpoonright A$ into $T_7 \upharpoonright S_4$. Reconsider $g_1 = g_2 \upharpoonright ((T_7 \upharpoonright A) \upharpoonright c_2)$ as a continuous function from $T_7 \upharpoonright C_2$ into $(T_7 \upharpoonright C_2) \upharpoonright s_2$. For every point w of $T_7 \upharpoonright C_2$ such that $w \in S_4$ holds $g_1(w) = w$ by [7, (11), (12)], [44, (61)], [7, (47)]. \square

(9) Suppose $p \in \text{Fr } A$ and A is closed. Suppose $r > 0$. Then there exists an open subset U of $\mathcal{E}_T^n \upharpoonright A$ such that

- (i) $p \in U$, and
- (ii) $U \subseteq \text{Ball}(p, r)$, and
- (iii) for every function f from $\mathcal{E}_T^n \upharpoonright (A \setminus U)$ into $\text{TopUnitCircle } n$ such that f is continuous there exists a function h from $\mathcal{E}_T^n \upharpoonright A$ into $\text{TopUnitCircle } n$ such that h is continuous and $h \upharpoonright (A \setminus U) = f$.

PROOF: $n > 0$ by [14, (77), (22)], [12, (33)]. Set $r_3 = \frac{r}{3}$. Set $r_2 = 2 \cdot r_3$. Set $B = \text{Ball}(p, r_3)$. Consider x being an object such that $x \in A^c$ and $x \in B$. Set $u = \text{Ball}(x, r_2)$. $u \subseteq \text{Ball}(p, r)$. \square

4. BROUWER INVARIANCE OF DOMAIN THEOREM – SPECIAL CASE

Let us consider a function h from $\mathcal{E}_T^n \upharpoonright A$ into $\mathcal{E}_T^n \upharpoonright B$. Now we state the propositions:

- (10) If A is closed and $p \in \text{Fr } A$, then if h is a homeomorphism, then $h(p) \in \text{Fr } B$. The theorem is a consequence of (9) and (8).
- (11) If B is closed and $p \in \text{Int } A$, then if h is a homeomorphism, then $h(p) \in \text{Int } B$. The theorem is a consequence of (8) and (9).
- (12) Suppose A is closed and B is closed. Then if h is a homeomorphism, then $h^\circ(\text{Int } A) = \text{Int } B$ and $h^\circ(\text{Fr } A) = \text{Fr } B$. PROOF: $h^\circ(\text{Int } A) = \text{Int } B$ by (11), (10), [46, (39)]. \square

5. TOPOLOGICAL INVARIANCE OF DIMENSION – AN INTRODUCTION TO MANIFOLDS

Now we state the proposition:

- (13) Suppose $r > 0$. Let us consider a subset U of $\text{Tdisk}(p, r)$. Suppose U is open and non empty. Let us consider a subset A of \mathcal{E}_T^n . If $A = U$, then $\text{Int } A$ is not empty.

Let us consider a non empty topological space T , subsets A, B of T , r, s , a point p_1 of \mathcal{E}_T^n , and a point p_2 of \mathcal{E}_T^m .

Let us assume that $r > 0$ and $s > 0$. Now we state the propositions:

- (14) Suppose $T \upharpoonright A$ and $\text{Tdisk}(p_1, r)$ are homeomorphic and $T \upharpoonright B$ and $\text{Tdisk}(p_2, s)$ are homeomorphic and $\text{Int } A$ meets $\text{Int } B$. Then $n = m$. The theorem is a consequence of (13) and (6).
- (15) Suppose $T \upharpoonright A$ and $\mathcal{E}_T^n \upharpoonright \text{Ball}(p_1, r)$ are homeomorphic and $T \upharpoonright B$ and $\text{Tdisk}(p_2, s)$ are homeomorphic and $\text{Int } A$ meets $\text{Int } B$. Then $n = m$. The theorem is a consequence of (13) and (6).

Now we state the propositions:

- (16) (i) $(\text{transl}(p, \mathcal{E}_T^n))^\circ(\text{Ball}(q, r)) = \text{Ball}(q + p, r)$, and
(ii) $(\text{transl}(p, \mathcal{E}_T^n))^\circ(\overline{\text{Ball}}(q, r)) = \overline{\text{Ball}}(q + p, r)$, and
(iii) $(\text{transl}(p, \mathcal{E}_T^n))^\circ(\text{Sphere}(q, r)) = \text{Sphere}((q + p), r)$.
- PROOF: Set $T_5 = \mathcal{E}_T^n$. Set $T = \text{transl}(p, T_5)$. $T^\circ(\text{Ball}(q, r)) = \text{Ball}(q + p, r)$ by [28, (7)], [42, (27)]. $T^\circ(\overline{\text{Ball}}(q, r)) = \overline{\text{Ball}}(q + p, r)$ by [28, (8)], [42, (27)]. $T^\circ(\text{Sphere}(q, r)) \subseteq \text{Sphere}((q + p), r)$ by [28, (9)]. \square
- (17) Suppose $s > 0$. Then
(i) $(s \bullet \mathcal{E}_T^n)^\circ(\text{Ball}(p, r)) = \text{Ball}(s \cdot p, r \cdot s)$, and
(ii) $(s \bullet \mathcal{E}_T^n)^\circ(\overline{\text{Ball}}(p, r)) = \overline{\text{Ball}}(s \cdot p, r \cdot s)$, and
(iii) $(s \bullet \mathcal{E}_T^n)^\circ(\text{Sphere}(p, r)) = \text{Sphere}((s \cdot p), (r \cdot s))$.

PROOF: Set $T_5 = \mathcal{E}_T^n$. Set $M = s \bullet T_5$. $M^\circ(\text{Ball}(p, r)) = \text{Ball}(s \cdot p, r \cdot s)$ by [42, (34)], [14, (11)], [28, (7)]. $M^\circ(\overline{\text{Ball}}(p, r)) = \overline{\text{Ball}}(s \cdot p, r \cdot s)$ by [42, (34)], [14, (11)], [28, (8)]. $M^\circ(\text{Sphere}(p, r)) \subseteq \text{Sphere}((s \cdot p), (r \cdot s))$ by [42, (34)], [14, (11)], [28, (9)]. \square

- (18) Let us consider a rotation homogeneous additive function f from \mathcal{E}_T^n into \mathcal{E}_T^n . Suppose f is onto. Then
(i) $f^\circ(\text{Ball}(p, r)) = \text{Ball}(f(p), r)$, and
(ii) $f^\circ(\overline{\text{Ball}}(p, r)) = \overline{\text{Ball}}(f(p), r)$, and
(iii) $f^\circ(\text{Sphere}(p, r)) = \text{Sphere}((f(p)), r)$.

PROOF: $f^\circ(\text{Ball}(p, r)) = \text{Ball}(f(p), r)$ by [28, (7)]. $f^\circ(\overline{\text{Ball}}(p, r)) = \overline{\text{Ball}}(f(p), r)$ by [28, (8)]. $f^\circ(\text{Sphere}(p, r)) \subseteq \text{Sphere}((f(p)), r)$ by [28, (9)]. Consider x being an object such that $x \in \text{dom } f$ and $f(x) = y$. \square

- (19) Let us consider points p, q of \mathcal{E}_T^{n+1} , r , and s . Suppose

- (i) $s \leq r \leq |p - q|$, and
(ii) $s < |p - q| < s + r$.

Then there exists a function h from $\mathcal{E}_T^{n+1} \upharpoonright (\text{Sphere}(p, r) \cap \overline{\text{Ball}}(q, s))$ into $\text{Tdisk}(0_{\mathcal{E}_T^n}, 1)$ such that

- (iii) h is a homeomorphism, and
(iv) $h^\circ(\text{Sphere}(p, r) \cap \text{Sphere}(q, s)) = \text{Sphere}(0_{\mathcal{E}_T^n}, 1)$.

PROOF: Set $n_1 = n + 1$. Set $T_6 = \mathcal{E}_T^{n_1}$. Set $y = \frac{1}{r} \cdot (q - p)$. Set $Y = \underbrace{\langle 0, \dots, 0 \rangle}_{n_1} + \cdot (n_1, |y|)$. There exists a homogeneous additive rotation function R from T_6 into T_6 such that R is a homeomorphism and $R(y) = Y$ by [34, (40), (41)]. Consider R being a homogeneous additive rotation function from T_6 into T_6 such that R is a homeomorphism and $R(y) = Y$. $s > 0$. \square

REFERENCES

- [1] Grzegorz Bancerek. Cardinal numbers. *Formalized Mathematics*, 1(2):377–382, 1990.
- [2] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [3] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [4] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [5] Leszek Borys. Paracompact and metrizable spaces. *Formalized Mathematics*, 2(4):481–485, 1991.
- [6] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(3):529–536, 1990.
- [7] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [8] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [9] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [10] Czesław Byliński. Introduction to real linear topological spaces. *Formalized Mathematics*, 13(1):99–107, 2005.
- [11] Czesław Byliński. The sum and product of finite sequences of real numbers. *Formalized Mathematics*, 1(4):661–668, 1990.
- [12] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [13] Czesław Byliński and Piotr Rudnicki. Bounding boxes for compact sets in \mathcal{E}^2 . *Formalized Mathematics*, 6(3):427–440, 1997.
- [14] Agata Darmochwał. The Euclidean space. *Formalized Mathematics*, 2(4):599–603, 1991.
- [15] Agata Darmochwał. Finite sets. *Formalized Mathematics*, 1(1):165–167, 1990.
- [16] Agata Darmochwał. Families of subsets, subspaces and mappings in topological spaces. *Formalized Mathematics*, 1(2):257–261, 1990.
- [17] Agata Darmochwał and Yatsuka Nakamura. Metric spaces as topological spaces – fundamental concepts. *Formalized Mathematics*, 2(4):605–608, 1991.
- [18] Roman Duda. *Wprowadzenie do topologii*. PWN, 1986.
- [19] Noboru Endou and Yasunari Shidama. Completeness of the real Euclidean space. *Formalized Mathematics*, 13(4):577–580, 2005.
- [20] Ryszard Engelking. *Dimension Theory*. North-Holland, Amsterdam, 1978.
- [21] Ryszard Engelking. *General Topology*. Heldermann Verlag, Berlin, 1989.
- [22] Zbigniew Karno. Continuity of mappings over the union of subspaces. *Formalized Mathematics*, 3(1):1–16, 1992.
- [23] Zbigniew Karno. Separated and weakly separated subspaces of topological spaces. *Formalized Mathematics*, 2(5):665–674, 1991.
- [24] Artur Korniłowicz. Homeomorphism between $[\mathcal{E}_T^i, \mathcal{E}_T^j]$ and \mathcal{E}_T^{i+j} . *Formalized Mathematics*, 8(1):73–76, 1999.
- [25] Artur Korniłowicz. On the continuity of some functions. *Formalized Mathematics*, 18(3):175–183, 2010. doi:10.2478/v10037-010-0020-z.
- [26] Artur Korniłowicz. Arithmetic operations on functions from sets into functional sets. *Formalized Mathematics*, 17(1):43–60, 2009. doi:10.2478/v10037-009-0005-y.

- [27] Artur Kornilowicz and Yasunari Shidama. Brouwer fixed point theorem for disks on the plane. *Formalized Mathematics*, 13(2):333–336, 2005.
- [28] Artur Kornilowicz and Yasunari Shidama. Intersections of intervals and balls in \mathcal{E}_T^n . *Formalized Mathematics*, 12(3):301–306, 2004.
- [29] Artur Kornilowicz and Yasunari Shidama. Some properties of circles on the plane. *Formalized Mathematics*, 13(1):117–124, 2005.
- [30] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Formalized Mathematics*, 1(2):335–342, 1990.
- [31] Roman Matuszewski and Yatsuka Nakamura. Projections in n -dimensional Euclidean space to each coordinates. *Formalized Mathematics*, 6(4):505–509, 1997.
- [32] Robert Milewski. Bases of continuous lattices. *Formalized Mathematics*, 7(2):285–294, 1998.
- [33] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(1):223–230, 1990.
- [34] Karol Pałk. The rotation group. *Formalized Mathematics*, 20(1):23–29, 2012. doi:10.2478/v10037-012-0004-2.
- [35] Karol Pałk. Small inductive dimension of topological spaces. *Formalized Mathematics*, 17(3):207–212, 2009. doi:10.2478/v10037-009-0025-7.
- [36] Karol Pałk. Small inductive dimension of topological spaces. Part II. *Formalized Mathematics*, 17(3):219–222, 2009. doi:10.2478/v10037-009-0027-5.
- [37] Andrzej Trybulec. Binary operations applied to functions. *Formalized Mathematics*, 1(2):329–334, 1990.
- [38] Andrzej Trybulec. On the sets inhabited by numbers. *Formalized Mathematics*, 11(4):341–347, 2003.
- [39] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers. *Formalized Mathematics*, 1(3):445–449, 1990.
- [40] Michał J. Trybulec. Integers. *Formalized Mathematics*, 1(3):501–505, 1990.
- [41] Wojciech A. Trybulec. Non-contiguous substrings and one-to-one finite sequences. *Formalized Mathematics*, 1(3):569–573, 1990.
- [42] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(2):291–296, 1990.
- [43] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [44] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.
- [45] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.
- [46] Mirosław Wysocki and Agata Darmochwał. Subsets of topological spaces. *Formalized Mathematics*, 1(1):231–237, 1990.
- [47] Mariusz Żynel and Adam Guzowski. T_0 topological spaces. *Formalized Mathematics*, 5(1):75–77, 1996.

Received February 11, 2014
