

## F. Riesz Theorem

Keiko Narita  
Hirosaki-city  
Aomori, Japan

Kazuhisa Nakasho  
Akita Prefectural University  
Akita, Japan

Yasunari Shidama  
Shinshu University  
Nagano, Japan

**Summary.** In this article, we formalize in the Mizar system [1, 4] the F. Riesz theorem. In the first section, we defined Mizar functor `ClstoCmp`, compact topological spaces as closed interval subset of real numbers. Then using the former definition and referring to the article [10] and the article [5], we defined the normed spaces of continuous functions on closed interval subset of real numbers, and defined the normed spaces of bounded functions on closed interval subset of real numbers. We also proved some related properties.

In Sec.2, we proved some lemmas for the proof of F. Riesz theorem. In Sec.3, we proved F. Riesz theorem, about the dual space of the space of continuous functions on closed interval subset of real numbers, finally. We applied Hahn-Banach theorem (36) in [7], to the proof of the last theorem. For the description of theorems of this section, we also referred to the article [8] and the article [6]. These formalizations are based on [2], [3], [9], and [11].

MSC: 46E15 46B10 03B35

Keywords: F. Riesz theorem; dual spaces; continuous functions

MML identifier: DUALSP05, version: 8.1.06 5.44.1305

### 1. THE NORMED SPACE OF CONTINUOUS FUNCTIONS ON CLOSED INTERVAL

Now we state the propositions:

- (1) Let us consider a real number  $d$ . Then  $|\operatorname{sgn} d| \leq 1$ .
- (2) Let us consider a real number  $x$ . Then  $|x| = \operatorname{sgn} x \cdot x$ .

Let  $A$  be a non empty, closed interval subset of  $\mathbb{R}$ . The functor  $\text{Cls2Cmp}(A)$  yielding a strict, compact, non empty topological space is defined by

(Def. 1) there exist real numbers  $a, b$  such that  $a \leq b$  and  $[a, b] = A$  and  $it = [a, b]_T$ .

Now we state the propositions:

- (3) Let us consider a strict, non empty subspace  $X$  of  $\mathbb{R}^1$ , a real map  $f$  of  $X$ , a partial function  $g$  from  $\mathbb{R}$  to  $\mathbb{R}$ , a point  $x$  of  $X$ , and a real number  $x_0$ . Suppose  $g = f$  and  $x = x_0$ . Then for every subset  $V$  of  $\mathbb{R}$  such that  $f(x) \in V$  and  $V$  is open there exists a subset  $W$  of  $X$  such that  $x \in W$  and  $W$  is open and  $f^\circ W \subseteq V$  if and only if  $g$  is continuous in  $x_0$ .
- (4) Let us consider a strict, non empty subspace  $X$  of  $\mathbb{R}^1$ , a real map  $f$  of  $X$ , and a partial function  $g$  from  $\mathbb{R}$  to  $\mathbb{R}$ . If  $g = f$ , then  $f$  is continuous iff  $g$  is continuous. The theorem is a consequence of (3).
- (5) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ . Then the carrier of  $\text{Cls2Cmp}(A) = A$ .
- (6) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , and a function  $u$ . Then  $u$  is a point of  $C(\text{Cls2Cmp}(A); \mathbb{R})$  if and only if  $\text{dom } u = A$  and  $u$  is a continuous partial function from  $\mathbb{R}$  to  $\mathbb{R}$ . The theorem is a consequence of (5) and (4).
- (7) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , and a point  $v$  of  $C(\text{Cls2Cmp}(A); \mathbb{R})$ . Then  $v \in \text{BoundedFunctions}(\text{the carrier of } \text{Cls2Cmp}(A))$ .

## 2. PRELIMINARIES

Now we state the proposition:

- (8) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , and real numbers  $a, b$ . Suppose  $A = [a, b]$ . Then there exists a function  $x$  from  $A$  into  $\text{BoundedFunctions } A$  such that for every real number  $s$  such that  $s \in [a, b]$  holds if  $s = a$ , then  $x(s) = [a, b] \mapsto 0$  and if  $s \neq a$ , then  $x(s) = ([a, s] \mapsto 1) + ([s, b] \mapsto 0)$ .

PROOF: Define  $\mathcal{C}[\text{object}] \equiv \$1 = a$ . Define  $\mathcal{F}(\text{object}) = [a, b] \mapsto 0$ . Define  $\mathcal{G}(\text{object}) = ([a, \$1 \in \mathbb{R}] \mapsto 1) + ([\$1 \in \mathbb{R}], b] \mapsto 0)$ . Set  $B = \text{BoundedFunctions } A$ . For every object  $s$  such that  $s \in [a, b]$  holds if  $\mathcal{C}[s]$ , then  $\mathcal{F}(s) \in B$  and if  $\mathcal{C}[s]$ , then  $\mathcal{G}(s) \in B$ . Consider  $x$  being a function from  $[a, b]$  into  $B$  such that for every object  $s$  such that  $s \in [a, b]$  holds if  $\mathcal{C}[s]$ , then  $x(s) = \mathcal{F}(s)$  and if  $\mathcal{C}[s]$ , then  $x(s) = \mathcal{G}(s)$ . For every real number  $s$  such that  $s \in [a, b]$  holds if  $s = a$ , then  $x(s) = [a, b] \mapsto 0$  and if  $s \neq a$ , then  $x(s) = ([a, s] \mapsto 1) + ([s, b] \mapsto 0)$ .  $\square$

Let  $A$  be a non empty, closed interval subset of  $\mathbb{R}$ ,  $D$  be a partition of  $A$ ,  $m$  be a function from  $A$  into BoundedFunctions  $A$ , and  $i$  be a natural number. Assume  $i \in \text{Seg}(\text{len } D + 1)$ . The functor  $\text{Dp1}(m, D, i)$  yielding a point of the  $\mathbb{R}$ -normed algebra of bounded functions on the carrier of  $\text{Cls2Cmp}(A)$  is defined by the term

$$\text{(Def. 2)} \quad \begin{cases} m(\inf A), & \text{if } i = 1, \\ m(D(i - 1)), & \text{otherwise.} \end{cases}$$

Let  $\varrho$  be a function from  $A$  into  $\mathbb{R}$ . The functor  $\text{Dp2}(\varrho, D, i)$  yielding a real number is defined by the term

$$\text{(Def. 3)} \quad \begin{cases} \varrho(\inf A), & \text{if } i = 1, \\ \varrho(D(i - 1)), & \text{otherwise.} \end{cases}$$

Now we state the propositions:

(9) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , a partition  $D$  of  $A$ , a function  $m$  from  $A$  into BoundedFunctions  $A$ , and a function  $\varrho$  from  $A$  into  $\mathbb{R}$ . Then there exists a finite sequence  $s$  of elements of the  $\mathbb{R}$ -normed algebra of bounded functions on the carrier of  $\text{Cls2Cmp}(A)$  such that

- (i)  $\text{len } s = \text{len } D$ , and
- (ii) for every natural number  $i$  such that  $i \in \text{dom } s$  holds  $s(i) = \text{sgn}(\text{Dp2}(\varrho, D, i + 1) - \text{Dp2}(\varrho, D, i)) \cdot (\text{Dp1}(m, D, i + 1) - \text{Dp1}(m, D, i))$ .

PROOF: Set  $V =$  the  $\mathbb{R}$ -normed algebra of bounded functions on the carrier of  $\text{Cls2Cmp}(A)$ . Define  $\mathcal{P}[\text{natural number, set}] \equiv \$_2 = \text{sgn}(\text{Dp2}(\varrho, D, \$_1 + 1) - \text{Dp2}(\varrho, D, \$_1)) \cdot (\text{Dp1}(m, D, \$_1 + 1) - \text{Dp1}(m, D, \$_1))$ . Consider  $s$  being a finite sequence of elements of  $V$  such that  $\text{dom } s = \text{Seg len } D$  and for every natural number  $i$  such that  $i \in \text{Seg len } D$  holds  $\mathcal{P}[i, s(i)]$ .  $\square$

(10) Let us consider a real linear space  $V$ , a functional  $f$  in  $V$ , and a finite sequence  $s$  of elements of  $V$ . If  $f$  is additive, then  $f(\sum s) = \sum(f \cdot s)$ .

PROOF: Define  $\mathcal{P}[\text{natural number}] \equiv$  for every real linear space  $V$  for every functional  $f$  in  $V$  for every finite sequence  $s$  of elements of  $V$  such that  $\text{len } s = \$_1$  and  $f$  is additive holds  $f(\sum s) = \sum(f \cdot s)$ .  $\mathcal{P}[0]$ . For every natural number  $n$  such that  $\mathcal{P}[n]$  holds  $\mathcal{P}[n + 1]$ . For every natural number  $n$ ,  $\mathcal{P}[n]$ .  $\square$

(11) Let us consider a non empty set  $A$ . Then every element of the  $\mathbb{R}$ -normed algebra of bounded functions on  $A$  is a function from  $A$  into  $\mathbb{R}$ .

(12) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , a finite sequence  $s$  of elements of the  $\mathbb{R}$ -normed algebra of bounded functions on the carrier of  $\text{Cls2Cmp}(A)$ , a finite sequence  $z$  of elements of  $\mathbb{R}$ , a function  $g$  from  $A$  into  $\mathbb{R}$ , and an element  $t$  of  $A$ . Suppose  $\text{len } s = \text{len } z$  and  $g = \sum s$

and for every natural number  $k$  such that  $k \in \text{dom } z$  there exists a function  $s_1$  from  $A$  into  $\mathbb{R}$  such that  $s_1 = s(k)$  and  $z(k) = s_1(t)$ . Then  $g(t) = \sum z$ .  
 PROOF: Define  $\mathcal{P}[\text{natural number}] \equiv$  for every non empty, closed interval subset  $A$  of  $\mathbb{R}$  for every finite sequence  $s$  of elements of the  $\mathbb{R}$ -normed algebra of bounded functions on the carrier of  $\text{Cls2Cmp}(A)$  for every finite sequence  $z$  of elements of  $\mathbb{R}$  for every function  $g$  from  $A$  into  $\mathbb{R}$  for every element  $t$  of  $A$  such that  $\text{len } s = \$_1$  and  $\text{len } s = \text{len } z$  and  $g = \sum s$  and for every natural number  $k$  such that  $k \in \text{dom } z$  there exists a function  $s_1$  from  $A$  into  $\mathbb{R}$  such that  $s_1 = s(k)$  and  $z(k) = s_1(t)$  holds  $g(t) = \sum z$ .  
 $\mathcal{P}[0]$ . For every natural number  $n$  such that  $\mathcal{P}[n]$  holds  $\mathcal{P}[n + 1]$ . For every natural number  $n$ ,  $\mathcal{P}[n]$ .  $\square$

- (13) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , a partition  $D$  of  $A$ , and an element  $t$  of  $A$ . Suppose  $\text{inf } A < D(1)$ . Then there exists an element  $i$  of  $\mathbb{N}$  such that
  - (i)  $i \in \text{dom } D$ , and
  - (ii)  $t \in \text{divset}(D, i)$ , and
  - (iii)  $i = 1$  or  $\text{inf divset}(D, i) < t \leq \text{sup divset}(D, i)$ .
- (14) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , a function  $\varrho$  from  $A$  into  $\mathbb{R}$ , and a real number  $B$ . Suppose  $0 < \text{vol}(A)$ . Suppose for every partition  $D$  of  $A$  and for every var-volume  $K$  of  $\varrho$  and  $D$  such that  $\text{inf } A < D(1)$  holds  $\sum K \leq B$ . Let us consider a partition  $D$  of  $A$ , and a var-volume  $K$  of  $\varrho$  and  $D$ . Then  $\sum K \leq B$ .

### 3. F. RIESZ THEOREM

Now we state the propositions:

- (15) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , a function  $\varrho$  from  $A$  into  $\mathbb{R}$ , and a point  $f$  of  $\text{DualSp } C(\text{Cls2Cmp}(A); \mathbb{R})$ . Suppose  $\varrho$  is bounded-variation and for every continuous partial function  $u$  from  $\mathbb{R}$  to  $\mathbb{R}$  such that  $\text{dom } u = A$  holds  $f(u) = \int_{\varrho} u(x)dx$ . Then  $\|f\| \leq \text{TotalVD}(\varrho)$ .  
 PROOF: Set  $X = C(\text{Cls2Cmp}(A); \mathbb{R})$ . For every continuous partial function  $u$  from  $\mathbb{R}$  to  $\mathbb{R}$  such that  $u \in$  the carrier of  $X$  holds  $f(u) = \int_{\varrho} u(x)dx$ . For every continuous partial function  $u$  from  $\mathbb{R}$  to  $\mathbb{R}$  and for every point  $v$  of  $X$  such that  $\text{dom } u = A$  and  $u = v$  holds  $|\int_{\varrho} u(x)dx| \leq \|v\| \cdot \text{TotalVD}(\varrho)$ .  
 $\square$

(16) Let us consider a non empty, closed interval subset  $A$  of  $\mathbb{R}$ , and a point  $x$  of  $\text{DualSp } C(\text{Cls2Cmp}(A); \mathbb{R})$ . Suppose  $0 < \text{vol}(A)$ . Then there exists a function  $\varrho$  from  $A$  into  $\mathbb{R}$  such that

- (i)  $\varrho$  is bounded-variation, and
- (ii) for every continuous partial function  $u$  from  $\mathbb{R}$  to  $\mathbb{R}$  such that  $\text{dom } u = A$  holds  $x(u) = \int_{\varrho} u(x)dx$ , and
- (iii)  $\|x\| = \text{TotalVD}(\varrho)$ .

PROOF: Set  $X = C(\text{Cls2Cmp}(A); \mathbb{R})$ . Set  $V =$  the  $\mathbb{R}$ -normed algebra of bounded functions on the carrier of  $\text{Cls2Cmp}(A)$ . Set  $A_1 =$  the carrier of  $\text{Cls2Cmp}(A)$ .  $A_1 = A$ . Reconsider  $h = x$  as a Lipschitzian linear functional in  $X$ . Consider  $f$  being a Lipschitzian linear functional in  $V$ ,  $F$  being a point of  $\text{DualSp } V$  such that  $f = F$  and  $f \upharpoonright (\text{the carrier of } X) = h$  and  $\|F\| = \|x\|$ . Consider  $a, b$  being real numbers such that  $a \leq b$  and  $[a, b] = A$  and  $\text{Cls2Cmp}(A) = [a, b]_{\mathbb{T}}$ . Consider  $m$  being a function from  $A$  into  $\text{BoundedFunctions } A$  such that for every real number  $s$  such that  $s \in [a, b]$  holds if  $s = a$ , then  $m(s) = [a, b] \mapsto 0$  and if  $s \neq a$ , then  $m(s) = ([a, s] \mapsto 1) + ([s, b] \mapsto 0)$ . The carrier of  $V = \text{BoundedFunctions } A$ . Reconsider  $\varrho = f \cdot m$  as a function from  $A$  into  $\mathbb{R}$ . For every partition  $D$  of  $A$  and for every var-volume  $K$  of  $\varrho$  and  $D$  such that  $a < D(1)$  holds  $\sum K \leq \|x\|$ . For every partition  $D$  of  $A$  and for every var-volume  $K$  of  $\varrho$  and  $D$ ,  $\sum K \leq \|x\|$ . Consider  $V_1$  being a non empty subset of  $\mathbb{R}$  such that  $V_1$  is upper bounded and  $V_1 = \{r, \text{ where } r \text{ is a real number} : \text{ there exists a partition } t \text{ of } A \text{ and there exists a var-volume } F_0 \text{ of } \varrho \text{ and } t \text{ such that } r = \sum F_0\}$  and  $\text{TotalVD}(\varrho) = \sup V_1$ . For every continuous partial function  $u$  from  $\mathbb{R}$  to  $\mathbb{R}$  such that  $\text{dom } u = A$  holds  $x(u) = \int_{\varrho} u(x)dx$ .  $\|x\| \leq \text{TotalVD}(\varrho)$ .  $\square$

REFERENCES

- [1] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Kornilowicz, Roman Matuszewski, Adam Naumowicz, Karol Pąk, and Josef Urban. Mizar: State-of-the-art and beyond. In Manfred Kerber, Jacques Carette, Cezary Kaliszyk, Florian Rabe, and Volker Sorge, editors, *Intelligent Computer Mathematics*, volume 9150 of *Lecture Notes in Computer Science*, pages 261–279. Springer International Publishing, 2015. ISBN 978-3-319-20614-1. doi:10.1007/978-3-319-20615-8\_17.
- [2] Haim Brezis. *Functional Analysis, Sobolev Spaces and Partial Differential Equations*. Springer, 2011.
- [3] Peter D. Dax. *Functional Analysis*. Pure and Applied Mathematics: A Wiley Series of Texts, Monographs and Tracts. Wiley Interscience, 2002.
- [4] Adam Grabowski, Artur Kornilowicz, and Adam Naumowicz. Four decades of Mizar. *Journal of Automated Reasoning*, 55(3):191–198, 2015. doi:10.1007/s10817-015-9345-1.

- [5] Katuhiko Kanazashi, Noboru Endou, and Yasunari Shidama. Banach algebra of continuous functionals and the space of real-valued continuous functionals with bounded support. *Formalized Mathematics*, 18(1):11–16, 2010. doi:10.2478/v10037-010-0002-1.
- [6] Kazuhisa Nakasho, Keiko Narita, and Yasunari Shidama. The basic existence theorem of Riemann-Stieltjes integral. *Formalized Mathematics*, 24(4):253–259, 2016. doi:10.1515/forma-2016-0021.
- [7] Keiko Narita, Noboru Endou, and Yasunari Shidama. Dual spaces and Hahn-Banach theorem. *Formalized Mathematics*, 22(1):69–77, 2014. doi:10.2478/forma-2014-0007.
- [8] Keiko Narita, Kazuhisa Nakasho, and Yasunari Shidama. Riemann-Stieltjes integral. *Formalized Mathematics*, 24(3):199–204, 2016. doi:10.1515/forma-2016-0016.
- [9] Walter Rudin. *Functional Analysis*. New York, McGraw-Hill, 2nd edition, 1991.
- [10] Yasunari Shidama, Hikofumi Suzuki, and Noboru Endou. Banach algebra of bounded functionals. *Formalized Mathematics*, 16(2):115–122, 2008. doi:10.2478/v10037-008-0017-z.
- [11] Kosaku Yoshida. *Functional Analysis*. Springer, 1980.

*Received August 30, 2017*

---



The English version of this volume of Formalized Mathematics was financed under agreement 548/P-DUN/2016 with the funds from the Polish Minister of Science and Higher Education for the dissemination of science.